




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification⁶ : C07D 295/22, C07C 311/46, C07D 213/42, 333/20, C07C 317/34, 323/49, 311/51, 311/58, C07D 211/46, 211/96, C07C 311/60, A61K 31/18</p>	<p>A1</p>	<p>(11) International Publication Number: WO 99/47508 (43) International Publication Date: 23 September 1999 (23.09.99)</p>
<p>(21) International Application Number: PCT/GB99/00739 (22) International Filing Date: 11 March 1999 (11.03.99) (30) Priority Data: 9805520.5 17 March 1998 (17.03.98) GB (71) Applicant (for all designated States except US): ZENECA LIMITED [GB/GB]; 15 Stanhope Gate, London W1Y 6LN (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): BUTLIN, Roger, John [GB/GB]; Alderley Park, Macclesfield, Cheshire SK10 4TG (GB). BURROWS, Jeremy, Nicholas [GB/GB]; Alderley Park, Macclesfield, Cheshire SK10 4TG (GB). CLARKE, David, Stephen [GB/GB]; Alderley Park, Macclesfield, Cheshire SK10 4TG (GB). NOWAK, Thorsten [DE/GB]; Alderley Park, Macclesfield, Cheshire SK10 4TG (GB). WHITTAMORE, Paul, Robert, Owen [GB/GB]; Alderley Park, Macclesfield, Cheshire SK10 4TG (GB).</p>	<p>(74) Agent: BILL, Kevin; Zeneca Pharmaceuticals, Intellectual Property Dept., Mereside, Alderley Park, Macclesfield, Cheshire SK10 4TG (GB). (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</p>	

(54) Title: BENZENESULFONAMIDE-DERIVATIVES AND THEIR USE AS MEDICAMENTS



(57) Abstract

Compounds of formula (I), pharmaceutically acceptable salts or *in vivo* hydrolysable esters thereof, wherein: Ring X is phenyl or a six membered heteroaryl ring containing one or two ring nitrogens where said nitrogens are optionally oxidised to form the N-oxide; R^1 and R^2 are substituents as defined within; R^3 and R^4 are defined within and are alkyl or halo alkyl or together form a cycloalkyl or halocycloalkyl ring; R^5 is a substituent as defined within; Y-Z is a linking group as defined within; are useful in the production of an elevation of PDH activity in a warm-blooded animal such as a human being. Pharmaceutical compositions, methods and processes for preparation of compounds of formula (I) are described.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

BENZENESULFONAMIDE-DERIVATIVES AND THEIR USE AS MEDICAMENTS

The present invention relates to compounds which elevate pyruvate dehydrogenase (PDH) activity; processes for their preparation, pharmaceutical compositions containing them as active ingredient, methods for the treatment of disease states associated with reduced PDH activity, to their use as medicaments and to their use in the manufacture of medicaments for use in the elevation of PDH activity in warm-blooded animals such as humans.

Within tissues adenosine triphosphate (ATP) provides the energy for synthesis of complex molecules and, in muscle, for contraction. ATP is generated from the breakdown of energy-rich substrates such as glucose or long chain free fatty acids. In oxidative tissues such as muscle the majority of the ATP is generated from acetyl CoA which enters the citric acid cycle, thus the supply of acetyl CoA is a critical determinant of ATP production in oxidative tissues. Acetyl CoA is produced either by β -oxidation of fatty acids or as a result of glucose metabolism by the glycolytic pathway. The key regulatory enzyme in controlling the rate of acetyl CoA formation from glucose is PDH which catalyses the oxidation of pyruvate to acetyl CoA and carbon dioxide with concomitant reduction of nicotinamide adenine dinucleotide (NAD) to NADH.

In disease states such as both non-insulin dependent (NIDDM) and insulin-dependent diabetes mellitus (IDDM), oxidation of lipids is increased with a concomitant reduction in utilisation of glucose, which contributes to the hyperglycaemia. Reduced glucose utilisation in both IDDM and NIDDM is associated with a reduction in PDH activity. In addition, a further consequence of reduced PDH activity may be that an increase in pyruvate concentration results in increased availability of lactate as a substrate for hepatic gluconeogenesis. It is reasonable to expect that increasing the activity of PDH could increase the rate of glucose oxidation and hence overall glucose utilisation, in addition to reducing hepatic glucose output. Another factor contributing to diabetes mellitus is impaired insulin secretion, which has been shown to be associated with reduced PDH activity in pancreatic β -cells (in a rodent genetic model of diabetes mellitus Zhou et al. (1996) Diabetes 45: 580-586).

Oxidation of glucose is capable of yielding more molecules of ATP per mole of oxygen than is oxidation of fatty acids. In conditions where energy demand may exceed energy supply, such as myocardial ischaemia, intermittent claudication, cerebral ischaemia and reperfusion,

(Zaidan et al., 1998; J. Neurochem. 70: 233-241), shifting the balance of substrate utilisation in favour of glucose metabolism by elevating PDH activity may be expected to improve the ability to maintain ATP levels and hence function.

An agent which is capable of elevating PDH activity may also be expected to be of benefit in treating conditions where an excess of circulating lactic acid is manifest such as in certain cases of sepsis.


The agent dichloroacetic acid (DCA) which increases the activity of PDH after acute administration in animals, (Vary et al., 1988; Circ. Shock, 24: 3-18), has been shown to have the predicted effects in reducing glycaemia, (Stacpoole et al., 1978; N. Engl. J. Med. 298: 526-530), and as a therapy for myocardial ischaemia (Bersin and Stacpoole 1997; American Heart Journal, 134: 841-855) and lactic acidemia, (Stacpoole et al., 1983; N. Engl. J. Med. 309: 390-396).

PDH is an intramitochondrial multienzyme complex consisting of multiple copies of several subunits including three enzyme activities E1, E2 and E3, required for the completion of the conversion of pyruvate to acetyl CoA (Patel and Roche 1990; FASEB J., 4: 3224-3233). E1 catalyses the non-reversible removal of CO₂ from pyruvate; E2 forms acetyl CoA and E3 reduces NAD to NADH. Two additional enzyme activities are associated with the complex: a specific kinase which is capable of phosphorylating E1 at three serine residues and a loosely-associated specific phosphatase which reverses the phosphorylation. Phosphorylation of a single one of the three serine residues renders the E1 inactive. The proportion of the PDH in its active (dephosphorylated) state is determined by a balance between the activity of the kinase and phosphatase. The activity of the kinase may be regulated in vivo by the relative concentrations of metabolic substrates such as NAD/NADH, CoA/acetylCoA and adenine diphosphate (ADP)/ATP as well as by the availability of pyruvate itself.

European Patent Publication No. 625516 refers to compounds which are capable of relaxing bladder smooth muscle and which may be used in the treatment of urge incontinence. We have found, surprisingly, that compounds also containing a sulphonamide moiety disclosed in the present invention are very good at elevating PDH activity, a property nowhere disclosed in EP 625516. The present invention is based on the surprising discovery that certain compounds elevate PDH activity, a property of value in the treatment of disease states associated with disorders of glucose utilisation such as diabetes mellitus, obesity, (Curto et al., 1997; Int. J. Obes. 21:

1137-1142), and lactic acidemia. Additionally the compounds may be expected to have utility in diseases where supply of energy-rich substrates to tissues is limiting such as peripheral vascular disease; (including intermittent claudication), cardiac failure and certain cardiac myopathies, muscle weakness, hyperlipidaemias and atherosclerosis (Stacpoole et al., 1978; N. Engl. J. Med. 5 298: 526-530). A compound that activates PDH may also be useful in treating Alzheimer disease (AD) (J Neural Transm (1998) 105: 855-870).

Accordingly, the present invention provides a compound of formula (I):



(I)

10 wherein:

Ring X is phenyl or a six membered heteroaryl ring containing one or two ring nitrogens where said nitrogens are optionally oxidised to form the *N*-oxide;

R¹ and **R²** are independently as defined in (a) or (b);

R³ and **R⁴** are as defined in (c) or (d);

15 **R⁵** is as defined in (e) or (f);

Y-Z is as defined in (g) or (h)

wherein:

(a) either **R¹** and **R²** are each selected independently from hydrogen, C₁₋₃alkyl, pyridyl and phenyl which is unsubstituted or substituted by one or two substituents selected independently

20 from C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkenyloxy, hydroxy, halo and cyano,

or **R¹** and **R²** together with the nitrogen atom to which they are attached form morpholino, thiomorpholino, piperidinyl, pyrrolidinyl or imidazolyl;

(b) either **R¹** and **R²** are each selected independently from phenyl substituted by one or more P (wherein P is as defined hereinbelow), phenyl substituted by one or more groups selected

25 from C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkenyloxy, hydroxy, halo and cyano and additionally substituted by one or more groups selected from P, a heterocyclic group other than unsubstituted pyridyl which is optionally substituted on a ring carbon by one or more Q (wherein Q is as defined

hereinbelow) and wherein if said heterocyclic group contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from D (wherein D is as defined hereinbelow), naphthyl optionally substituted by one or more Q, C₄₋₆alkyl, C₃₋₆cycloalkyl optionally substituted with one or more Q, C₂₋₆alkenyl, C₂₋₆alkynyl, C₁₋₆alkyl substituted by one or more V (wherein V is as defined hereinbelow), R⁶T- (wherein R⁶ and T are as defined hereinbelow) and R⁷C₁₋₆alkylT- (where R⁷ is as defined hereinbelow),

or R¹ and R² together with the nitrogen atom to which they are attached form a heterocyclic group other than unsubstituted morpholino, unsubstituted thiomorpholino, unsubstituted piperidinyl, unsubstituted pyrrolidinyl or unsubstituted imidazolyl which is optionally substituted on a ring carbon by one or more Q (wherein Q is as defined hereinbelow) and wherein if said heterocyclic group contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from D (wherein D is as defined hereinbelow);

(c) either R³ and R⁴ are independently C_kalkyl optionally substituted by from 1 to 2k+1 atoms selected from fluoro and chloro wherein k is 1-3, provided that R³ and R⁴ are not both methyl;

or R³ and R⁴ together with the carbon atom to which they are attached, form a C_mcycloalkyl ring optionally substituted by from 1 to 2m-2 fluorine atoms wherein m is 3-5;

(d) R³ and R⁴ are both methyl;

(e) R⁵ is hydrogen, C₁₋₄alkyl, haloC₁₋₄alkyl, C₁₋₄alkoxy, haloC₁₋₄alkoxy, cyano, nitro, C₂₋₄alkenyloxy or trifluoromethylthio;

(f) R⁵ is halo, hydroxy, amino, C₁₋₆alkylamino, (C₁₋₆alkyl)₂amino, C₁₋₆alkanoylamino, C₁₋₆alkanoyl(N-C₁₋₆alkyl)amino, C₁₋₆alkylsulphonylamino, C₁₋₆alkylsulphonyl(N-C₁₋₆alkyl)amino, thiol, C₁₋₆alkylsulphanyl, C₁₋₆alkylsulphiny, C₁₋₆alkylsulphonyl, sulphamoyl, N-(C₁₋₆alkyl)aminosulphonyl, N-(C₁₋₆alkyl)₂aminosulphonyl, carboxy, carbamoyl, N-(C₁₋₆alkyl)carbamoyl, N-(C₁₋₆alkyl)₂carbamoyl, C₁₋₆alkoxycarbonyl, formyl, C₁₋₆alkanoyl, C₂₋₆alkenyl, C₂₋₆alkynyl, C₃₋₆alkyl, haloC₃₋₆alkyl, C₃₋₆alkoxy, haloC₃₋₆alkoxy or C₃₋₆alkenyloxy;

(g) Y-Z is -NHC(O)-, -OCH₂-, -SCH₂-, -NHCH₂-, trans-vinylene, and ethynylene;

(h) Y-Z is -NHC(S)-;

R⁶ is selected from C₁₋₆alkyl (optionally substituted with one or more R⁸), C₃₋₆cycloalkyl optionally substituted with one or more R⁸, a heterocyclic group optionally substituted on a ring

carbon by one or more R^8 and if said heterocyclic group contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from D (wherein D is as defined hereinbelow), phenyl optionally substituted by one or more R^8 , naphthyl optionally substituted by one or more R^8 ;

- 5 R^7 is a heterocyclic group optionally substituted on a ring carbon by one or more R^8 and if said heterocyclic group contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from D (wherein D is as defined hereinbelow), phenyl optionally substituted by one or more R^8 , naphthyl optionally substituted by one or more R^8 ;

R^8 is trifluoromethyl, C_{1-6} alkyl, halo, hydroxy, trifluoromethoxy, cyano, C_{1-6} alkoxy,

- 10 formyl, C_{1-6} alkanoyl, C_{1-6} alkanoyloxy, amino, C_{1-6} alkylamino, $(C_{1-6}alkyl)_2$ amino, C_{1-6} alkanoylamino, C_{1-6} alkanoyl(N - $C_{1-6}alkyl$)amino, nitro, carboxy, carbamoyl, C_{1-6} alkoxycarbonyl, thiol, C_{1-6} alkylsulphanyl, C_{1-6} alkylsulphinyl, C_{1-6} alkylsulphonyl, sulphamoyl, N -($C_{1-6}alkyl$)aminosulphonyl, N -($C_{1-6}alkyl$) $_2$ aminosulphonyl, carbamoyl $C_{1-6}alkyl$, N -($C_{1-6}alkyl$)carbamoyl $C_{1-6}alkyl$, N -($C_{1-6}alkyl$) $_2$ carbamoyl- $C_{1-6}alkyl$, hydroxy $C_{1-6}alkyl$,
15 $C_{1-6}alkoxyC_{1-6}alkyl$, phenyl $C_{1-6}alkyl$ or phenyl $C_{1-6}alkoxy$;

P is selected from - $C_{2-6}alkyl$ -M- substituted with one or more R^9 , - $C_{2-6}alkenyl$ -M- optionally substituted with one or more R^9 , - $C_{2-6}alkynyl$ -M- optionally substituted with one or more R^9 (with the proviso that in the three previous groups R^9 is not a substituent on the carbon atom attached to M), R^{10} -CH₂-M-, R^{11} -M-, thiol, $C_{1-6}alkylsulphanyl$, $C_{1-6}alkylsulphinyl$,

- 20 $C_{1-6}alkylsulphonyl$, sulphamoyl, nitro, carboxy, $C_{1-6}alkoxycarbonyl$, amino, $C_{1-6}alkylamino$, $(C_{1-6}alkyl)_2$ amino, carbamoyl, N -($C_{1-6}alkyl$)carbamoyl, N -($C_{1-6}alkyl$) $_2$ carbamoyl, $C_{1-6}alkanoylamino$, $C_{1-6}alkanoyl(N$ - $C_{1-6}alkyl$)amino, trifluoromethyl, trifluoromethoxy, formyl, $C_{1-6}alkanoyl$, $C_{5-6}alkyl$, $C_{2-6}alkenyl$, $C_{2-6}alkynyl$, N -($C_{1-6}alkyl$)aminosulphonyl, hydroxymethyl, hydroxyacetyl, N -($C_{1-6}alkyl$) $_2$ aminosulphonyl, $C_{1-6}alkanoylaminosulphonyl$,
25 $C_{1-6}alkanoyl(N$ - $C_{1-6}alkyl$)aminosulphonyl, $C_{1-6}alkylsulphonylaminocarbonyl$, $C_{1-6}alkylsulphonyl(N$ - $C_{1-6}alkyl$)aminocarbonyl, $C_{5-6}alkoxy$, $C_{5-6}alkenyloxy$, phenyl optionally substituted by one or more R^8 , naphthyl optionally substituted by one or more R^8 and a heterocyclic group optionally substituted on a ring carbon by one or more R^8 and if said heterocyclic group contains an -NH- moiety that nitrogen may be optionally substituted by a
30 group selected from D (wherein D is as defined hereinbelow);

Q is selected from any of the values defined for P; C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkenyloxy, hydroxy, halo and cyano;

V is selected from any of the values defined for Q, phenyl optionally substituted by one or more Q, naphthyl optionally substituted by one or more Q, a heterocyclic group optionally substituted on a ring carbon by one or more Q and if said heterocyclic group contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from D (wherein D is as defined hereinbelow) or C₃₋₆cycloalkyl optionally substituted with one or more Q;

T is selected from -O-, -C(O)-, -NH-, -N(N-C₁₋₆alkyl)-, -C(O)NH-, -NHC(O)-, -C(O)N(N-C₁₋₆alkyl)-, -N(N-C₁₋₆alkyl)C(O)-, -SO₂-, -C(S)-, -C(S)NH-, -NHC(S)-, -C(S)N(N-C₁₋₆alkyl)- and -N(N-C₁₋₆alkyl)C(S)-;

M is selected from -O-, -N(R¹²)-, -C(O)-, -N(R¹²)C(O)-, -C(O)N(R¹²)-, -S(O)_n-, -OC(O)-, -C(O)O-, -N(R¹²)C(O)O-, -OC(O)N(R¹²)-, -C(S)N(R¹²)-, -N(R¹²)C(S)-, -SO₂N(R¹²)-, -N(R¹²)SO₂- and -N(R¹²)C(O)N(R¹²)-, -N(R¹²)C(S)N(R¹²)-, -SO₂NHC(O)-, -SO₂N(R¹²)C(O)-, -C(O)NHSO₂-, -C(O)N(R¹²)SO₂- or M is a direct bond;

D is selected from C₁₋₆alkyl, C₁₋₆alkanoyl, C₁₋₆alkylsulphonyl, C₁₋₆alkoxycarbonyl, carbamoyl, *N*-(C₁₋₆alkyl)carbamoyl, *N,N*-(C₁₋₆alkyl)₂carbamoyl, benzoyl, (heterocyclic group)carbonyl, phenylsulphonyl, (heterocyclic group)sulphonyl, phenyl or a carbon linked heterocyclic group, and wherein any C₁₋₆alkyl group may be optionally substituted by one or more R⁹, and wherein any phenyl or heterocyclic group may be optionally substituted on a ring carbon by one or more groups selected from R⁸ and if a heterocyclic group contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from E;

E is selected from C₁₋₆alkyl, C₁₋₆alkanoyl, C₁₋₆alkylsulphonyl, C₁₋₆alkoxycarbonyl, carbamoyl, *N*-(C₁₋₆alkyl)carbamoyl, *N,N*-(C₁₋₆alkyl)₂carbamoyl, C₁₋₆alkoxyC₁₋₆alkanoyl, phenylC₁₋₆alkyl, benzoyl, phenylC₁₋₆alkanoyl, phenylC₁₋₆alkoxycarbonyl and phenylsulphonyl.

R⁹ is selected from hydroxy, amino, C₁₋₆alkylamino, (C₁₋₆alkyl)₂amino, carboxy, C₁₋₆alkoxy, carbamoyl, *N*-(C₁₋₆alkyl)carbamoyl, *N*-(C₁₋₆alkyl)₂carbamoyl, formyl, sulphamoyl, *N*-C₁₋₆alkylaminosulphonyl, *N*-(C₁₋₆alkyl)₂aminosulphonyl, C₁₋₆alkylsulphonylamino, C₁₋₆alkanoylamino, a heterocyclic group optionally substituted on a ring carbon by one or more R⁸ and if said heterocyclic group contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from D (wherein D is as defined hereinabove), phenyl optionally

substituted by one or more R^8 , naphthyl optionally substituted by one or more R^8 , C_{1-6} alkylsulphanyl, C_{1-6} alkylsulphinyl and C_{1-6} alkylsulphonyl;

R^{10} is carboxy, carbamoyl, N -(C_{1-6} alkyl)carbamoyl, N -(C_{1-6} alkyl)₂carbamoyl, sulphamoyl, N -(C_{1-6} alkyl)aminosulphonyl, N -(C_{1-6} alkyl)₂aminosulphonyl, C_{1-6} alkylsulphanyl,

- 5 C_{1-6} alkylsulphinyl, C_{1-6} alkylsulphonyl, C_{1-6} alkoxycarbonyl, C_{1-6} alkanoylamino; a heterocyclic group optionally substituted on a ring carbon by one or more R^8 and if said heterocyclic group contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from D (wherein D is as defined hereinabove), phenyl optionally substituted by one or more R^8 or naphthyl optionally substituted by one or more R^8 ;

- 10 R^{11} is a heterocyclic group optionally substituted on a ring carbon by one or more R^8 and if said heterocyclic group contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from D (wherein D is as defined hereinabove), phenyl optionally substituted by one or more R^8 , C_{3-6} cycloalkyl optionally substituted by one or more R^8 , or naphthyl optionally substituted by one or more R^8 ;

- 15 R^{12} is hydrogen or C_{1-6} alkyl optionally substituted with R^{13} with the proviso that R^{13} is not a substituent on the carbon attached to the nitrogen atom of M;

R^{13} is halo, hydroxy, amino, cyano, nitro, trifluoromethyl, trifluoromethoxy, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkylamino, (C_{1-6} alkyl)₂amino, C_{1-6} alkanoylamino, C_{1-6} alkanoyl(N - C_{1-6} alkyl)amino, C_{1-6} alkylsulphonylamino, C_{1-6} alkylsulphonyl(N - C_{1-6} alkyl)amino,

20 thiol, C_{1-6} alkylsulphanyl, C_{1-6} alkylsulphinyl, C_{1-6} alkylsulphonyl, sulphamoyl, N -(C_{1-6} alkyl)aminosulphonyl, N -(C_{1-6} alkyl)₂aminosulphonyl, carboxy, carbamoyl, N -(C_{1-6} alkyl)carbamoyl, N -(C_{1-6} alkyl)₂carbamoyl, C_{1-6} alkoxycarbonyl, C_{1-6} alkanoyl or formyl;

n is 0-2;

with the proviso that where R^1 and R^2 are both as defined in (a), R^3 and R^4 are both as defined in

- 25 (c), R^5 is as defined in (e) and Ring X is phenyl, Y-Z must be -NHC(S)-;

and pharmaceutically acceptable salts or *in vivo* hydrolysable esters thereof,

and with the proviso that when R^3 and R^4 are both methyl, R^5 is hydrogen, fluoro or chloro, Y-Z is ethynylene, X is phenyl and one of R^1 and R^2 is hydrogen and the other is pyrimidyl-NH-C(O)- or triazinyl-NH-C(O)- (wherein said triazine or pyrimidine is substituted by methyl, methoxy or

dimethylamino) then the $-\text{SO}_2\text{NR}^1\text{R}^2$ moiety cannot be ortho to Y-Z; and provided said compound is not:

4-(3-hydroxy-3-methyl-1-butynyl)-N-(3-methyl-2-pyridinyl)-benzenesulphonamide,

N-{4-[N,N-bis-(*sec*-butyl)aminosulphonyl]phenyl}-2-hydroxy-2-methyl-3,3,3-

5 trifluoropropanamide, or

N-{4-[N,N-bis-(*iso*-butyl)aminosulphonyl]phenyl}-2-hydroxy-2-methyl-3,3,3-

trifluoropropanamide.

In this specification the term "alkyl" includes both straight and branched chain alkyl groups but references to individual alkyl groups such as "propyl" are specific for the straight
10 chain version only. For example, " C_{1-4} alkyl" includes propyl, isopropyl and *t*-butyl. However, references to individual alkyl groups such as 'propyl' are specific for the straight chained version only and references to individual branched chain alkyl groups such as 'isopropyl' are specific for the branched chain version only. A similar convention applies to other radicals, for example " haloC_{1-4} alkyl" includes 1-chloroethyl and 2-fluoroethyl. The term "halo" refers to fluoro, chloro,
15 bromo and iodo. Where a phrase such as "any C_{1-6} alkyl group may be optionally substituted by one or more groups" for the avoidance of doubt, it is to be understood that this refers to all groups that contains a C_{1-6} alkyl group, for example this phrase would also relate to a C_{1-6} alkanoyl group if that was listed in the paragraph.

A "heterocyclic group" is a saturated, partially saturated or unsaturated, mono or bicyclic
20 ring containing 4-12 atoms of which at least one atom is chosen from nitrogen, sulphur or oxygen, which may, unless otherwise specified, be carbon or nitrogen linked, wherein a $-\text{CH}_2-$ group can optionally be replaced by a $-\text{C}(\text{O})-$, a ring nitrogen atom may optionally bear a C_{1-6} alkyl group and form a quaternary compound or a ring nitrogen and/or sulphur atom may be optionally oxidised to form the *N*-oxide and or the S-oxides. Examples and suitable values of the
25 term "heterocyclic group" are morpholino, piperidyl, pyridyl, pyranyl, pyrrolyl, isothiazolyl, indolyl, quinolyl, thienyl, 1,3-benzodioxolyl, thiadiazolyl, piperazinyl, thiazolidinyl, pyrrolidinyl, thiomorpholino, pyrrolinyl, homopiperazinyl, tetrahydropyranyl, imidazolyl, pyrimidyl, pyrazinyl, pyridazinyl, isoxazolyl, *N*-methylpyrrolyl, 4-pyridone, 1-isoquinolone, 2-pyrrolidone, 4-thiazolidone, pyridine-*N*-oxide and quinoline-*N*-oxide.

A "six membered heteroaryl ring containing one or two ring nitrogens where said nitrogens are optionally oxidised to form the *N*-oxide" is an unsaturated monocyclic ring containing six atoms. Examples are pyridine, pyrimidine, pyrazine and pyridine-*N*-oxide.

- Suitable values when R¹ and R² together with the nitrogen atom to which they are attached form a heterocyclic group are morpholino, piperidyl, piperazinyl, indolyl, thiazolidinyl, pyrrolidinyl, thiomorpholino, pyrrolinyl, homopiperazinyl, pyrrolyl, imidazolyl, pyrazolyl, pyrazolinyl, pyrazolidinyl, triazolyl, indolyl, isoindolyl and benzimidazolyl. Especially morpholino, piperidyl, piperazinyl, indolyl, thiazolidinyl, pyrrolidinyl, thiomorpholino, pyrrolinyl and homopiperazinyl.
- 10 An example of "C₁₋₆alkanoyloxy" is acetoxy. Examples of "C₁₋₆alkoxycarbonyl" include methoxycarbonyl, ethoxycarbonyl, *n*- and *t*-butoxycarbonyl. Examples of "C₁₋₆alkoxy" include methoxy, ethoxy and propoxy. Examples of "C₁₋₆alkanoylamino" include formamido, acetamido and propionylamino. Examples of "C₁₋₄alkylsulphanyl" include methylthio and ethylthio. Examples of "C₁₋₄alkylsulphinyl" include methylsulphinyl and ethylsulphinyl. Examples of
- 15 "C₁₋₄alkylsulphonyl" include mesyl and ethylsulphonyl. Examples of "C₁₋₆alkanoyl" include propionyl and acetyl. Examples of "C₁₋₆alkylamino" include methylamino and ethylamino. Examples of "(C₁₋₆alkyl)₂amino" include di-*N*-methylamino, di-(*N*-ethyl)amino and *N*-ethyl-*N*-methylamino. Examples of "C₁₋₆alkoxyC₁₋₆alkyl" methoxymethyl and propoxyethyl. Examples of "carbamoylC₁₋₆alkyl" are carbamoylmethyl and 2-carbamoyl ethyl. Examples of
- 20 "*N*-(C₁₋₆alkyl)carbamoylC₁₋₆alkyl" are *N*-(methyl)aminocarbonyl ethyl and *N*-(ethyl)aminocarbonyl propyl. Examples of "*N*-(C₁₋₆alkyl)₂carbamoylC₁₋₆alkyl" are *N,N*-(dimethyl)aminocarbonyl ethyl and *N*-(methyl)-*N*-(ethyl)aminocarbonyl propyl. Examples of "C₂₋₄alkenyloxy" are vinyloxy and allyloxy. Examples of "C₃₋₆cycloalkyl" are cyclopropyl and cyclohexyl. Examples of "C₂₋₆alkenyl" are vinyl, allyl and 1-propenyl. Examples of "C₂₋₆alkynyl"
- 25 are ethynyl, 1-propynyl and 2-propynyl. Examples of "haloC₁₋₄alkoxy" are 2-fluoroethoxy and 1-bromopropoxy. Examples of "C₁₋₆alkanoyl(*N*-C₁₋₆alkyl)amino" are (*N*-methyl)formamido and (*N*-propyl)acetamido. Examples of "C₁₋₆alkylsulphonylamino" are methylsulphonylamino and ethylsulphonylamino. Examples of "C₁₋₆alkylsulphonyl(*N*-C₁₋₆alkyl)amino" are methylsulphonyl-(*N*-ethyl)-amino and ethylsulphonyl-(*N*-butyl)-amino. Examples of
- 30 "*N*-(C₁₋₆alkyl)aminosulphonyl" are *N*-(methyl)aminosulphonyl and *N*-(ethyl)aminosulphonyl.

- Examples of "*N*-(C₁₋₆alkyl)₂aminosulphonyl" are *N,N*-(dimethyl)aminosulphonyl and *N*-(methyl)-*N*-(ethyl)aminosulphonyl. Examples of "*N*-(C₁₋₆alkyl)carbamoyl" are methylaminocarbonyl and ethylaminocarbonyl. Examples of "*N*-(C₁₋₆alkyl)₂carbamoyl" are dimethylaminocarbonyl and methylethylaminocarbonyl. Examples of "phenylC₁₋₆alkyl" are benzyl and phenethyl. Examples of "phenylC₁₋₆alkoxy" are benzyloxy and phenylethoxy. Examples of "C₁₋₆alkanoylaminosulphonyl" are acetylaminosulphonyl and propionylaminosulphonyl. Examples of "C₁₋₆alkanoyl(*N*-C₁₋₆alkyl)aminosulphonyl" are acetyl(*N*-methyl)aminosulphonyl and propionyl(*N*-ethyl)aminosulphonyl. Examples of "C₁₋₆alkylsulphonylaminocarbonyl" are mesylaminocarbonyl and ethanesulphonylaminocarbonyl.
- 10 Examples of "C₁₋₆alkylsulphonyl(*N*-C₁₋₆alkyl)aminocarbonyl" are *N*-(methyl)mesylaminocarbonyl and *N*-(methyl)ethanesulphonylaminocarbonyl. Examples of "C₁₋₆alkoxyC₁₋₆alkanoyl" are methoxyacetyl and ethoxypropionyl. Examples of "phenylC₁₋₆alkanoyl" are phenylacetyl and phenylpropionyl. Examples of "(heterocyclic group)carbonyl" are pyrid-3-ylcarbonyl and pyrimid-2-ylcarbonyl. Examples of "(heterocyclic group)C₁₋₆alkanoyl" are pyrid-3-ylacetyl and
- 15 pyrimid-2-ylpropionyl. Examples of "phenylC₁₋₆alkoxycarbonyl" are benzyloxycarbonyl and phenethyloxycarbonyl.

Another aspect of the invention provides a compound of formula (I) as depicted above wherein:

- Ring X is phenyl or a six membered heteroaryl ring containing one or two ring nitrogens where
- 20 said nitrogens are optionally oxidised to form the *N*-oxide;
- R¹ and R² are independently as defined in (a) or (b);
- R³ and R⁴ are as defined in (c) or (d);
- R⁵ is as defined in (e) or (f);
- Y-Z is as defined in (g) or (h)
- 25 wherein:

(a) either R¹ and R² are each selected independently from hydrogen, C₁₋₃alkyl, pyridyl and phenyl which is unsubstituted or substituted by one or two substituents selected independently from C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkenyloxy, hydroxy, halo and cyano,

- or R¹ and R² together with the nitrogen atom to which they are attached form morpholino,
- 30 thiomorpholino, piperidinyl, pyrrolidinyl or imidazolyl;

(b) either R^1 and R^2 are each selected independently from phenyl substituted by one or more P (wherein P is as defined hereinbelow), a heterocyclic group other than unsubstituted pyridyl which is optionally substituted by one or more Q (wherein Q is as defined hereinbelow), naphthyl optionally substituted by one or more Q, C_{4-6} alkyl, C_{3-6} cycloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkyl substituted by one or more V (wherein V is as defined hereinbelow), R^6T - (wherein R^6 and T are as defined hereinbelow) and $R^7C_{1-6}alkylT$ - (where R^7 is as defined hereinbelow),

or R^1 and R^2 together with the nitrogen atom to which they are attached form a heterocyclic group other than morpholino, thiomorpholino, piperidinyl, pyrrolidinyl or

imidazolyl;

(c) either R^3 and R^4 are independently C_k alkyl optionally substituted by from 1 to $2k+1$ atoms selected from fluoro and chloro wherein k is 1-3, provided that R^3 and R^4 are not both methyl;

or R^3 and R^4 together with the carbon atom to which they are attached, form a

C_m cycloalkyl ring optionally substituted by from 1 to $2m-2$ fluorine atoms wherein m is 3-5;

(d) R^3 and R^4 are both methyl;

(e) R^5 is hydrogen, C_{1-4} alkyl, halo C_{1-4} alkyl, C_{1-4} alkoxy, halo C_{1-4} alkoxy, cyano, nitro, C_{2-4} alkenyloxy or trifluoromethylthio;

(f) R^5 is halo, hydroxy, amino, C_{1-6} alkylamino, $(C_{1-6}alkyl)_2$ amino, C_{1-6} alkanoylamino,

C_{1-6} alkanoyl(N - $C_{1-6}alkyl$)amino, $C_{1-6}alkylsulphonamido$, $C_{1-6}alkylsulphonyl(N$ - $C_{1-6}alkyl$)amino, thiol, $C_{1-6}alkylsulphanyl$, $C_{1-6}alkylsulphiny$, $C_{1-6}alkylsulphonyl$, sulphamoyl, N -($C_{1-6}alkyl$)aminosulphonyl, N -($C_{1-6}alkyl$)₂aminosulphonyl, carboxy, carbamoyl, N -($C_{1-6}alkyl$)carbamoyl, N -($C_{1-6}alkyl$)₂carbamoyl, $C_{1-6}alkyloxycarbonyl$, $C_{1-6}alkanoyl$, C_{2-6} alkenyl, C_{2-6} alkynyl, $C_{5-6}alkyl$, halo $C_{5-6}alkyl$, $C_{5-6}alkoxy$, halo $C_{5-6}alkoxy$ or $C_{5-6}alkenyloxy$;

(g) Y-Z is -NHC(O)-, -OCH₂-, -SCH₂-, -NHCH₂-, trans-vinylene, and ethynylene;

(h) Y-Z is -NHC(S)-;

R^6 is selected from $C_{1-6}alkyl$, $C_{3-6}cycloalkyl$, a heterocyclic group optionally substituted by one or more R^8 , phenyl optionally substituted by one or more R^8 , naphthyl optionally substituted by one or more R^8 ;

R^7 is a heterocyclic group optionally substituted by one or more R^8 , phenyl optionally substituted by one or more R^8 , naphthyl optionally substituted by one or more R^8 ;

R^8 is trifluoromethyl, C_{1-6} alkyl, halo, hydroxy, trifluoromethoxy, cyano, C_{1-6} alkoxy, C_{1-6} alkanoyl, C_{1-6} alkanoyloxy, amino, C_{1-6} alkylamino, $(C_{1-6}alkyl)_2$ amino, C_{1-6} alkanoylamino, C_{1-6} alkanoyl(N - $C_{1-6}alkyl$)amino, nitro, carboxy, carbamoyl, C_{1-6} alkoxycarbonyl, thiol, C_{1-6} alkylsulphanyl, C_{1-6} alkylsulphinyl, C_{1-6} alkylsulphonyl, sulphamoyl, N -($C_{1-6}alkyl$)aminosulphonyl, N -($C_{1-6}alkyl$) $_2$ aminosulphonyl, carbamoyl $C_{1-6}alkyl$, N -($C_{1-6}alkyl$)carbamoyl $C_{1-6}alkyl$, N -($C_{1-6}alkyl$) $_2$ carbamoyl- $C_{1-6}alkyl$, hydroxy $C_{1-6}alkyl$ or $C_{1-6}alkoxyC_{1-6}alkyl$;

10 P is selected from R^9 - $C_{2-6}alkyl$ -M-, R^9 - $C_{2-6}alkenyl$ -M-, R^9 - $C_{2-6}alkynyl$ -M-, R^{10} -CH $_2$ -M-, R^{11} -M-, thiol, $C_{1-6}alkylsulphanyl$, $C_{1-6}alkylsulphinyl$, $C_{1-6}alkylsulphonyl$, sulphamoyl, nitro, carboxy, $C_{1-6}alkoxycarbonyl$, amino, $C_{1-6}alkylamino$, $(C_{1-6}alkyl)_2$ amino, carbamoyl, N -($C_{1-6}alkyl$)carbamoyl, N -($C_{1-6}alkyl$) $_2$ carbamoyl, C_{1-6} alkanoylamino, C_{1-6} alkanoyl(N - $C_{1-6}alkyl$)amino, trifluoromethyl, trifluoromethoxy, C_{1-6} alkanoyl, $C_{5-6}alkyl$, $C_{2-6}alkenyl$, $C_{2-6}alkynyl$, N -($C_{1-6}alkyl$)aminosulphonyl, N -($C_{1-6}alkyl$) $_2$ aminosulphonyl, $C_{5-6}alkoxy$, $C_{5-6}alkenyloxy$, phenyl optionally substituted by one or more R^8 , naphthyl optionally substituted by one or more R^8 , and a heterocyclic group optionally substituted by one or more R^8 ;

Q is selected from any of the values defined for P , $C_{1-4}alkyl$, $C_{1-4}alkoxy$, $C_{2-4}alkenyloxy$, hydroxy, halo and cyano;

20 V is selected from any of the values defined for Q , phenyl optionally substituted by one or more Q , naphthyl optionally substituted by one or more Q or a heterocyclic group optionally substituted by one or more Q ;

T is selected from -O-, -C(O)-, -NH-, -N(N - $C_{1-6}alkyl$)-, -C(O)NH-, -C(O)N(N - $C_{1-6}alkyl$)-, -SO $_2$ -, -C(S)-, -C(S)NH-, -C(S)N(N - $C_{1-6}alkyl$)-

25 M is selected from O, -N(R^{12})-, -N(R^{12})C(O)-, -C(O)N(R^{12})-, -S(O) $_n$ -, -OC(O)-, -C(O)O-, -N(R^{12})C(O)O-, -OC(O)N(R^{12})-, -C(S)N(R^{12})-, -N(R^{12})C(S)-, -SO $_2$ N(R^{12})-, -N(R^{12})SO $_2$ - and -N(R^{12})C(O)N(R^{12})-, -N(R^{12})C(S)N(R^{12})-, -SO $_2$ NHC(O)-, -C(O)NH(SO $_2$)- or M is a direct bond;

R^9 is selected from hydroxy, amino, $C_{1-6}alkylamino$, $(C_{1-6}alkyl)_2$ amino, carboxy, $C_{1-6}alkoxy$, carbamoyl, N -($C_{1-6}alkyl$)carbamoyl, N -($C_{1-6}alkyl$) $_2$ carbamoyl, sulphamoyl,

30 N - $C_{1-6}alkylaminosulphonyl$, N -($C_{1-6}alkyl$) $_2$ aminosulphonyl, a heterocyclic group optionally

substituted by one or more R^8 , phenyl optionally substituted by one or more R^8 , naphthyl optionally substituted by one or more R^8 , C_{1-6} alkylsulphanyl, C_{1-6} alkylsulphinyl and C_{1-6} alkylsulphonyl;

R^{10} is carboxy, carbamoyl, N -(C_{1-6} alkyl)carbamoyl, N -(C_{1-6} alkyl)₂carbamoyl, sulphamoyl,
5 N -(C_{1-6} alkyl)aminosulphonyl, N -(C_{1-6} alkyl)₂aminosulphonyl, C_{1-6} alkylsulphonyl, a heterocyclic group optionally substituted by one or more R^8 , phenyl optionally substituted by one or more R^8 or naphthyl optionally substituted by one or more R^8 ;

R^{11} is a heterocyclic group optionally substituted by one or more R^8 , phenyl optionally substituted by one or more R^8 or naphthyl optionally substituted by one or more R^8 ;

10 R^{12} is hydrogen or C_{1-6} alkyl;

n is 0-2;

with the proviso that where R^1 and R^2 are both as defined in (a), R^3 and R^4 are both as defined in (c), R^5 is as defined in (e) and Ring X is phenyl, Y-Z must be -NHC(S)-; and pharmaceutically acceptable salts or *in vivo* hydrolysable esters thereof,

15 provided said compound is not

4-(3-hydroxy-3-methyl-1-butyryl)- N -(3-methyl-2-pyridinyl)-benzenesulphonamide;

N -{4-[N,N -bis-(*sec*-butyl)aminosulphonyl]phenyl}-2-hydroxy-2-methyl-3,3,3-trifluoropropanamide, or

N -{4-[N,N -bis-(*iso*-butyl)aminosulphonyl]phenyl}-2-hydroxy-2-methyl-3,3,3-

20 trifluoropropanamide.

A further aspect of the invention provides a compound of formula (I) as depicted above wherein:

Ring X is phenyl or a six membered heteroaryl ring containing one or two ring nitrogens where said nitrogens are optionally oxidised to form the *N*-oxide;

25 either R^1 and R^2 are each selected independently from hydrogen, C_{1-6} alkyl, phenyl optionally substituted by one or more P, a heterocyclic group optionally substituted on a ring carbon by one or more P and wherein if said heterocyclic group contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from D, naphthyl optionally substituted by one or more P, C_{3-6} cycloalkyl optionally substituted with one or more P,
30 C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkyl substituted by one or more V, R^6T - and $R^7C_{1-6}alkylT$ -,

or R^1 and R^2 together with the nitrogen atom to which they are attached form a heterocyclic group optionally substituted on a ring carbon by one or more P and wherein if said heterocyclic group contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from D;

- 5 either R^3 and R^4 are independently C_k alkyl optionally substituted by from 1 to $2k+1$ atoms selected from fluoro and chloro wherein k is 1-3;

or R^3 and R^4 together with the carbon atom to which they are attached, form a C_m cycloalkyl ring optionally substituted by from 1 to $2m-2$ fluorine atoms wherein m is 3-5;

R^5 is halo, hydroxy, amino, C_{1-6} alkylamino, $(C_{1-6}$ alkyl) $_2$ amino, C_{1-6} alkanoylamino,

- 10 C_{1-6} alkanoyl(N - C_{1-6} alkyl)amino, C_{1-6} alkylsulphonylamino, C_{1-6} alkylsulphonyl(N - C_{1-6} alkyl)amino, thiol, C_{1-6} alkylsulphanyl, C_{1-6} alkylsulphinyl, C_{1-6} alkylsulphonyl, sulphamoylsulphonamido, N -(C_{1-6} alkyl)aminosulphonyl, N -(C_{1-6} alkyl) $_2$ aminosulphonyl, carboxy, carbamoyl, N -(C_{1-6} alkyl)carbamoyl, N -(C_{1-6} alkyl) $_2$ carbamoyl, C_{1-6} alkoxyalkyloxycarbonyl, formyl, C_{1-6} alkanoyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{5-6} alkyl, halo C_{5-6} alkyl, C_{5-6} alkoxy, halo C_{5-6} alkoxy or

- 15 C_{5-6} alkenyloxy;

$Y-Z$ is -NHC(O)-, -NHC(S)-, -OCH $_2$ -, -SCH $_2$ -, -NHCH $_2$ -, trans-vinylene, and ethynylene;

R^6 is selected from C_{1-6} alkyl (optionally substituted with one or more R^8), C_{3-6} cycloalkyl optionally substituted with one or more R^8 , a heterocyclic group optionally substituted on a ring carbon by one or more R^8 and if said heterocyclic group contains an -NH- moiety that nitrogen

- 20 may be optionally substituted by a group selected from D, phenyl optionally substituted by one or more R^8 , naphthyl optionally substituted by one or more R^8 ;

R^7 is a heterocyclic group optionally substituted on a ring carbon by one or more R^8 and if said heterocyclic group contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from D, phenyl optionally substituted by one or more R^8 , naphthyl optionally

- 25 substituted by one or more R^8 ;

R^8 is trifluoromethyl, C_{1-6} alkyl, halo, hydroxy, trifluoromethoxy, cyano, C_{1-6} alkoxy, formyl, C_{1-6} alkanoyl, C_{1-6} alkanoyloxy, amino, C_{1-6} alkylamino, $(C_{1-6}$ alkyl) $_2$ amino,

C_{1-6} alkanoylamino, C_{1-6} alkanoyl(N - C_{1-6} alkyl)amino, nitro, carboxy, carbamoyl,

C_{1-6} alkoxycarbonyl, thiol, C_{1-6} alkylsulphanyl, C_{1-6} alkylsulphinyl, C_{1-6} alkylsulphonyl, sulphamoyl,

- 30 N -(C_{1-6} alkyl)aminosulphonyl, N -(C_{1-6} alkyl) $_2$ aminosulphonyl, carbamoyl C_{1-6} alkyl,

$N-(C_{1-6}\text{alkyl})\text{carbamoyl}C_{1-6}\text{alkyl}$, $N-(C_{1-6}\text{alkyl})_2\text{carbamoyl}-C_{1-6}\text{alkyl}$, hydroxy $C_{1-6}\text{alkyl}$, $C_{1-6}\text{alkoxy}C_{1-6}\text{alkyl}$, phenyl $C_{1-6}\text{alkyl}$ or phenyl $C_{1-6}\text{alkoxy}$;

P is selected from $C_{2-6}\text{alkyl}-M$ - substituted with one or more R^9 , $C_{2-6}\text{alkenyl}-M$ - optionally substituted with one or more R^9 , $C_{2-6}\text{alkynyl}-M$ - optionally substituted with one or more R^9 (with the proviso that in the three previous groups R^9 is not a substituent on the carbon atom attached to M), $R^{10}-CH_2-M$ -, $R^{11}-M$ -, thiol, $C_{1-6}\text{alkylsulphanyl}$, $C_{1-6}\text{alkylsulphinyl}$, $C_{1-6}\text{alkylsulphonyl}$, sulphamoyl, nitro, carboxy, hydroxy, halo, cyano $C_{1-6}\text{alkoxycarbonyl}$, amino, $C_{1-6}\text{alkylamino}$, $(C_{1-6}\text{alkyl})_2\text{amino}$, carbamoyl, $N-(C_{1-6}\text{alkyl})\text{carbamoyl}$, $N-(C_{1-6}\text{alkyl})_2\text{carbamoyl}$, $C_{1-6}\text{alkanoylamino}$, $C_{1-6}\text{alkanoyl}(N-C_{1-6}\text{alkyl})\text{amino}$, trifluoromethyl, trifluoromethoxy, formyl, $C_{1-6}\text{alkanoyl}$, $C_{1-6}\text{alkyl}$, $C_{2-6}\text{alkenyl}$, $C_{2-6}\text{alkynyl}$, $N-(C_{1-6}\text{alkyl})\text{aminosulphonyl}$, hydroxymethyl, hydroxyacetyl, $N-(C_{1-6}\text{alkyl})_2\text{aminosulphonyl}$, $C_{1-6}\text{alkanoylaminosulphonyl}$, $C_{1-6}\text{alkanoyl}(N-C_{1-6}\text{alkyl})\text{aminosulphonyl}$, $C_{1-6}\text{alkylsulphonylaminocarbonyl}$, $C_{1-6}\text{alkylsulphonyl}(N-C_{1-6}\text{alkyl})\text{aminocarbonyl}$, $C_{1-6}\text{alkoxy}$, $C_{2-6}\text{alkenyloxy}$, phenyl optionally substituted by one or more R^8 , naphthyl optionally substituted by one or more R^8 and a heterocyclic group optionally substituted on a ring carbon by one or more R^8 and if said heterocyclic group contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from D;

V is selected from any of the values defined for **P**, phenyl optionally substituted by one or more **P**, naphthyl optionally substituted by one or more **P**, a heterocyclic group optionally substituted on a ring carbon by one or more **P** and if said heterocyclic group contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from D or $C_{3-6}\text{cycloalkyl}$ optionally substituted with one or more **P**;

T is selected from -O-, -C(O)-, -NH-, -N($N-C_{1-6}\text{alkyl}$)-, -C(O)NH-, -NHC(O)-, -C(O)N($N-C_{1-6}\text{alkyl}$)-, -N($N-C_{1-6}\text{alkyl}$)C(O)-, -SO₂-, -C(S)-, -C(S)NH-, -NHC(S)-, -C(S)N($N-C_{1-6}\text{alkyl}$)- and -N($N-C_{1-6}\text{alkyl}$)C(S)-;

M is selected from -O-, -N(R^{12})-, -C(O)-, -N(R^{12})C(O)-, -C(O)N(R^{12})-, -S(O)_n-, -OC(O)-, -C(O)O-, -N(R^{12})C(O)O-, -OC(O)N(R^{12})-, -C(S)N(R^{12})-, -N(R^{12})C(S)-, -SO₂N(R^{12})-, -N(R^{12})SO₂- and -N(R^{12})C(O)N(R^{12})-, -N(R^{12})C(S)N(R^{12})-, -SO₂NHC(O)-, -SO₂N(R^{12})C(O)-, -C(O)NHSO₂-, -C(O)N(R^{12})SO₂- or **M** is a direct bond;

D is selected from C_{1-6} alkyl, C_{1-6} alkanoyl, C_{1-6} alkylsulphonyl, C_{1-6} alkoxycarbonyl, carbamoyl, N -(C_{1-6} alkyl)carbamoyl, N,N -(C_{1-6} alkyl)₂carbamoyl, benzoyl, (heterocyclic group)carbonyl, phenylsulphonyl, (heterocyclic group)sulphonyl, phenyl or a carbon linked heterocyclic group; and wherein any C_{1-6} alkyl group may be optionally substituted by one or more R^9 ; and wherein any phenyl or heterocyclic group may be optionally substituted on a ring carbon by one or more groups selected from R^8 and if a heterocyclic group contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from E;

E is selected from C_{1-6} alkyl, C_{1-6} alkanoyl, C_{1-6} alkylsulphonyl, C_{1-6} alkoxycarbonyl, carbamoyl, N -(C_{1-6} alkyl)carbamoyl, N,N -(C_{1-6} alkyl)₂carbamoyl, C_{1-6} alkoxy C_{1-6} alkanoyl, phenyl C_{1-6} alkyl, benzoyl, phenyl C_{1-6} alkanoyl, phenyl C_{1-6} alkoxycarbonyl and phenylsulphonyl.

R^9 is selected from hydroxy, amino, C_{1-6} alkylamino, (C_{1-6} alkyl)₂amino, carboxy, C_{1-6} alkoxy, carbamoyl, N -(C_{1-6} alkyl)carbamoyl, N -(C_{1-6} alkyl)₂carbamoyl, formyl, sulphamoyl, N - C_{1-6} alkylaminosulphonyl, N -(C_{1-6} alkyl)₂aminosulphonyl, C_{1-6} alkylsulphonylamino, C_{1-6} alkanoylamino, a heterocyclic group optionally substituted on a ring carbon by one or more R^8 and if said heterocyclic group contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from D, phenyl optionally substituted by one or more R^8 , naphthyl optionally substituted by one or more R^8 , C_{1-6} alkylsulphanyl, C_{1-6} alkylsulphanyl and C_{1-6} alkylsulphonyl;

R^{10} is carboxy, carbamoyl, N -(C_{1-6} alkyl)carbamoyl, N -(C_{1-6} alkyl)₂carbamoyl, sulphamoyl, N -(C_{1-6} alkyl)aminosulphonyl, N -(C_{1-6} alkyl)₂aminosulphonyl, C_{1-6} alkylsulphanyl, C_{1-6} alkylsulphanyl, C_{1-6} alkylsulphonyl, C_{1-6} alkoxycarbonyl, C_{1-6} alkanoylamino, a heterocyclic group optionally substituted on a ring carbon by one or more R^8 and if said heterocyclic group contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from D, phenyl optionally substituted by one or more R^8 or naphthyl optionally substituted by one or more R^8 ;

R^{11} is a heterocyclic group optionally substituted on a ring carbon by one or more R^8 and if said heterocyclic group contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from D, phenyl optionally substituted by one or more R^8 , C_{3-6} cycloalkyl optionally substituted by one or more R^8 , or naphthyl optionally substituted by one or more R^8 ;

R^{12} is hydrogen or C_{1-6} alkyl optionally substituted with R^{13} with the proviso that R^{13} is not a substituent on the carbon attached to the nitrogen atom of M;

R^{13} is halo, hydroxy, amino, cyano, nitro, trifluoromethyl, trifluoromethoxy, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkylamino, $(C_{1-6}alkyl)_2$ amino, C_{1-6} alkanoylamino,

5 C_{1-6} alkanoyl(N - C_{1-6} alkyl)amino, C_{1-6} alkylsulphonylamino, C_{1-6} alkylsulphonyl(N - C_{1-6} alkyl)amino, thiol, C_{1-6} alkylsulphanyl, C_{1-6} alkylsulphinyl, C_{1-6} alkylsulphonyl; sulphamoyl, N -(C_{1-6} alkyl)aminosulphonyl, N -(C_{1-6} alkyl) $_2$ aminosulphonyl, carboxy, carbamoyl, N -(C_{1-6} alkyl)carbamoyl, N -(C_{1-6} alkyl) $_2$ carbamoyl, C_{1-6} alkoxycarbonyl, C_{1-6} alkanoyl or formyl;

n is 0-2;

10 and pharmaceutically acceptable salts or *in vivo* hydrolysable esters thereof.

Preferred values for R^1 , R^2 , R^3 , R^4 , R^5 , X and Y-Z are as follows.

Preferably R^1 and R^2 are each independently selected from hydrogen, C_{1-6} alkyl, C_{2-6} alkenyl, C_{3-6} cycloalkyl, C_{1-6} alkyl substituted by a heterocyclic group, C_{1-6} alkyl substituted by phenyl (which phenyl is optionally substituted by one or more substituents selected from halo,

15 trifluoromethoxy, a heterocyclic group and trifluoromethyl), and phenyl which is optionally substituted by one or more substituents selected from halo, C_{1-4} alkoxy, carbamoyl, trifluoromethyl, C_{1-4} alkyl, nitro, hydroxy, cyano, C_{1-6} alkanolyamino, C_{1-6} alkylsulphonyl, C_{1-6} alkanoyl and sulphamoyl,

or R^1 and R^2 together with the nitrogen group to which they are attached form morpholino
20 or piperazine.

More preferably R^1 and R^2 are each independently selected from hydrogen, C_{1-5} alkyl, C_3 alkenyl, C_{3-6} cycloalkyl, pyridylCH $_2$ -, thienylCH $_2$ -, 1,3-benzodioxylCH $_2$ -, phenylCH $_2$ - (which phenyl is optionally substituted by one or more substituents selected from fluoro, chloro, trifluoromethoxy, thiadiazole, and trifluoromethyl) and phenyl which is optionally substituted by
25 one or more substituents selected from fluoro, chloro, bromo, iodo, methoxy, hydroxy, carbamoyl, C_{1-4} alkyl, trifluoromethyl, nitro, cyano, sulphamoyl, C_{1-2} alkanolyamino, mesyl and C_{1-2} alkanoyl,

or R^1 and R^2 together with the nitrogen group to which they are attached form morpholino or piperazine.

Particularly R^1 and R^2 are each independently selected from hydrogen, methyl, ethyl, propenyl and phenyl which is substituted by one substituent selected from methoxy, chloro, iodo, hydroxy, carbamoyl, cyano, acetylamino, mesyl, acetyl and sulphamoyl;

or R^1 and R^2 together with the nitrogen group to which they are attached form morpholino.

More particularly R^1 and R^2 are each independently selected from hydrogen and phenyl which is substituted by one substituent selected from methoxy, hydroxy, carbamoyl, cyano, acetyl, mesyl and sulphamoyl.

Preferred combinations of R^1 and R^2 are as follows.

Preferably R^1 and R^2 are both methyl or ethyl, or one of R^1 and R^2 is methyl, ethyl or optionally substituted phenyl and the other is hydrogen, or R^1 and R^2 together with the nitrogen to which they are attached form morpholino.

More preferably one of R^1 and R^2 is hydrogen and the other is phenyl substituted with one substituent selected from methoxy, hydroxy, carbamoyl, cyano, acetyl, mesyl and sulphamoyl.

In another aspect of the invention, preferably R^1 and R^2 are each independently selected from

i) hydrogen;

ii) C_{1-6} alkyl optionally substituted with one or more hydroxy, halo, C_{1-6} alkoxy (optionally substituted with one or more hydroxy), C_{1-6} alkoxycarbonyl, C_{1-6} alkanoylamino, N,N -(C_{1-6} alkyl)₂amino, C_{1-6} alkylsulphanyl, N -(C_{1-6} alkyl)carbamoyl (optionally substituted with one or more C_{1-6} alkoxycarbonyl), phenyl (optionally substituted with one or more halo, trifluoromethyl, C_{1-6} alkoxy, C_{1-6} alkyl, trifluoromethoxy, thiadiazolyl or sulphamoyl), pyridyl, thienyl, 1,3-benzodioxolyl, morpholino, piperidinyl, tetrahydrofuran and imidazolyl;

iii) phenyl optionally substituted with one or more halo, trifluoromethyl, nitro, cyano, hydroxy, sulphamoyl, carbamoyl, amino, formyl, carboxy, C_{1-4} alkyl (optionally substituted with one or more hydroxy), C_{1-4} alkoxy (optionally substituted with one or more phenyl), C_{1-6} alkanoyl, C_{1-6} alkoxycarbonyl, C_{1-6} alkanoylamino, C_{1-6} alkylsulphanyl (optionally substituted with one or more hydroxy or C_{1-6} alkoxy), C_{1-6} alkylsulphonyl (optionally substituted with one or more hydroxy or C_{1-6} alkoxy), N -(C_{1-6} alkyl)amino (optionally substituted with one or more hydroxy), N -(C_{1-6} alkyl)carbamoyl (optionally substituted with one or more C_{1-6} alkoxy, N,N -(C_{1-6} alkyl)₂amino, hydroxy or tetrahydrofuran), N,N -(C_{1-6} alkyl)₂carbamoyl (optionally

- substituted with one or more hydroxy or C_{1-6} alkoxy), $N-(C_{2-6}$ alkenyl)carbamoyl, $N-(C_{3-6}$ cycloalkyl)carbamoyl (optionally substituted with one or more hydroxy), $N-(C_{2-6}$ alkenyl)aminosulphonyl, $N-(C_{3-6}$ cycloalkyl)aminosulphonyl (optionally substituted with one or more hydroxy), $N-(C_{1-6}$ alkyl)aminosulphonyl (optionally substituted with one or more hydroxy, amino or $N,N-(C_{1-6}$ alkyl)₂amino), $N,N-(C_{1-6}$ alkyl)₂aminosulphonyl (optionally substituted with one or more hydroxy), C_{1-6} alkanoylamino sulphonyl, C_{1-6} alkylsulphonylamino carbonyl, morpholin sulphonyl, piperazinyl carbonyl (optionally substituted on a ring nitrogen by C_{1-6} alkyl), morpholino carbonyl, pyrrolidinyl sulphonyl (optionally substituted with one or more hydroxy);
- 10 iv) C_{3-6} cycloalkyl (optionally substituted with one or more hydroxy);
- v) C_{2-6} alkenyl,
- vi) C_{1-6} alkanoyl [optionally substituted with one or more amino or phenyl (wherein said phenyl is optionally substituted with one or more halo)];
- vii) benzoyl (optionally substituted with one or more halo, C_{1-6} alkylsulphonyl, C_{1-6} alkylsulphonyl, cyano, benzyloxy, C_{1-6} alkoxy, trifluoromethyl, $N,N-(C_{1-6}$ alkyl)₂aminosulphonyl, hydroxy);
- 15 viii) N -phenylcarbamoyl; or
- ix) a heterocyclic group selected from tetrahydrofuran, piperidinyl (optionally substituted on a ring nitrogen with C_{1-6} alkyl), pyridyl (optionally substituted with one or more trifluoromethyl, C_{1-6} alkyl or halo), pyrimidinyl (optionally substituted with one or more C_{1-6} alkyl or
- 20 $N,N-(C_{1-6}$ alkyl)₂amino), thienyl (optionally substituted with one or more carbamoyl), isoxazolyl (optionally substituted with one or more C_{1-6} alkyl) and pyrazinyl (optionally substituted with one or more C_{1-6} alkyl);

or R^1 and R^2 together with the nitrogen atom to which they are attached form morpholino, piperidinyl [optionally substituted with one or more hydroxy, C_{1-6} alkyl (wherein said C_{1-6} alkyl is optionally substituted with hydroxy) or pyrrolidinyl], piperazinyl [optionally substituted on a ring nitrogen by C_{1-6} alkoxycarbonyl, C_{1-6} alkanoyl (wherein said C_{1-6} alkanoyl is optionally substituted with one or more C_{1-6} alkoxy, C_{1-6} alkanoylamino, hydroxy or amino), C_{1-6} alkyl (wherein said C_{1-6} alkyl is optionally substituted with one or more hydroxy), C_{1-6} alkylsulphonyl, phenylsulphonyl (wherein said phenylsulphonyl is optionally substituted with one or more

25 C_{1-6} alkylsulphonyl) morpholino carbonyl or tetrahydrofuryl carbonyl], indolinyl, pyrrolyl,

30

thiazolidinyl, pyrrolidinyl (optionally substituted with one or more hydroxy), thiomorpholino, 3-pyrrolinyl or homopiperazinyl (optionally substituted on a ring nitrogen with C₁₋₆alkyl).

More preferably R¹ and R² are each independently selected from:

- i) hydrogen;
- 5 ii) C₁₋₆alkyl optionally substituted with one or more hydroxy, halo, C₁₋₆alkoxy (optionally substituted with one or more hydroxy), C₁₋₆alkoxycarbonyl, *N,N*-(C₁₋₆alkyl)₂amino, C₁₋₆alkylsulphanyl, *N*-(C₁₋₆alkyl)carbamoyl (optionally substituted with one or more C₁₋₆alkoxycarbonyl), phenyl (optionally substituted with one or more halo or sulphamoyl), pyridyl, thienyl, 1,3-benzodioxolyl, morpholino, piperidinyl, tetrahydrofuran and imidazolyl;
- 10 iii) phenyl optionally substituted with one or more halo, trifluoromethyl, nitro, cyano, hydroxy, sulphamoyl, carbamoyl, amino, formyl, carboxy, C₁₋₄alkyl (optionally substituted with one or more hydroxy), C₁₋₄alkoxy (optionally substituted with one or more phenyl), C₁₋₆alkanoyl, C₁₋₆alkoxycarbonyl, C₁₋₆alkanoylamino, C₁₋₆alkylsulphanyl (optionally substituted with one or more hydroxy or C₁₋₆alkoxy), C₁₋₆alkylsulphonyl (optionally substituted with one or more
- 15 hydroxy or C₁₋₆alkoxy), *N*-(C₁₋₆alkyl)amino (optionally substituted with one or more hydroxy), *N*-(C₁₋₆alkyl)carbamoyl (optionally substituted with one or more C₁₋₆alkoxy, *N,N*-(C₁₋₆alkyl)₂amino, hydroxy or tetrahydrofuran), *N,N*-(C₁₋₆alkyl)₂carbamoyl (optionally substituted with one or more hydroxy or C₁₋₆alkoxy), *N*-(C₂₋₆alkenyl)carbamoyl, *N*-(C₃₋₆cycloalkyl)carbamoyl (optionally substituted with one or more hydroxy),
- 20 *N*-(C₂₋₆alkenyl)aminosulphonyl, *N*-(C₃₋₆cycloalkyl)aminosulphonyl (optionally substituted with one or more hydroxy), *N*-(C₁₋₆alkyl)aminosulphonyl (optionally substituted with one or more hydroxy, or *N,N*-(C₁₋₆alkyl)₂amino), *N,N*-(C₁₋₆alkyl)₂aminosulphonyl (optionally substituted with one or more hydroxy), C₁₋₆alkanoylaminosulphonyl, C₁₋₆alkylsulphonylaminocarbonyl, morpholinosulphonyl, piperazinylcarbonyl (optionally substituted on a ring nitrogen by
- 25 C₁₋₆alkyl), morpholinocarbonyl, pyrrolidinylsulphonyl (optionally substituted with one or more hydroxy);
- iv) C₃₋₆cycloalkyl (optionally substituted with one or more hydroxy);
- v) C₂₋₆alkenyl,
- vi) C₁₋₆alkanoyl [optionally substituted with one or more phenyl (wherein said phenyl is
- 30 optionally substituted with one or more halo)];

- vii) benzoyl (optionally substituted with one or more halo, C₁₋₆alkylsulphanyl, C₁₋₆alkylsulphonyl, cyano, C₁₋₆alkoxy, trifluoromethyl, *N,N*-(C₁₋₆alkyl)₂aminosulphonyl, hydroxy);
- viii) a heterocyclic group selected from tetrahydrofuran, pyridyl (optionally substituted with one or more trifluoromethyl, C₁₋₆alkyl or halo), pyrimidinyl (optionally substituted with one or more C₁₋₆alkyl or *N,N*-(C₁₋₆alkyl)₂amino), thienyl (optionally substituted with one or more carbamoyl), isoxazolyl (optionally substituted with one or more C₁₋₆alkyl) and pyrazinyl (optionally substituted with one or more C₁₋₆alkyl);
- ix) R¹ and R² together with the nitrogen atom to which they are attached form morpholino, piperidinyl [optionally substituted with one or more hydroxy, C₁₋₆alkyl (wherein said C₁₋₆alkyl is optionally substituted with hydroxy) or pyrrolidinyl], piperazinyl [optionally substituted on a ring nitrogen by C₁₋₆alkoxycarbonyl, C₁₋₆alkanoyl (wherein said C₁₋₆alkanoyl is optionally substituted with one or more C₁₋₆alkoxy, C₁₋₆alkanoylamino, hydroxy or amino), C₁₋₆alkyl (wherein said C₁₋₆alkyl is optionally substituted with one or more hydroxy), C₁₋₆alkylsulphonyl, phenylsulphonyl (wherein said phenylsulphonyl is optionally substituted with one or more C₁₋₆alkylsulphonyl) or morpholinocarbonyl], indolinyl, thiazolidinyl, pyrrolidinyl (optionally substituted with one or more hydroxy), thiomorpholino or 3-pyrrolinyl.

Particularly R¹ and R² are each independently selected from:

- i) hydrogen;
- ii) C₁₋₆alkyl optionally substituted with one or more hydroxy, C₁₋₆alkoxy, C₁₋₆alkoxycarbonyl, C₁₋₆alkylsulphanyl, *N*-(C₁₋₆alkyl)carbamoyl (optionally substituted with one or more C₁₋₆alkoxycarbonyl), phenyl (optionally substituted with one or more sulphamoyl), morpholino and tetrahydrofuran;
- iii) phenyl optionally substituted with one or more halo, trifluoromethyl, hydroxy, sulphamoyl, amino, C₁₋₄alkyl, C₁₋₄alkoxy, C₁₋₆alkoxycarbonyl, C₁₋₆alkylsulphanyl (optionally substituted with one or more hydroxy or C₁₋₆alkoxy), C₁₋₆alkylsulphonyl (optionally substituted with one or more hydroxy or C₁₋₆alkoxy), *N*-(C₁₋₆alkyl)amino (optionally substituted with one or more hydroxy), *N*-(C₁₋₆alkyl)carbamoyl (optionally substituted with one or more C₁₋₆alkoxy, hydroxy or tetrahydrofuran), *N,N*-(C₁₋₆alkyl)₂carbamoyl (optionally substituted with one or more hydroxy), *N*-(C₂₋₆alkenyl)carbamoyl, *N*-(C₂₋₆alkenyl)aminosulphonyl, *N*-(C₁₋₆alkyl)aminosulphonyl (optionally substituted with one or more hydroxy or

- N,N*-(C₁₋₆alkyl)₂amino), *N,N*-(C₁₋₆alkyl)₂aminosulphonyl, C₁₋₆alkylsulphonylaminocarbonyl, morpholinosulphonyl, pyrrolidinylsulphonyl (optionally substituted with one or more hydroxy);
- iv) C₃₋₆cycloalkyl (optionally substituted with one or more hydroxy);
- v) C₂₋₆alkenyl,
- 5 vi) C₁₋₆alkanoyl [optionally substituted with one or more phenyl,
- vii) benzoyl (optionally substituted with one or more halo, cyano, C₁₋₆alkoxy, hydroxy);
- viii) a heterocyclic group selected from pyridyl (optionally substituted with one or more trifluoromethyl, C₁₋₆alkyl or halo), pyrimidinyl (optionally substituted with one or more C₁₋₆alkyl or *N,N*-(C₁₋₆alkyl)₂amino), thienyl (optionally substituted with one or more carbamoyl),
- 10 isoxazolyl (optionally substituted with one or more C₁₋₆alkyl) and pyrazinyl (optionally substituted with one or more C₁₋₆alkyl);

or R¹ and R² together with the nitrogen atom to which they are attached form morpholino, piperidinyl [optionally substituted with one or more hydroxy or C₁₋₆alkyl (wherein said C₁₋₆alkyl is optionally substituted with hydroxy)], piperazinyl [optionally substituted on a ring nitrogen by

15 C₁₋₆alkoxycarbonyl, C₁₋₆alkanoyl (wherein said C₁₋₆alkanoyl is optionally substituted with one or more C₁₋₆alkoxy, C₁₋₆alkanoylamino, hydroxy or amino), C₁₋₆alkylsulphonyl or morpholinocarbonyl], indolinyl or 3-pyrrolinyl.

More particularly R¹ and R² are each independently selected from hydrogen, methyl, cyclopropyl, 4-hydroxycyclohexyl, 2-hydroxyethyl, 2-hydroxypropyl, 2,3-dihydroxypropyl,

20 4-(morpholinosulphonyl)phenyl, pyrid-3-yl, 2-carbamoylthien-3-yl, 2-chloropyrid-3-yl, 5-chloropyrid-2-yl, 5-methylpyrid-2-yl, pyrimid-2-yl, 4,6-dimethylpyrimid-2-yl or 5,6-dimethylpyrazin-2-yl;

or R¹ and R² together with the nitrogen atom to which they are attached form 4-hydroxypiperidinyl or 1-(hydroxyacetyl)piperazin-4-yl.

- 25 In a further aspect of the invention, preferably R¹ and R² are independently selected from
- i) hydrogen;
- ii) C₁₋₆alkyl or C₃₋₆cycloalkyl optionally substituted with one or more C₁₋₆alkyl, C₁₋₆alkoxy, C₂₋₆alkenyloxy, hydroxy, halo, cyano, C₁₋₆alkylsulphanyl, C₁₋₆alkylsulphinyl, C₁₋₆alkylsulphonyl, sulphamoyl, carboxy, C₁₋₆alkoxycarbonyl, amino, C₁₋₆alkylamino, (C₁₋₆alkyl)₂amino, carbamoyl,
- 30 *N*-(C₁₋₆alkyl)carbamoyl, *N*-(C₁₋₆alkyl)₂carbamoyl, C₁₋₆alkanoylamino,

- C_{1-6} alkanoyl(N - C_{1-6} alkyl)amino, C_{1-6} alkanoyl, C_{2-6} alkenyl, C_{2-6} alkynyl, N -(C_{1-6} alkyl)aminosulphonyl, hydroxymethyl, hydroxyacetyl or N -(C_{1-6} alkyl)₂aminosulphonyl;
- (iii) a heterocyclic group selected from pyridyl, pyrimidyl, pyridazinyl or pyrazinyl, wherein said heterocyclic group is optionally substituted with one or more C_{1-6} alkyl, C_{1-6} alkoxy, C_{2-6} alkenyloxy, hydroxy, halo, cyano, C_{1-6} alkylsulphanyl, C_{1-6} alkylsulphinyl, C_{1-6} alkylsulphonyl, sulphamoyl, carboxy, C_{1-6} alkoxycarbonyl, amino, C_{1-6} alkylamino, (C_{1-6} alkyl)₂amino, carbamoyl, N -(C_{1-6} alkyl)carbamoyl, N -(C_{1-6} alkyl)₂carbamoyl, C_{1-6} alkanoylamino, C_{1-6} alkanoyl(N - C_{1-6} alkyl)amino, trifluoromethyl, trifluoromethoxy, C_{1-6} alkanoyl, C_{2-6} alkenyl, C_{2-6} alkynyl, N -(C_{1-6} alkyl)aminosulphonyl, hydroxymethyl, hydroxyacetyl or
- 10 N -(C_{1-6} alkyl)₂aminosulphonyl; and R^1 and R^2 together with the nitrogen atom to which they are attached form piperidinyl or piperazinyl; wherein said piperidinyl and piperazinyl may be optionally substituted on a ring carbon by one or more groups selected from C_{1-6} alkyl, C_{1-6} alkoxy, C_{2-6} alkenyloxy, hydroxy, halo, cyano, C_{1-6} alkylsulphanyl, C_{1-6} alkylsulphinyl, C_{1-6} alkylsulphonyl, sulphamoyl, carboxy,
- 15 C_{1-6} alkoxycarbonyl, amino, C_{1-6} alkylamino, (C_{1-6} alkyl)₂amino, carbamoyl, N -(C_{1-6} alkyl)carbamoyl, N -(C_{1-6} alkyl)₂carbamoyl, C_{1-6} alkanoylamino, C_{1-6} alkanoyl(N - C_{1-6} alkyl)amino, C_{1-6} alkanoyl, C_{2-6} alkenyl, C_{2-6} alkynyl, N -(C_{1-6} alkyl)aminosulphonyl, hydroxymethyl, hydroxyacetyl or N -(C_{1-6} alkyl)₂aminosulphonyl; and said piperazinyl may be optionally substituted on the ring nitrogen by a group selected from
- 20 C_{1-6} alkanoyl, C_{1-6} alkylsulphonyl, C_{1-6} alkoxycarbonyl, carbamoyl, N -(C_{1-6} alkyl)carbamoyl and N,N -(C_{1-6} alkyl)₂carbamoyl; and wherein any C_{1-6} alkyl group may be optionally substituted by one or more groups selected from hydroxy, amino, C_{1-6} alkylamino, (C_{1-6} alkyl)₂amino, carboxy, C_{1-6} alkoxy, carbamoyl, N -(C_{1-6} alkyl)carbamoyl, N -(C_{1-6} alkyl)₂carbamoyl, sulphamoyl, N - C_{1-6} alkylaminosulphonyl, N -(C_{1-6} alkyl)₂aminosulphonyl, C_{1-6} alkylsulphonylamino,
- 25 C_{1-6} alkanoylamino, C_{1-6} alkylsulphanyl, C_{1-6} alkylsulphinyl and C_{1-6} alkylsulphonyl.

Preferably the $R^1R^2NSO_2$ - group is para to Y-Z.

In one aspect of the invention R^3 and R^4 are independently C_k alkyl optionally substituted by from 1 to $2k+1$ atoms selected from fluoro and chloro wherein k is 1-3.

Preferably R^3 and R^4 are independently C_k alkyl optionally substituted by from 1 to $2k+1$

30 atoms selected from fluoro and chloro, wherein k is 1-3,

or R^3 and R^4 , together with the carbon atom to which they are attached, form a cyclopropane ring optionally substituted by from 1 to 4 fluorine atoms.

More preferably R^3 and R^4 are independently C_k alkyl optionally substituted by from 1 to $2k+1$ fluorine atoms, wherein k is 1-2,

5 or R^3 and R^4 , together with the carbon atom to which they are attached, form a cyclopropane ring optionally substituted by from 1 to 4 fluorine atoms.

Particularly R^3 and R^4 are independently methyl, fluoromethyl, difluoromethyl, trifluoromethyl, 2,2,2-trifluoroethyl and perfluoroethyl,

10 or R^3 and R^4 , together with the carbon atom to which they are attached, form a cyclopropane ring optionally substituted by from 1 to 4 fluorine atoms.

More particularly R^3 and R^4 are independently methyl, fluoromethyl, difluoromethyl and trifluoromethyl,

or R^3 and R^4 , together with the carbon atom to which they are attached, form a cyclopropane ring optionally substituted by from 1 to 4 fluorine atoms.

15 Preferred combinations of R^3 and R^4 are as follows.

Preferably R^3 and R^4 are both methyl or one of R^3 and R^4 is methyl and the other is trifluoromethyl.

More preferably one of R^3 and R^4 is methyl and the other is trifluoromethyl.

20 Where applicable, the R-configuration generally represents a preferred stereochemistry for compounds of formula (I).

Preferably R^5 is ortho to Y-Z.

Preferably R^5 is selected from halo, nitro, C_{1-4} alkyl, C_{1-4} alkoxy, hydroxy, hydrogen, amino, carboxy and sulphamoyl.

25 More preferably R^5 is selected from fluoro, chloro, bromo, nitro, methyl, ethyl, methoxy, ethoxy, hydroxy, hydrogen, amino, carboxy and sulphamoyl.

Particularly R^5 is selected from fluoro, chloro, nitro and methyl.

More particularly R^5 is selected from fluoro and chloro.

Particularly preferred R^5 is chloro ortho to Y-Z.

30 In another aspect of the invention preferably R^5 is selected from halo, nitro, C_{1-4} alkyl, C_{1-4} alkoxy, C_{2-4} alkenyl, C_{2-4} alkynyl, hydroxy, hydrogen, amino, carboxy and sulphamoyl.

More preferably R^5 is selected from fluoro, chloro, bromo, nitro, methyl, ethyl, methoxy, ethoxy, ethenyl, ethynyl, hydroxy, hydrogen, amino, carboxy and sulphamoyl.

Particularly R^5 is selected from fluoro, chloro, nitro, methyl, ethenyl and ethynyl.

Preferably X is phenyl.


5 Preferably Y-Z is selected from -NHC(O)-, -NHC(S)-, trans-vinylene and ethynylene.

More preferably Y-Z is selected from -NHC(O)- and -NHC(S)-.

Particularly Y-Z is -NHC(O)-.

In a particular embodiment of the present invention there is provided a compound of formula (I) or a pharmaceutically acceptable salt or an *in vivo* hydrolysable ester thereof (as herein
10 before defined). Particular and preferred values are those mentioned above.

According to another aspect of the present invention there is provided a compound of the formula (Ia)



(Ia)

15 wherein

R^{1a} and R^{2a} are independently selected from hydrogen, C_{1-6} alkyl, C_{2-6} alkenyl, C_{3-6} cycloalkyl, C_{1-6} alkyl substituted by a heterocyclic group, C_{1-6} alkyl substituted by phenyl (which phenyl is optionally substituted by one or more halo, trifluoromethoxy, a heterocyclic group or trifluoromethyl) and phenyl which is optionally substituted by one or more halo,

20 C_{1-4} alkoxy, carbamoyl, trifluoromethyl, C_{1-4} alkyl, nitro, hydroxy, cyano, C_{1-6} alkanolyamino, C_{1-6} alkylsulphonyl, C_{1-6} alkanoyl or sulphamoyl,

or R^{1a} and R^{2a} together with the nitrogen group to which they are attached form morpholino or piperazine;

R^{3a} and R^{4a} are independently methyl optionally substituted by from 1 - 3 fluoro;

25 R^{5a} is selected from halo, nitro, C_{1-6} alkyl, C_{1-6} alkoxy, hydroxy, hydrogen, amino, carboxy and sulphamoyl;


G is O or S;

with the proviso that where R^{1a} and R^{2a} are selected from hydrogen, C_{1-3} alkyl or phenyl (which phenyl is optionally substituted by one or two substituents selected from halo, C_{1-4} alkoxy, C_{1-4} alkyl, hydroxy or cyano), R^{5a} is nitro, C_{1-4} alkyl, C_{1-4} alkoxy or hydrogen and R^{3a} and R^{4a} are not both methyl, G must be S;

5 and pharmaceutically acceptable salts or *in vivo* hydrolysable esters thereof.

Where applicable, the R-configuration generally represents a preferred stereochemistry for compounds of formula (Ia).

A further preferred class of compounds is that of formula (Ib):



(Ib)

wherein

R^{1b} and R^{2b} are independently selected from hydrogen, methyl, ethyl, C_3 alkenyl or phenyl which is substituted by one of methoxy, chloro, iodo, hydroxy, carbamoyl, cyano, acetylamino, mesyl, acetyl or sulphamoyl,

15 or R^{1b} and R^{2b} together with the nitrogen group to which they are attached form morpholino.

and pharmaceutically acceptable salts or *in vivo* hydrolysable esters thereof.

In a further aspect of the invention there is provided a compound of formula (Ib) wherein R^{1b} and R^{2b} are independently selected from:

20 i) hydrogen;

ii) C_{1-6} alkyl or C_{3-6} cycloalkyl optionally substituted with one or more C_{1-6} alkyl, C_{1-6} alkoxy, C_{2-6} alkenyloxy, hydroxy, halo, cyano, C_{1-6} alkylsulphanyl, C_{1-6} alkylsulphinyl, C_{1-6} alkylsulphonyl, sulphamoyl, carboxy, C_{1-6} alkoxycarbonyl, amino, C_{1-6} alkylamino, $(C_{1-6}alkyl)_2$ amino, carbamoyl, *N*-(C_{1-6} alkyl)carbamoyl, *N*-(C_{1-6} alkyl)₂carbamoyl, C_{1-6} alkanoylamino,

25 C_{1-6} alkanoyl(*N*- C_{1-6} alkyl)amino, C_{1-6} alkanoyl, C_{2-6} alkenyl, C_{2-6} alkynyl,

N-(C_{1-6} alkyl)aminosulphonyl, hydroxymethyl, hydroxyacetyl or *N*-(C_{1-6} alkyl)₂aminosulphonyl;


iii) a heterocyclic group selected from pyridyl, pyrimidyl, pyridazinyl or pyrazinyl, wherein said heterocyclic group is optionally substituted with one or more C₁₋₆alkyl, C₁₋₆alkoxy, C₂₋₆alkenyloxy, hydroxy, halo, cyano, C₁₋₆alkylsulphanyl, C₁₋₆alkylsulphinyl, C₁₋₆alkylsulphonyl, sulphamoyl, carboxy, C₁₋₆alkoxycarbonyl, amino, C₁₋₆alkylamino, (C₁₋₆alkyl)₂amino, carbamoyl, N-(C₁₋₆alkyl)carbamoyl, N-(C₁₋₆alkyl)₂carbamoyl, C₁₋₆alkanoylamino, C₁₋₆alkanoyl(N-C₁₋₆alkyl)amino, trifluoromethyl, trifluoromethoxy, C₁₋₆alkanoyl, C₂₋₆alkenyl, C₂₋₆alkynyl, N-(C₁₋₆alkyl)aminosulphonyl, hydroxymethyl, hydroxyacetyl or N-(C₁₋₆alkyl)₂aminosulphonyl;

or R^{1b} and R^{2b} together with the nitrogen atom to which they are attached form piperidinyl or piperazinyl; wherein said piperidinyl and piperazinyl may be optionally substituted on a ring carbon by one or more groups selected from C₁₋₆alkyl, C₁₋₆alkoxy, C₂₋₆alkenyloxy, hydroxy, halo, cyano, C₁₋₆alkylsulphanyl, C₁₋₆alkylsulphinyl, C₁₋₆alkylsulphonyl, sulphamoyl, carboxy, C₁₋₆alkoxycarbonyl, amino, C₁₋₆alkylamino, (C₁₋₆alkyl)₂amino, carbamoyl, N-(C₁₋₆alkyl)carbamoyl, N-(C₁₋₆alkyl)₂carbamoyl, C₁₋₆alkanoylamino, C₁₋₆alkanoyl(N-C₁₋₆alkyl)amino, C₁₋₆alkanoyl, C₂₋₆alkenyl, C₂₋₆alkynyl, N-(C₁₋₆alkyl)aminosulphonyl, hydroxymethyl, hydroxyacetyl or N-(C₁₋₆alkyl)₂aminosulphonyl; and said piperazinyl may be optionally substituted on the ring nitrogen by a group selected from C₁₋₆alkanoyl, C₁₋₆alkylsulphonyl, C₁₋₆alkoxycarbonyl, carbamoyl, N-(C₁₋₆alkyl)carbamoyl and N,N-(C₁₋₆alkyl)₂carbamoyl; and wherein any C₁₋₆alkyl group may be optionally substituted by one or more groups selected from hydroxy, amino, C₁₋₆alkylamino, (C₁₋₆alkyl)₂amino, carboxy, C₁₋₆alkoxy, carbamoyl, N-(C₁₋₆alkyl)carbamoyl, N-(C₁₋₆alkyl)₂carbamoyl, sulphamoyl, N-C₁₋₆alkylaminosulphonyl, N-(C₁₋₆alkyl)₂aminosulphonyl, C₁₋₆alkylsulphonylamino, C₁₋₆alkanoylamino, C₁₋₆alkylsulphanyl, C₁₋₆alkylsulphinyl and C₁₋₆alkylsulphonyl.

and pharmaceutically acceptable salts or *in vivo* hydrolysable esters thereof.

Where applicable, the R-configuration generally represents a preferred stereochemistry for compounds of formula (Ib).

A further preferred class of compounds is that of formula (Ic):



wherein:

K is methoxy, hydroxy, carbamoyl, cyano, acetyl, mesyl, or sulphamoyl.

5 and pharmaceutically acceptable salts or *in vivo* hydrolysable esters thereof.

Where applicable; the R-configuration generally represents a preferred stereochemistry for compounds of formula (Ic).

Preferred compounds having formula (I) are Examples 1-37, 61-94 and 270 and pharmaceutically acceptable salts or *in vivo* hydrolysable esters thereof.

10 Most preferred compounds having formula (I) are Examples 17; 34, 35, 36, 62, 78, 81 and 270 and pharmaceutically acceptable salts or *in vivo* hydrolysable esters thereof.

In another aspect of the invention, preferred compounds of the invention are any one of Examples 1-272 and pharmaceutically acceptable salts or *in vivo* hydrolysable esters thereof.

15 In a further aspect of the invention preferred compounds of the invention are Examples 27, 43, 44, 123, 143, 144, 145, 150, 166, 251, 252, 253, 255, 258, 259, 263 and 261 and pharmaceutically acceptable salts or *in vivo* hydrolysable esters thereof.

In an additional aspect of the invention preferred compounds of the invention are Examples 143, 145, 251, 252 and 258 and pharmaceutically acceptable salts or *in vivo* hydrolysable esters thereof.

20 Preferred aspects of the invention are those which relate to the compound or a pharmaceutically acceptable salt thereof.

Within the present invention it is to be understood that a compound of the formula (I) or a salt thereof may exhibit the phenomenon of tautomerism and that the formulae drawings within this specification can represent only one of the possible tautomeric forms. It is to be understood
25 that the invention encompasses any tautomeric form which elevates PDH activity and is not to be limited merely to any one tautomeric form utilized within the formulae drawings. The formulae

drawings within this specification can represent only one of the possible tautomeric forms and it is to be understood that the specification encompasses all possible tautomeric forms of the compounds drawn not just those forms which it has been possible to show graphically herein.


It will be appreciated by those skilled in the art that certain compounds of formula (I) contain one or more asymmetrically substituted carbon and/or sulphur atoms, and accordingly may exist in, and be isolated as enantiomerically pure, a mixture of diastereoisomers or as a racemate-. Some compounds may exhibit polymorphism. It is to be understood that the present invention encompasses any racemic, optically-active, enantiomerically pure, mixture of diastereoisomers, polymorphic or stereoisomeric form, or mixtures thereof, which form possesses properties useful in the elevation of PDH activity, it being well known in the art how to prepare optically-active forms (for example, by resolution of the racemic form by recrystallisation techniques, by synthesis from optically-active starting materials, by chiral synthesis, by enzymatic resolution, (for example WO 9738124), by biotransformation, or by chromatographic separation using a chiral stationary phase) and how to determine efficacy for the elevation of PDH activity by the standard tests described hereinafter.

It is also to be understood that certain compounds of the formula (I) and salts thereof can exist in solvated as well as unsolvated forms such as, for example, hydrated forms. It is to be understood that the invention encompasses all such solvated forms which elevate PDH activity.

A compound of the formula (I), or salt thereof, and other compounds of the invention (as hereinafter defined) may be prepared by any process known to be applicable to the preparation of chemically-related compounds. Such processes include, for example, those illustrated in European Patent Applications, Publication Nos. 0524781, 0617010, 0625516, and in GB 2278054, WO 9323358 and WO 9738124.

Another aspect of the present invention provides a process for preparing a compound of formula (I) or a pharmaceutically acceptable salt or an *in vivo* hydrolysable ester thereof, which process (in which variable groups are as defined for formula (I) unless otherwise stated) comprises of:


(a) deprotecting a protected compound of formula (II)



(II)


where Pg is an alcohol protecting group;

- (b) for a compound of formula (I) in which Y-Z is -NHC(O)-, by coupling an aniline of formula
5 (III):



(III)

with an acid of formula (IV):



(IV)


10

wherein G is a hydroxyl group;

- (c) by coupling an aniline of formula (III) with an activated acid derivative of formula (IV)


wherein G is a hydroxyl group which may be protected as an ester or ether;

- (d) for a compound of formula (I) in which Y-Z is ethynylene, by reacting a alkyne of formula
15 (V):



(V)


with a base, followed by treatment with a ketone of formula (VI):



(VI);


(e) for a compound of formula (I) in which Y-Z is *trans*-vinylene, by reducing a compound of
 5 formula (I) in which Y-Z is ethynylene;

(f) for a compound of formula (I) in which Y-Z is *trans*-vinylene, by dehydration of a diol of
 formula (VII):



(VII);


10 (g) for a compound of formula (I) in which Y-Z is *trans*-vinylene, by base catalysed opening of
 an epoxide of formula (VIII):



(VIII);


(h) for a compound of formula (I) in which Y-Z is -NHCH₂-, by reducing a compound of formula
 15 (I) in which Y-Z is -NHC(O)-;

(i) for a compound of formula (I) in which Y-Z is -OCH₂-, -SCH₂- or -NHCH₂- by reacting an
 ethylene oxide of formula (IX):



(IX)


20 with a compound of formula (III) or a compound of formula (X):



(X)

where J is -OH, -NH₂ or -SH;

(j) by reacting a compound of formula (XI):



(XI)

where K is a leaving atom or group, and in which Y-Z is OCH₂, SCH₂ or NHCH₂ or -NHC(O)- with an amine of formula R¹R²NH;

(k) for a compound of formula (I) in which Y-Z is -NHC(S)-, by reacting a compound of formula

10 (I) in which Y-Z is -NHC(O)- with a sulphur reagent;

and thereafter if necessary:

i) converting a compound of the formula (I) into another compound of the formula (I);

ii) removing any protecting groups; or

iii) forming a pharmaceutically acceptable salt or *in vivo* hydrolysable ester.

15 Suitable values for Pg are a benzyl group, a silyl group or an acetyl protecting group.

K is a leaving atom or group, suitable values for K are for example a halogen atom such as fluoro or chloro.

Specific conditions of the above reactions are as follows:

a) Suitable reagents for deprotecting an alcohol of formula (II) are for example:

20 1) when Pg is benzyl:

(i) hydrogen in the presence of palladium/carbon catalyst, i.e. hydrogenolysis; or

(ii) hydrogen bromide or hydrogen iodide;

2) when Pg is a silyl protecting group:

(i) tetrabutylammonium fluoride; or

25 (ii) aqueous hydrofluoric acid;

3) when Pg is acetyl:

i) mild aqueous base for example lithium hydroxide.

The reaction can be conducted in a suitable solvent such as ethanol, methanol, acetonitrile, or dimethylsulphoxide and may conveniently be performed at a temperature in the range of -40 to 5 100°C.

(b) An aniline of formula (III) and an acid of formula (IV) may be coupled together in the presence of a suitable coupling reagent. Standard peptide coupling reagents known in the art can be employed as suitable coupling reagents, for example thionyl chloride (or oxalyl chloride), carbonyldiimidazole and dicyclohexyl-carbodiimide, optionally in the presence of a catalyst such 10 as dimethylaminopyridine or 4-pyrrolidinopyridine, optionally in the presence of a base for example triethylamine, pyridine, or 2,6-di-*alkyl*-pyridines such as 2,6-lutidine or 2,6-di-*tert*-butylpyridine. Suitable solvents include dimethylacetamide, dichloromethane, benzene, tetrahydrofuran, and dimethylformamide. The coupling reaction may conveniently be performed at a temperature in the range of -40 to 40°C.

15 (c) An aniline of formula (III) may be coupled with an activated acid derivative of formula (IV) for example acid chlorides, acid anhydrides, or phenyl esters, wherein G is a hydroxyl group which may be suitably protected as a stable ester or ether. This coupling may be achieved optionally in the presence of a base for example triethylamine, pyridine, or 2,6-di-*alkyl*-pyridines such as 2,6-lutidine or 2,6-di-*tert*-butylpyridine. Suitable solvents include dimethylacetamide, 20 dichloromethane, benzene, tetrahydrofuran, and dimethylformamide. The coupling reaction may conveniently be performed at a temperature in the range of -40 to 40°C;

(d) suitable bases for reacting with a corresponding alkyne of formula (V) are for example lithium diisopropylamide (LDA), *n*-butyllithium or *tert*-butyllithium. The reaction with a ketone of formula (IV) may be performed at a temperature in the range of -100 to -40°C preferably at a 25 temperature in the range of -70 to -40°C and in a solvent such as tetrahydrofuran, diethyl ether, or 1,2-dimethoxyethane.

(e) A suitable reducing agent for a compound of formula (I) in which Y-Z is *trans*-vinylene is, for example, lithium aluminium hydride or sodium bis(methoxyethoxy)aluminium hydride. The reaction can be conducted in a suitable solvent such as tetrahydrofuran or diethyl ether, and at a 30 temperature in the range of 0 to 50°C.

(f) Dehydration of a diol of formula (VII) may be conducted in the presence of an acid catalyst (for example *p*-toluenesulphonic acid), neat or with a solvent such as toluene or dichloromethane at a temperature in the range of 0 to 200°C preferably a temperature in the range of 20 to 100°C.

(g) Base catalysed opening of an epoxide of formula (VIII) may be carried out in a suitable organic solvent for example, ethers or toluene. Ethers such as tetrahydrofuran are preferred. Suitable bases include potassium *tert*-butoxide or sodium hydride. The opening may be carried out at a temperature in the range of -50 to 100°C, preferably at a temperature in the range of 0 to 50°C for example room temperature.

(h) A compound of formula (I) in which Y-Z is -NHC(O)- may be reduced with a suitable reducing agent such as lithium aluminium hydride or borane. The reaction can conveniently be carried out at a temperature in the range of 0°C to reflux, in solvents such as for example diethyl ether, tetrahydrofuran, or 1,2-dimethoxyethane.

(i) An ethylene oxide of formula (IX) may be reacted with a corresponding compound of formula (III) or a compound of formula (X) in the presence of a base for example sodium hydride or triethylamine. The reaction can be conducted at reflux in a solvent such as dichloromethane, tetrahydrofuran, or diethyl ether.

(j) A compound of formula (XI) where K is a leaving atom or group, for example a halogen atom such as fluoro or chloro and in which Y-Z is OCH₂, SCH₂ or NHCH₂ or -NHC(O)- may be reacted with an amine of formula R¹R²NH in the presence of a base, for example a tertiary amine such as triethylamine and in the presence of a catalyst for example dimethylaminopyridine. Suitable solvents for the reaction include nitriles such as acetonitrile and amides such as dimethylformamide. The reaction is conveniently performed at a temperature in the range of from 0 to 120°C.

(k) A compound of formula (I) in which Y-Z is -NHC(O)- may be reacted with a reagent such as for example phosphorus pentasulphide or Lawesson's reagent (2,4-bis(4-methoxyphenyl)-1,3-dithia-2,4-diphosphetane-2,4-disulphide), optionally in the presence of a suitable base such as for example pyridine or triethylamine. Suitable solvents for the reaction include for example toluene, tetrahydrofuran, 1,3-dioxane or acetonitrile. The reaction is conveniently performed at a temperature in the range of from 0 to reflux.


If not commercially available, the necessary starting materials for the procedures such as that described above may be made by procedures which are selected from standard organic chemical techniques, techniques which are analogous to the synthesis of known, structurally similar compounds, or techniques which are analogous to the above described procedure or the
5 procedures described in the examples.

For example, it will be appreciated that certain of the optional aromatic substituents in the compounds of the present invention may be introduced by standard aromatic substitution reactions or generated by conventional functional group modifications either prior to or immediately following the processes mentioned above, and as such are included in the process
10 aspect of the invention. Such reactions and modifications include, for example, introduction of a substituent by means of an aromatic substitution reaction, reduction of substituents, alkylation of substituents and oxidation of substituents. The reagents and reaction conditions for such procedures are well known in the chemical art. Particular examples of aromatic substitution reactions include the introduction of a nitro group using concentrated nitric acid, the introduction
15 of an acyl group using, for example, an acylhalide and Lewis acid (such as aluminium trichloride) under Friedel Crafts conditions; the introduction of an alkyl group using an alkyl halide and Lewis acid (such as aluminium trichloride) under Friedel Crafts conditions; and the introduction of a halogeno group. Particular examples of modifications include the reduction of a nitro group to an amino group by, for example, catalytic hydrogenation with a nickel catalyst or treatment
20 with iron in the presence of hydrochloric acid with heating; oxidation of alkylthio to alkylsulphinyl or alkylsulphonyl using, for example, hydrogen peroxide in acetic acid with heating or 3-chloroperbenzoic acid.

Specific examples of the techniques used to make that starting materials described above are illustrated, but not limited by, the following examples in which variable groups are as defined
25 for formula (I) unless otherwise stated.

1) Preparation of compounds of formula (II).

a) compounds of formula (II) in which Y-Z is OCH₂, SCH₂ or NHCH₂ may be made by treating the corresponding compound of formula (X) wherein J is -OH, -SH, -NH₂ or a compound of
30 formula (III) with a compound of formula (XII):



(XII)

where Y is a leaving group for example mesylate; in the presence of a base such as an alkali metal hydride (e.g. sodium hydride), in a solvent such as tetrahydrofuran,


5 *N,N*-dimethylformamide, dimethyl sulphoxide, or

1,3-Dimethyl-3,4,5,6-tetrahydro-2(1*H*)-pyrimidinone, and at a temperature of 20°C to reflux.

b) A compound of formula (II), wherein Y-Z is -NHC(O)-, may be made by coupling a compound of formula (III) with a compound of formula (IV) (where G is hydroxy protected with a protecting group) in a manner analogous to that described for procedure (b) of preparations of a

10 compound of formula (I) above.

Compounds of formula (IV) where G is hydroxy protected with a protecting group may be made by conventional procedures. For example, cleavage of the ester group of a compound of formula (XIII):



(XIII)


where E is a carboxy protecting group (e.g. Me);

under standard conditions such as mild alkaline conditions, for example, aqueous lithium hydroxide.

Compounds of formula (XIII) where G is protected hydroxy are prepared by protecting a
 20 compound of formula (XIII) where G is hydroxy by reaction with a compound such as benzyl chloride or benzyl bromide (in the presence of a suitable base such as sodium hydride and optionally with a catalyst such as sodium iodide, to provide a benzyl protecting group) or any of the conventional silylating agents known and used for such purpose (for example 2-trimethylsilylethoxymethyl chloride, in the presence of a suitable base such as triethylamine
 25 optionally in the presence of a catalyst such as dimethylaminopyridine).


Compounds of formula (XIII) where G is hydroxy are prepared by esterifying an acid of formula (IV) by a conventional esterification procedure such as reaction with a C₁₋₆ alcohol (e.g. methanol) in the presence of an acid catalyst (for example sulphuric acid).

- c) A compound of formula (II), wherein Y-Z is ethynylene, may be made by reacting a
5 compound of formula (XIV):



(XIV)

wherein L is a leaving group such as bromo, iodo, or triflate, with an acetylene of formula (XV)




(XV)

in the presence of a catalyst such as a combination of copper (I) iodide and bis(triphenylphosphine)palladium dichloride or palladium (II) acetate. The reaction can be conducted in an inert solvent such as tetrahydrofuran, benzene, or toluene, or in a basic solvent

- 15 such as diethylamine or triethylamine, and at a temperature in the range of -20 to 110°C.

A compound of formula (XV) may be made by reacting a compound of formula (XVI)




(XVI)

with an agent such as:

- 20 i) benzyl bromide (to provide a benzyl protecting group), this reaction may conveniently be conducted in the presence of a base such as sodium hydride and optionally in the presence of a catalyst such as sodium iodide in a solvent such as tetrahydrofuran at a temperature of about -78 to about 100 °C; or

ii) any of the conventional silylating agents known and used for such purpose (such as for example *tert*-butyl dimethylsilylchloride or triflate, in the presence of a suitable base such as 1,8-Diazabicyclo[5.4.0]undec-7-ene or triethylamine optionally in the presence of a catalyst such as dimethylaminopyridine) at a temperature of about -78 to about 100 °C.

- 5 d) A compound of formula (II), wherein Y-Z is *trans*-vinylene, may be made by reacting a compound of formula (XVII):



(XVII)

- where M is an alkylmetal group such as a trialkyltin (for example tributyl- or trimethyl-tin) or a
 10 bisalkoxyborane (for example catecholborane);
 with a compound of formula (X), wherein J may be a leaving group for example iodide, bromide or triflate in the presence of a catalyst such as bis(triphenylphosphine)palladium dichloride or tetrakis(triphenylphosphine)palladium (0). The reaction may conveniently be conducted in a suitable inert solvent such as a tetrahydrofuran or dimethylformamide at a temperature of 0 -
 15 150°C.

A compound of formula (XVII) may be made by a reaction of a compound of formula (XV)

- i) with an agent such as catecholborane, to form the vinylborane compound; or
 ii) a trialkyltinhydride in the presence of a catalytic amount of a radical chain initiator such as, for
 20 example, *aza-bis*-isobutyronitrile or by using trialkyltinhydride pre-treated with a strong base (such as an alkyllithium) and copper (I) cyanide, or by using a transition metal catalyst such as, for example, tetrakis(triphenylphosphine)palladium(0) to form a compound of formula (XVII) where M is trialkyltin.

- These reactions may conveniently be conducted in a suitable inert solvent such as
 25 tetrahydrofuran, toluene or xylene at a temperature of from 0 - 150°C.


Compounds of formula (XVI) may be made by reacting a compound of formula (VI) with an alkali metal acetylide (for example lithium acetylide) or alkaline earth metal acetylide (for example magnesium acetylide). The reaction may be conducted in a solvent such as

tetrahydrofuran, diethyl ether, or 1,2-dimethoxyethane and at a temperature of -100 to 25 °C.

2) Preparation of compounds of formula (III).

A compound of formula (III) may be prepared:


- 5 i) from a compound of formula (XVIII)



(XVIII)

wherein Pg is a protective group such as for example acetyl;


- a) by treatment with chlorosulphonic acid under standard conditions, and then
 10 b) formation of the sulphonamide under standard conditions as described above in process (j) for preparation of a compound of formula (I) and then
 c) cleavage of the protecting group under mild alkaline conditions (for example when Pg is acetyl with a base such as aqueous sodium hydroxide); or
 ii) by reducing a compound of formula (XIX):



(XIX)

under standard conditions for example by a reducing agent such as tin (II) chloride or iron dust in conjunction with concentrated acid to give a compound of formula (III).

A compound of formula (XIX) may be made by reacting a compound of formula (XX):




(XX)

with an amine of formula R^1R^2NH- in a procedure analogous to that used in process (j) for preparation of a compound of formula (I) above.

A compound of formula (XX) may be prepared:


a) by oxidising a compound of formula (XXI):



(XXI)

under standard conditions for example with chlorine in a suitable solvent such as acetic acid at a temperature of -78 to about 100°C; or

b) by diazotizing a compound of formula (XXII):



(XXII)

under standard conditions for example with nitrous acid and sulphuric acid followed by reaction with a mixture of sulphur dioxide and copper (II) chloride in a suitable solvent such as water or a water/acetic acid solution.

3) Resolution of compounds of formula (IV)

If the resolved acid is required it may be prepared by any of the known methods for preparation of optically-active forms (for example, by recrystallisation of the chiral salt {for example WO 9738124}, by enzymatic resolution, by biotransformation, or by chromatographic separation using a chiral stationary phase). For example if an (R)-(+)-resolved acid is required it may be prepared by the method of Scheme 2 in World Patent Application Publication No. WO 9738124 for preparation of the (S)-(-) acid, i.e. using the classical resolution method described in European Patent Application Publication No. EP 0524781, also for preparation of the (S)-(-) acid, except that (1S,2R)-norephedrine may be used in place of (S)-(-)-1-phenylethylamine.


4) Preparation of compounds of formula (V).

A compound of formula (V) may be prepared by reacting a compound of formula (XIV), wherein L is bromo, iodo or triflate with trimethylsilylacetylene in the presence of a catalyst such

- 5 as a combination of bis(triphenylphosphine)palladium dichloride and copper(I) iodide in diethylamine or triethylamine, followed by treatment with a base (for example potassium carbonate) in a C₁₋₆ alcohol (such as methanol) as the solvent to remove the trimethylsilyl group.

5) Preparation of compounds of formula (VII).


- 10 A compound of formula (VII) may be prepared from a compound of formula (XXIII):



(XXIII)


by reduction under standard conditions for example by using a hydride, such as sodium borohydride.

- 15 A compound of formula (XXIII) may be prepared by deprotonation of a compound of formula (XXIV),



(XXIV)

- with a strong base, for example lithium diisopropyl amide in an organic solvent, for example tetrahydrofuran at a temperature of -78 to 100 °C followed by addition of an amide of formula (XXV):
- 20



(XXV)

in which R^{19} and R^{20} are each independently C_{1-6} alkyl or together with the atoms to which they are
5 attached form a 5-7 membered ring.

An amide of formula (XXV) may be prepared from an acid of formula (IV), or a reactive derivative thereof, by reaction with an amine of formula $R^{19}(R^{20}O)NH$ under standard conditions such as those described in process (b) for preparation of a compound of formula (I) above.

6) Preparation of compounds of formula (VIII).

10 A compound of formula (VIII) may be prepared from a diol of formula (VII) using a suitable dehydrating agent, for example
bis[α,α -bis(trifluoromethyl)benzenemethanolato]diphenyl sulphur.

7) Preparation of compounds of formula (IX).

15 A compound of formula (IX) may be made by treating a compound of formula (VI) with a trimethylsulphonium salt (such as trimethylsulphonium iodide) and a base (such as an alkali metal hydroxide) in a solvent such as dichloromethane.

8) Preparation of compounds of formula (X).


20 a) A compound of formula (X) wherein J is -OH, may be prepared by diazotizing a compound of formula (III) under standard conditions such as those described above in 2(ii) above followed by heating the resulting compound in dilute sulphuric acid.
b) A compound of formula (X), wherein J is -SH, may be prepared by reacting a compound of formula (XIV) where L is a leaving group (for example chloro) with an excess of methanethiol in
25 the presence of sodium hydride.

9) Preparation of compounds of formula (XI).

A compound of formula (XI) wherein K is chloro, in which Y-Z is OCH₂, SCH₂, NHCH₂ or -NHC(O)- may be prepared by

1) either

a) coupling a compound of formula (XXVI)



(XXVI)

wherein J is -OH, -SH or NH₂ with a compound of formula (XII) where Y is a leaving group for example mesylate; in the presence of a base such as an alkali metal hydride (e.g. sodium hydride), in a solvent such as tetrahydrofuran, *N,N*-dimethylformamide, dimethyl sulphoxide, or

10 1,3-Dimethyl-3,4,5,6-tetrahydro-2(1*H*)-pyrimidinone, and at a temperature of 20°C to reflux; or


b) where Y-Z is -NHC(O)- coupling with a compound of formula (XXVI) where J is NH₂ with a compound of formula (IV), following a method analogous to that of process (b) for preparation of a compound of formula (I) above.

Route a or b is then followed by:

15 2) treatment with chlorosulphonic acid.

10) Preparation of compounds of formula (XII).

A compound of formula (XII), wherein Y is mesylate may be prepared by reacting a compound of formula (XXVII):



(XXVII)

with methanesulphonic acid chloride in the presence of a base such as triethylamine, in a solvent such as dichloromethane, and at a temperature of about -78 to 25°C.

Compounds of formula (XXVII) are prepared by reducing a compound of formula (XIII)

25 with a suitable reducing agent such as lithium aluminium hydride in a solvent such as diethyl

ether or THF and at a temperature of about 0 to about 25 °C.

It is noted that many of the starting materials for synthetic methods as described above are commercially available and/or widely reported in the scientific literature, or could be made from commercially available compounds using adaptations of processes reported in the scientific
5 literature.

It will also be appreciated that in some of the reactions mentioned herein it may be necessary/desirable to protect any sensitive groups in the compounds. The instances where protection is necessary or desirable and suitable methods for protection are known to those skilled in the art. Thus, if reactants include groups such as amino, carboxy or hydroxy it may be
10 desirable to protect the group in some of the reactions mentioned herein.

A suitable protecting group for an amino or alkylamino group is, for example, an acyl group, for example an alkanoyl group such as acetyl, an alkoxycarbonyl group, for example a methoxycarbonyl, ethoxycarbonyl or *t*-butoxycarbonyl group, an arylmethoxycarbonyl group, for example benzyloxycarbonyl, or an aroyl group, for example benzoyl. The deprotection conditions
15 for the above protecting groups necessarily vary with the choice of protecting group. Thus, for example, an acyl group such as an alkanoyl or alkoxycarbonyl group or an aroyl group may be removed for example, by hydrolysis with a suitable base such as an alkali metal hydroxide, for example lithium or sodium hydroxide. Alternatively an acyl group such as a *t*-butoxycarbonyl group may be removed, for example, by treatment with a suitable acid as hydrochloric, sulphuric
20 or phosphoric acid or trifluoroacetic acid and an arylmethoxycarbonyl group such as a benzyloxycarbonyl group may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon, or by treatment with a Lewis acid for example boron tris(trifluoroacetate). A suitable alternative protecting group for a primary amino group is, for example, a phthaloyl group which may be removed by treatment with an alkylamine, for example
25 dimethylaminopropylamine, or with hydrazine.

A suitable protecting group for a hydroxy group is, for example, an acyl group, for example an alkanoyl group such as acetyl, an aroyl group, for example benzoyl, or an arylmethyl group, for example benzyl. The deprotection conditions for the above protecting groups will necessarily vary with the choice of protecting group. Thus, for example, an acyl group such as an
30 alkanoyl or an aroyl group may be removed, for example, by hydrolysis with a suitable base such

as an alkali metal hydroxide, for example lithium or sodium hydroxide. Alternatively an arylmethyl group such as a benzyl group may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon.

A suitable protecting group for a carboxy group is, for example, an esterifying group, for example a methyl or an ethyl group which may be removed, for example, by hydrolysis with a base such as sodium hydroxide, or for example a *t*-butyl group which may be removed, for example, by treatment with an acid, for example an organic acid such as trifluoroacetic acid, or for example a benzyl group which may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon.

10 The protecting groups may be removed at any convenient stage in the synthesis using conventional techniques well known in the chemical art.

In cases where compounds of formula (I) are sufficiently basic or acidic to form stable acid or basic salts, administration of the compound as a salt may be appropriate, and pharmaceutically acceptable salts may be made by conventional methods such as those described following. Examples of suitable pharmaceutically acceptable salts are organic acid addition salts formed with acids which form a physiologically acceptable anion, for example, tosylate, methanesulphonate, acetate, tartrate, citrate, succinate, benzoate, ascorbate, α -ketoglutarate, and α -glycerophosphate. Suitable inorganic salts may also be formed such as sulphate, nitrate, and hydrochloride.

20 Pharmaceutically acceptable salts may be obtained using standard procedures well known in the art, for example by reacting a sufficiently basic compound of formula (I) (or its ester) with a suitable acid affording a physiologically acceptable anion. It is also possible with most compounds of the invention to make a corresponding alkali metal (e.g. sodium, potassium, or lithium) or alkaline earth metal (e.g. calcium) salt by treating a compound of formula (I) (and in some cases the ester) with one equivalent of an alkali metal or alkaline earth metal hydroxide or alkoxide (e.g. the ethoxide or methoxide) in aqueous medium followed by conventional purification techniques.

In vivo cleavable esters of compounds of the invention may be made by coupling with a pharmaceutically acceptable carboxylic acid or an activated derivative thereof. For example, the coupling may be carried out by treating a compound of formula (I) with an appropriate acid

chloride (for example, acetyl chloride, propionyl chloride, or benzoyl chloride) or acid anhydride (for example, acetic anhydride, propionic anhydride, or benzoic anhydride) in the presence of a suitable base such as triethylamine. Those skilled in the art will appreciate that other suitable carboxylic acids (including their activated derivatives) for the formation of *in vivo* cleavable esters are known to the art and these are also intended to be included within the scope of the invention. Catalysts such as 4-dimethylaminopyridine may also be usefully employed.

Many of the intermediates defined herein are novel and these are provided as a further feature of the invention.

The identification of compounds which elevate PDH activity is the subject of the present invention. These properties may be assessed, for example, using one or more of the procedures set out below:

(a) In vitro elevation of PDH activity

This assay determines the ability of a test compound to elevate PDH activity. cDNA encoding PDH kinase may be obtained by Polymerase Chain Reaction (PCR) and subsequent cloning. This may be expressed in a suitable expression system to obtain polypeptide with PDH kinase activity. For example rat PDHkinaseII (rPDHKII) obtained by expression of recombinant protein in *Escherichia coli* (*E. Coli*), was found to display PDH kinase activity.

In the case of the rPDHKII (Genbank accession number U10357) a 1.3kb fragment encoding the protein was isolated by PCR from rat liver cDNA and cloned into a vector (for example pQE32 - Quiagen Ltd.). The recombinant construct was transformed into *E. coli* (for example M15pRep4 - Quiagen Ltd.). Recombinant clones were identified, plasmid DNA was isolated and subjected to DNA sequence analysis. One clone which had the expected nucleic acid sequence was selected for the expression work. Details of the methods for the assembly of recombinant DNA molecules and the expression of recombinant proteins in bacterial systems can be found in standard texts for example Sambrook et al, 1989, *Molecular Cloning - A Laboratory Manual*, 2nd edition, Cold Spring Harbour Laboratory Press. Other known PDH kinases for use in assays, may be cloned and expressed in a similar manner.

For expression of rPDHKII activity, *E. coli* strain M15pRep4 cells were transformed with the pQE32 vector containing rPDHKII cDNA. This vector incorporates a 6-His tag onto the protein at its N-terminus. *E. coli* were grown to an optical density of 0.6 (600 nM) and protein

expression was induced by the addition of 10 μ M isopropylthio- β -galactosidase. Cells were grown for 18 hours at 18°C and harvested by centrifugation. The resuspended cell paste was lysed by homogenisation and insoluble material removed by centrifugation at 24000xg for 1 hour. The 6-His tagged protein was removed from the supernatant using a nickel chelating
5 nitrilotriacetic acid resin (Ni-NTA: Quiagen Ltd.) matrix (Quiagen) which was washed with 20 mM tris(hydroxymethyl)aminomethane-hydrogen chloride, 20 mM imidazole, 0.5 M sodium chloride pH 8.0; prior to elution of bound protein using a buffer containing 20 mM tris(hydroxymethyl)aminomethane-hydrogen chloride, 200 mM imidazole, 0.15 M sodium
10 chloride pH 8.0. Eluted fractions containing 6-His protein were pooled and stored in aliquots at -80°C in 10% glycerol.

Each new batch of stock enzyme was titrated in the assay to determine a concentration giving approximately 90% inhibition of PDH in the conditions of the assay. For a typical batch, stock enzyme was diluted to 7.5 μ g/ml.

For assay of the activity of novel compounds, compounds were diluted with 10%
15 dimethylsulphoxide (DMSO) and 10 μ l transferred to individual wells of 96-well assay plates. Control wells contained 20 μ l 10% DMSO instead of compound. 40 μ l Buffer containing 50mM potassium phosphate buffer pH 7.0, 10mM ethylene glycol-bis(β -aminoethyl ether)-N,N,N,N-tetracetic acid (EGTA), 1mM benzamidine, 1mM phenylmethylsulphonyl fluoride (PMSF), 0.3mM tosyl-L-lysine chloromethyl ketone (TLCK), 2mM dithiothreitol (DTT), recombinant
20 rPDHKII and compounds were incubated in the presence of PDH kinase at room temperature for 45 minutes. In order to determine the maximum rate of the PDH reaction a second series of control wells were included containing 10% DMSO instead of compound and omitting rPDHKII. PDH kinase activity was then initiated by the addition of 5 μ M ATP, 2 mM magnesium chloride and 0.04 U/ml PDH (porcine heart PDH Sigma P7032) in a total volume of 50 μ l and plates
25 incubated at ambient temperature for a further 45 minutes. The residual activity of the PDH was then determined by the addition of substrates (2.5mM coenzyme A, 2.5mM thiamine pyrophosphate (cocarboxylase), 2.5mM sodium pyruvate, 6mM NAD in a total volume of 80 μ l and the plates incubated for 90 minutes at ambient temperature. The production of reduced NAD (NADH) was established by measured optical density at 340nm using a plate reading
30 spectrophotometer. The ED₅₀ for a test compound was determined in the usual way using results

from 12 concentrations of the compound.

(b) In vitro elevation of PDH activity in isolated primary cells

This assay determines the ability of compounds to stimulate pyruvate oxidation in primary rat hepatocytes.

- 5 Hepatocytes were isolated by the two-step collagenase digestion procedure described by Seglen (Methods Cell Biol. (1976) 13, 29-33) and plated out in 6-well culture plates (Falcon Primaria) at 600000 viable cells per well in Dulbecco's Modified Eagles Medium (DMEM, Gibco BRL) containing 10% foetal calf serum (FCS), 10% penicillin/streptomycin (Gibco BRL) and 10% non-essential amino acids (NEAA, Gibco BRL). After 4 hours incubation at 37°C in 5% CO₂, the medium was replaced with Minimum Essential Medium (MEM, Gibco BRL) containing NEAA and penicillin/streptomycin as above in addition to 10nM dexamethasone and 10nM insulin.

- The following day cells were washed with phosphate buffered saline (PBS) and medium replaced with 1ml HEPES-buffered Krebs solution (25mM HEPES, 0.15M sodium chloride, 25 mM sodium hydrogen carbonate, 5mM potassium chloride, 2mM calcium chloride, 1mM magnesium sulphate, 1 mM potassium dihydrogen phosphate) containing the compound to be tested at the required concentration in 0.1% DMSO. Control wells contained 0.1% DMSO only and a maximum response was determined using a 10 µM treatment of a known active compound. After a preincubation period of 40 minutes at 37°C in 5% CO₂, cells were pulsed with sodium pyruvate to a final concentration of 0.5mM (containing 1-¹⁴C sodium pyruvate (Amersham product CFA85) 0.18Ci/mmol) for 12 minutes. The medium was then removed and transferred to a tube which was immediately sealed with a bung containing a suspended centre well. Absorbent within the centre well was saturated with 50% phenylethylamine, and CO₂ in the medium released by the addition of 0.2µl 60% (w/v) perchloric acid (PCA). Released ¹⁴CO₂ trapped in the absorbent was determined by liquid scintillation counting. The ED₅₀ for a test compound was determined in the usual way using results from 7 concentrations of the compound.
- (c) In vivo elevation of PDH activity

- The capacity of compounds to increase the activity of PDH in relevant tissues of rats may be measured using the test described hereinafter. Typically an increase in the proportion of PDH in its active, nonphosphorylated form may be detected in muscle, heart, liver and adipose tissue

after a single administration of an active compound. This may be expected to lead to a decrease in blood glucose after repeated administration of the compound. For example a single administration of DCA, a compound known to activate PDH by inhibition of PDH kinase (Whitehouse, Cooper and Randle (1974) *Biochem. J.* 141, 761-774) 150 mg/kg, intraperitoneally, increased the proportion of PDH in its active form (Vary et al. (1988) *Circ. Shock* 24, 3-18) and after repeated administration resulted in a significant decrease in plasma glucose (Evans and Stacpoole (1982) *Biochem. Pharmacol.* 31, 1295-1300).

Groups of rats (weight range 140-180g) are treated with a single dose or multiple doses of the compound of interest by oral gavage in an appropriate vehicle. A control group of rats is treated with vehicle only. At a fixed time after the final administration of compound, animals are terminally anaesthetised, tissues are removed and frozen in liquid nitrogen. For determination of PDH activity, muscle samples are disrupted under liquid nitrogen prior to homogenisation by one thirty-second burst in a Polytron homogenizer in 4 volumes of a buffer containing 40 mM potassium phosphate pH 7.0, 5 mM EDTA, 2mM DTT, 1% Triton X-100, 10mM sodium pyruvate, 10µM phenylmethylsulphonyl chloride (PMSF) and 2µg/ml each of leupeptin, pepstatin A and aprotinin. Extracts are centrifuged before assay. A portion of the extract is treated with PDH phosphatase prepared from pig hearts by the method of Siess and Wieland (*Eur. J. Biochem* (1972) 26, 96): 20 µl extract, 40 µl phosphatase (1:20 dilution), in a final volume of 125 µl containing 25 mM magnesium chloride, 1 mM calcium chloride. The activity of the untreated sample is compared with the activity of the dephosphorylated extract thus prepared. PDH activity is assayed by the method of Stansbie et al., (*Biochem. J.* (1976) 154, 225). 50 µl Extract is incubated with 0.75 mM NAD, 0.2 mM CoA, 1.5 mM thiamine pyrophosphate (TPP) and 1.5mM sodium pyruvate in the presence of 20 µg/ml p-(p-amino-phenylazo) benzene sulphonic acid (AABS) and 50 mU/ml arylamine transferase (AAT) in a buffer containing 100 mM tris(hydroxymethyl)aminomethane, 0.5 mM EDTA, 50mM sodium fluoride, 5mM 2-mercaptoethanol and 1mM magnesium chloride pH 7.8. AAT is prepared from pigeon livers by the method of Tabor et al. (*J. Biol. Chem.* (1953) 204, 127). The rate of acetyl CoA formation is determined by the rate of reduction of AABS which is indicated by a decrease in optical density at 460 nm.

Liver samples are prepared by an essentially similar method, except that sodium pyruvate is excluded from the extraction buffer and added to the phosphatase incubation to a final concentration of 5mM.

Treatment of an animal with an active compound results in an increase in the activity of
5 PDH complex in tissues. This is indicated by an increase in the amount of active PDH (determined by the activity of untreated extract as a percentage of the total PDH activity in the same extract after treatment with phosphatase).

According to a further aspect of the invention there is provided a pharmaceutical composition which comprises a compound of the formula (I) as defined hereinbefore or a
10 pharmaceutically acceptable salt or an *in vivo* hydrolysable ester thereof, in association with a pharmaceutically acceptable excipient or carrier.

According to a further aspect of the invention there is provided a pharmaceutical composition which comprises a compound of the formula (I) which is selected from:
i) a compound of the formula (I) wherein R³ and R⁴ are both methyl, R⁵ is hydrogen, fluoro or
15 chloro, Y-Z is ethynylene, X is phenyl and one of R¹ and R² is hydrogen and the other is pyrimidyl-NH-C(O)- or triazinyl-NH-C(O)- (wherein said triazine or pyrimidine is substituted by methyl, methoxy or dimethylamino) and the -SO₂NR¹R² moiety is ortho to Y-Z;
ii) 4-(3-hydroxy-3-methyl-1-butynyl)-N-(3-methyl-2-pyridinyl)-benzenesulphonamide;
iii) *N*-{4-[N,N-bis-(*sec*-butyl)aminosulphonyl]phenyl}-2-hydroxy-2-methyl-3,3,3-
20 trifluoropropanamide; or
iv) *N*-{4-[N,N-bis-(*iso*-butyl)aminosulphonyl]phenyl}-2-hydroxy-2-methyl-3,3,3-trifluoropropanamide;
or a pharmaceutically acceptable salt or an *in vivo* hydrolysable ester thereof, in association with a pharmaceutically acceptable excipient or carrier.

25 The composition may be in a form suitable for oral administration, for example as a tablet or capsule, for parenteral injection (including intravenous, subcutaneous, intramuscular, intravascular or infusion) for example as a sterile solution, suspension or emulsion, for topical administration for example as an ointment or cream or for rectal administration for example as a suppository. In general the above compositions may be prepared in a conventional manner using
30 conventional excipients.

The compositions of the present invention are advantageously presented in unit dosage form. The compound will normally be administered to a warm-blooded animal at a unit dose within the range 5-5000 mg per square metre body area of the animal, i.e. approximately 0.1-100 mg/kg. A unit dose in the range, for example, 1-100 mg/kg, preferably 1-50 mg/kg is envisaged and this normally provides a therapeutically-effective dose. A unit dose form such as a tablet or capsule will usually contain, for example 1-250 mg of active ingredient.

According to a further aspect of the present invention there is provided a compound of the formula (I) or a pharmaceutically acceptable salt or an *in vivo* hydrolysable ester thereof as defined hereinbefore for use in a method of treatment of the human or animal body by therapy.

10 We have found that compounds of the present invention elevate PDH activity and are therefore of interest for their blood glucose-lowering effects.

A further feature of the present invention is a compound of formula (I) and pharmaceutically acceptable salts or *in vivo* hydrolysable esters thereof for use as a medicament.

Conveniently this is a compound of formula (I), or a pharmaceutically acceptable salt or an *in vivo* hydrolysable ester thereof, for use as a medicament for producing an elevation of PDH activity in a warm-blooded animal such as a human being.

Thus according to a further aspect of the invention there is provided the use of a compound of the formula (I), or a pharmaceutically acceptable salt or an *in vivo* hydrolysable ester thereof in the manufacture of a medicament for use in the production of an elevation of PDH activity in a warm-blooded animal such as a human being.

According to a further feature of the invention there is provided a method for producing an elevation of PDH activity in a warm-blooded animal, such as a human being, in need of such treatment which comprises administering to said animal an effective amount of a compound of formula (I) or a pharmaceutically acceptable salt or an *in vivo* hydrolysable ester thereof as defined hereinbefore.

For the avoidance of doubt, in aspects of the invention concerning the use of compounds of formula (I) or pharmaceutically acceptable salts or *in vivo* hydrolysable esters thereof in medicine, the definition of compounds includes compounds selected from:

i) a compound of the formula (I) wherein R³ and R⁴ are both methyl, R⁵ is hydrogen, fluoro or chloro, Y-Z is ethynylene, X is phenyl and one of R¹ and R² is hydrogen and the other is

pyrimidyl-NH-C(O)- or triazinyl-NH-C(O)- (wherein said triazine or pyrimidine is substituted by methyl, methoxy or dimethylamino) and the -SO₂NR¹R² moiety is ortho to Y-Z;

ii) 4-(3-hydroxy-3-methyl-1-butynyl)-N-(3-methyl-2-pyridinyl)-benzenesulphonamide;

iii) N-{4-[N,N-bis-(*sec*-butyl)aminosulphonyl]phenyl}-2-hydroxy-2-methyl-3,3,3-

5 trifluoropropanamide; or

iv) N-{4-[N,N-bis-(*iso*-butyl)aminosulphonyl]phenyl}-2-hydroxy-2-methyl-3,3,3-

trifluoropropanamide;

and their pharmaceutically acceptable salts and *in vivo* hydrolysable esters.

As stated above the size of the dose required for the therapeutic or prophylactic treatment of
 10 a particular disease state will necessarily be varied depending on the host treated, the route of administration and the severity of the illness being treated. Preferably a daily dose in the range of 1-50 mg/kg is employed. However the daily dose will necessarily be varied depending upon the host treated, the particular route of administration, and the severity of the illness being treated. Accordingly the optimum dosage may be determined by the practitioner who is treating any
 15 particular patient.

The elevation of PDH activity described herein may be applied as a sole therapy or may involve, in addition to the subject of the present invention, one or more other substances and/or treatments. Such conjoint treatment may be achieved by way of the simultaneous, sequential or separate administration of the individual components of the treatment. For example in the
 20 treatment of diabetes mellitus chemotherapy may include the following main categories of treatment:

i) insulin;

ii) insulin secretagogue agents designed to stimulate insulin secretion (for example glibenclamide, tolbutamide, other sulphonylureas);

25 iii) oral hypoglycaemic agents such as metformin, thiazolidinediones;

iv) agents designed to reduce the absorption of glucose from the intestine (for example acarbose);

v) agents designed to treat complications of prolonged hyperglycaemia;

vi) other agents used to treat lactic acidemia;

vii) inhibitors of fatty acid oxidation;

viii) lipid lowering agents;

ix) agents used to treat coronary heart disease and peripheral vascular disease such as aspirin, pentoxifylline, cilostazol; and/or

x) thiamine.

- 5 As stated above the compounds defined in the present invention are of interest for their ability to elevate the activity of PDH. Such compounds of the invention may therefore be useful in a range of disease states including diabetes mellitus, peripheral vascular disease, (including intermittent claudication), cardiac failure and certain cardiac myopathies, myocardial ischaemia, cerebral ischaemia and reperfusion, muscle weakness, hyperlipidaemias, Alzheimers disease and/or
- 10 atherosclerosis.

In addition to their use in therapeutic medicine, the compounds of formula (I) and their pharmaceutically acceptable salts are also useful as pharmacological tools in the development and standardisation of *in vitro* and *in vivo* test systems for the evaluation of the effects of elevators of PDH activity in laboratory animals such as cats, dogs, rabbits, monkeys, rats and mice, as part of

15 the search for new therapeutic agents.

The invention will now be illustrated by the following non-limiting examples in which, unless stated otherwise:

- (i) temperatures are given in degrees Celsius (°C); operations were carried out at room or ambient temperature, that is, at a temperature in the range of 18-25°C;
- 20 (ii) organic solutions were dried over anhydrous magnesium sulphate; evaporation of solvent was carried out using a rotary evaporator under reduced pressure (600-4000 Pascals; 4.5-30 mm Hg) with a bath temperature of up to 60°C;
- (iii) chromatography unless otherwise stated means flash chromatography on silica gel; thin layer chromatography (TLC) was carried out on silica gel plates; where a "Bond Elut" column is
- 25 referred to, this means a column containing 10g or 20g of silica of 40 micron particle size, the silica being contained in a 60ml disposable syringe and supported by a porous disc, obtained from Varian, Harbor City, California, USA under the name "Mega Bond Elut SI"
- (iv) in general, the course of reactions was followed by TLC and reaction times are given for illustration only;

(v) yields are given for illustration only and are not necessarily those which can be obtained by diligent process development; preparations were repeated if more material was required;

(vi) when given, ¹H NMR data is quoted and is in the form of delta values for major diagnostic protons, given in parts per million (ppm) relative to tetramethylsilane (TMS) as an internal

5 standard, determined at 300 MHz using perdeuterio dimethyl sulphoxide (DMSO-d₆) as the solvent unless otherwise stated; coupling constants (J) are given in Hz;

(vii) chemical symbols have their usual meanings; SI units and symbols are used;

(viii) solvent ratios are given in percentage by volume;

(ix) mass spectra (MS) were run with an electron energy of 70 electron volts in the chemical
10 ionisation (CI) mode using a direct exposure probe; where indicated ionisation was effected by electron impact (EI) or fast atom bombardment (FAB); where values for m/z are given, generally only ions which indicate the parent mass are reported, and unless otherwise stated the mass ion quoted is the negative mass ion - (M-H)⁻; and

(x) the following abbreviations are used:

15	DMSO	dimethyl sulphoxide;
	DMF	<i>N,N</i> -dimethylformamide;
	DCM	dichloromethane; and
	EtOAc	ethyl acetate.

20 Example 1

N-[2-Chloro-4-(morpholinosulphonyl)phenyl]-2-hydroxy-2-methylpropanamide

2-Acetoxyisobutyryl chloride (660 mg, 3.6 mmol) was added to a stirred mixture of 2-chloro-4-(morpholinosulphonyl)anilino (Method A) (1.0 g, 3.6 mmol) and pyridine (0.34 ml, 4.2 mmol) in DCM (10 ml). The resultant mixture was stirred at ambient temperature overnight,
25 then washed with 1M aqueous hydrochloric acid and brine, dried and evaporated to dryness. The residue was dissolved in 1% aqueous ethanol (10 ml) and lithium hydroxide monohydrate was added (300 mg, 7.5 mmol). The mixture was stirred for 2h at ambient temperature, then EtOAc (25 ml) was added and the resulting organic layer was washed with water, dried and evaporated to dryness to yield the title compound (226 mg, 0.6 mmol). NMR: 1.4 (s, 6H), 2.9 (m, 4H), 3.6
30 (m, 4H), 6.3 (s, 1H), 7.7 (d, 1H), 7.8 (m, 1H), 8.6 (d, 1H), 9.8 (s, 1H); MS: 363 (M+H)⁺.

Example 2**N-[2-Chloro-4-(piperidinosulphonyl)phenyl]-2-hydroxy-2-methylpropanamide**

A solution of *N*-[2-chloro-4-(fluorosulphonyl)phenyl]-2-acetoxy-2-methylpropanamide (Method B) (340 mg, 1 mmol), 4-dimethylaminopyridine (10 mg, 0.08 mmol) and piperidine (0.1 ml, 1 mmol) in acetonitrile (5 ml) was heated under reflux for 18h. After evaporation to dryness, the residue was dissolved in 1% aqueous ethanol (10 ml) and lithium hydroxide monohydrate (84 mg, 2 mmol) was added. The mixture was stirred for 2h at ambient temperature, then EtOAc (25 ml) was added. The resulting solution was washed with water, dried and evaporated to dryness. The residue was purified by column chromatography using 50% EtOAc/ isohexane to yield the title compound as an oil (226 mg, 0.6 mmol). NMR: 1.4 (s, 6H), 1.5 (m, 6H), 2.9 (m, 4H), 6.2 (d, 1H), 7.7 (d, 1H), 7.8 (m, 1H), 8.6 (d, 1H), 9.8 (s, 1H); MS: 361 (M+H)⁺.

Examples 3-21

The procedures described in Examples 1 and 2 were repeated using the appropriate amine to replace the morpholine or piperidine to obtain the compounds described below in yields of 30-65%. "Meth" refers to whether the Example was made by the procedure of Example 1 or 2.

Ex	Compound	MS	NMR	Meth
3	<i>N</i> -[2-Chloro-4-(methylamino sulphonyl)phenyl]-2-hydroxy-2-methylpropanamide	305	1.4 (s, 6H), 2.5 (d, 3H), 6.2 (s, 1H), 7.3 (m, 1H), 7.7 (m, 1H), 7.8 (d, 1H), 8.5 (d, 1H), 9.7 (s, 1H).	1
4	<i>N</i> -[2-Chloro-4-(dimethyl-aminosulphonyl)phenyl]-2-hydroxy-2-methylpropanamide	319	1.4 (s, 6H), 2.6 (m, 6H), 6.2 (s, 1H), 7.7 (m, 1H), 7.8 (d, 1H), 8.6 (d, 1H), 9.8 (s, 1H).	1
5	<i>N</i> -[2-Chloro-4-(allylamino-sulphonyl)phenyl]-2-hydroxy-2-methylpropanamide	331	1.4 (s, 6H), 3.2 (m, 2H), 5.1 (m, 2H), 5.6 (m, 1H), 6.2 (br, 1H), 7.7 (m, 1H), 7.8 (m, 1H), 7.85 (1H, m), 8.5 (d, 1H), 9.7 (s, 1H).	2

6	<i>N</i> -[2-Chloro-4-(isopropyl-aminosulphonyl)phenyl]-2-hydroxy-2-methylpropanamide	333	0.9 (d, 6H), 1.4 (s, 6H), 2.7 (m, 1H), 6.2 (s, 1H), 7.6 (m, 1H), 7.8 (m, 1H), 7.9 (d, 1H), 8.6 (d, 1H), 9.8 (s, 1H).	2
7	<i>N</i> -[2-Chloro-4-(<i>n</i> -butylamino-sulphonyl)phenyl]-2-hydroxy-2-methylpropanamide	347	0.8 (m, 3H), 1.2 (m, 4H), 1.4 (s, 6H), 2.7 (m, 2H), 6.2 (s, 1H), 7.6 (m, 1H), 7.8 (m, 1H), 7.9 (d, 1H), 8.6 (d, 1H), 9.8 (s, 1H).	2
8	<i>N</i> -[2-Chloro-4-(<i>t</i> -butylamino-sulphonyl)phenyl]-2-hydroxy-2-methylpropanamide	347	1.0 (s, 9H), 1.4 (s, 6H), 6.2 (s, 1H), 7.6 (m, 1H), 7.8 (m, 1H), 7.9 (d, 1H), 8.6 (d, 1H), 9.8 (s, 1H).	2
9	<i>N</i> -[2-Chloro-4-(<i>n</i> -pentyl-aminosulphonyl)phenyl]-2-hydroxy-2-methylpropanamide	361	0.8 (m, 3H), 1.2 (m, 6H), 1.4 (s, 6H), 2.7 (m, 2H), 6.2 (s, 1H), 7.6 (m, 1H), 7.8 (m, 1H), 7.9 (d, 1H), 8.5 (d, 1H), 9.8 (s, 1H).	2
10	<i>N</i> -[2-Chloro-4-(cyclohexyl-aminosulphonyl)phenyl]-2-hydroxy-2-methylpropanamide	373	1.0-1.2 (m, 10H), 1.4 (s, 6H), 2.7 (m, 1H), 6.2 (s, 1H), 7.6 (m, 1H), 7.8 (m, 1H), 7.9 (d, 1H), 8.5 (d, 1H), 9.8 (s, 1H).	2
11	<i>N</i> -[2-Chloro-4-(benzylamino-sulphonyl)phenyl]-2-hydroxy-2-methylpropanamide	381	1.4 (s, 6H), 4.0 (s, 2H), 6.2 (br, 1H), 7.2 (m, 5H), 7.7 (m, 1H), 7.8 (m, 1H), 8.2 (br, 1H), 8.4 (d, 1H), 9.7 (s, 1H)	2
12	<i>N</i> -[2-Chloro-4-(pyrid-3-yl-methylaminosulphonyl)phenyl]-2-hydroxy-2-methylpropanamide	382	1.4 (s, 6H), 3.9 (m, 1H), 4.0 (d, 2H), 6.2 (s, 1H), 7.3 (br, 1H), 7.4 (m, 1H), 7.6 (d, 1H), 7.8 (m, 1H), 8.3 (m, 1H), 8.4 (m, 2H), 9.7 (m, 1H)	1
13	<i>N</i> -(2-Chloro-4-anilino-sulphonylphenyl)-2-hydroxy-2-methylpropanamide	367	1.4 (s, 6H), 6.2 (s, 1H), 7.0 (m, 3H), 7.2 (m, 3H), 7.7 (q, 1H), 7.8 (d, 1H), 8.4 (d, 1H), 9.7 (s, 1H).	1

14	<i>N</i> -[2-Chloro-4-(2-chloroanilino-sulphonyl)phenyl]-2-hydroxy-2-methylpropanamide	401	1.4 (s, 6H), 6.2 (s, 1H), 7.2 (m, 3H), 7.4 (m, 2H), 7.6 (q, 1H), 7.8 (d, 1H), 8.4 (d, 1H), 9.8 (s, 1H).	1
15	<i>N</i> -[2-Chloro-4-(3-iodoanilino-sulphonyl)phenyl]-2-hydroxy-2-methylpropanamide	493	1.4 (s, 6H), 6.2 (s, 1H), 7.1 (m, 2H), 7.4 (m, 3H), 7.7 (q, 1H), 7.8 (s, 1H), 8.4 (d, 1H), 9.8 (s, 1H).	1
16	<i>N</i> -[2-Chloro-4-(4-chloroanilino-sulphonyl)phenyl]-2-hydroxy-2-methylpropanamide	401	1.4 (s, 6H), 6.2 (s, 1H), 7.1 (d, 2H), 7.3 (d, 2H), 7.7 (m, 2H), 7.8 (d, 1H), 8.4 (d, 1H), 9.7 (s, 1H).	1
17	<i>N</i> -[2-Chloro-4-(4-methoxy-anilinosulphonyl)phenyl]-2-hydroxy-2-methylpropanamide	397	1.4 (s, 6H), 3.6 (s, 3H), 6.2 (br, 1H), 6.8 (d, 2H), 7.0 (d, 2H), 7.6 (m, 2H), 7.7 (d, 1H), 8.4 (d, 1H), 9.7 (s, 1H).	1
18	<i>N</i> -[2-Chloro-4-(2-chloro-4-iodoanilinosulphonyl)phenyl]-2-hydroxy-2-methylpropanamide	527	1.4 (s, 6H), 6.3 (s, 1H), 7.0 (d, 1H), 7.6 (m, 3H), 7.8 (d, 2H), 8.4 (d, 1H), 9.8 (s, 1H).	1
19	<i>N</i> -[2-Chloro-4-(2-fluoro-4-iodoanilinosulphonyl)phenyl]-2-hydroxy-2-methylpropanamide	511	1.4 (s, 6H), 6.3 (s, 1H), 7.0 (m, 1H), 7.5 (m, 1H), 7.6 (m, 2H), 7.8 (m, 2H), 8.5 (m, 1H), 9.8 (s, 1H).	1
20	<i>N</i> -[2-Chloro-4-(2,4-dibromo-anilinosulphonyl)phenyl]-2-hydroxy-2-methylpropanamide	523/ 525	1.4 (s, 6H), 6.2 (s, 1H), 7.6 (m, 3H), 7.8 (m, 2H), 8.2 (d, 2H), 9.6 (s, 1H).	1
21	<i>N</i> -[2-Chloro-4-(3-bromo-4-chloroanilinosulphonyl)phenyl]-2-hydroxy-2-methylpropanamide	479/ 481	1.4 (s, 6H), 6.2 (s, 1H), 7.1 (m, 1H), 7.5 (m, 3H), 7.8 (m, 2H), 8.3 (m, 1H), 9.8 (m, 1H)	1

Example 22**R/S-N-[2-Chloro-4-(3-bromo-4-chloroanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide**

A solution of R/S-3,3,3-trifluoro-2-hydroxy-2-methylpropanoylchloride (Method E1) (110 mg, 0.62 mmol) in DCM (5 ml) was added to a stirred mixture of 2-chloro-4-(3-bromo-4-chloroanilinosulphonyl)aniline (Method C) (270 mg, 0.68 mmol) in DCM (10 ml). The resultant mixture was stirred at ambient temperature overnight and then washed with aqueous sodium hydrogen carbonate solution and water, dried and evaporated to dryness. The residue was purified by column chromatography using 50% EtOAc/isohexane to yield the title compound as a foam (100 mg, 0.19 mmol). NMR: 1.9 (s, 3H), 6.2 (s, 1H), 7.1 (m, 1H), 7.5 (m, 3H), 7.8 (m, 2H), 8.4 (d, 1H), 9.6 (s, 1H); MS: 533/ 535.

Example 23**R-N-[2-Chloro-4-(4-methoxyanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide**

A solution of S-3,3,3-trifluoro-2-hydroxy-2-methylpropanoylchloride (Method P) (446 mg, 2.5 mmol) in DCM (15 ml) was added to a stirred mixture of 2-chloro-4-(4-methoxyanilino-sulphonyl)aniline (Method D) (650 mg, 2.1 mmol) and 2, 6-di-*t*-butylpyridine (0.56 ml, 2.5 mmol) in DCM (50 ml). The resultant mixture was stirred at ambient temperature overnight and then washed with 1M aqueous hydrochloric acid, aqueous sodium hydrogen carbonate solution and brine, dried and evaporated to dryness. The residue was purified by column chromatography using 5% EtOAc in DCM to yield the title compound as a foam (680 mg, 1.5 mmol). EA: found: C, 45.1; H, 3.8; N, 5.8%; $C_{17}H_{16}N_2F_3ClSO_3$, requires: C, 45.1; H, 3.5; N, 6.2%; NMR ($CDCl_3$): 1.9 (s, 3H), 3.75 (s, 3H), 3.8 (s, 1H), 6.4 (s, 1H), 6.8 (d, 2H), 7.0 (d, 2H), 7.6 (dd, 1H), 7.8 (d, 1H), 8.5 (d, 1H), 9.3 (s, 1H); MS: 451.

Examples 24-105

The procedure described in Example 23 was repeated using the appropriate 4-aminobenzenesulphonamide to replace the 2-chloro-4-[(4-methoxyanilino)sulphonyl]aniline to

obtain the compounds described below. "Meth" refers to the Method (see section on Starting Materials below) used to make said appropriate sulphonamide.

Ex	Compound	MS	NMR	Meth
24	R-N-[2-Chloro-4-(allylamino-sulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	385	(CDCl ₃): 1.8 (s, 3H), 3.6 (m, 2H), 3.7 (s, 1H), 4.6 (t, 1H), 5.2-5.5 (m, 2H), 5.7-5.8 (m, 1H), 7.8 (dd, 1H), 7.9 (d, 1H), 8.6 (d, 1H), 9.3 (s, 1H)	E
25	R-N-[2-Chloro-4-(thien-2-yl-methylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide *	441	1.6 (s, 3H), 4.2 (d, 2H), 6.9 (m, 1H), 7.1 (m, 1H), 7.4 (m, 1H), 7.7 (d, 1H), 7.9 (s, 1H), 8.0 (s, 1H), 8.2 (d, 1H), 8.4 (m, 1H), 9.8 (s, 1H)	F
26	R-N-[2-Chloro-4-(2-chloro-anilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	455	(CDCl ₃): 1.8 (s, 3H), 3.65 (s, 1H), 7.0 (s, 1H), 7.1 (m, 1H), 7.3 (m, 2H), 7.6 (d, 2H), 7.8 (d, 1H), 8.5 (d, 1H), 9.3 (s, 1H).	D
27	R-N-[2-Chloro-4-(methylamino-sulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	359	(CDCl ₃): 1.8 (s, 3H), 2.7 (d, 3H), 3.6 (s, 1H), 4.4 (m, 1H), 7.8 (dd, 1H), 7.9 (d, 1H), 8.6 (d, 1H), 9.3 (s, 1H).	E
28	R-N-[2-Chloro-4-(dimethylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	373	(CDCl ₃): 1.8 (s, 3H), 2.7 (s, 6H), 3.8 (s, 1H), 7.7 (dd, 1H), 7.9 (d, 1H), 8.6 (d, 1H), 9.3 (s, 1H).	E
29	R-N-[2-Chloro-4-(morpholino-sulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	415	(CDCl ₃): 1.8 (s, 3H), 3.0 (m, 4H), 3.6 (s, 1H), 3.8 (m, 4H), 7.7 (dd, 1H), 7.9 (d, 1H), 8.6 (d, 1H), 9.3 (s, 1H).	E
30	R-N-[2-Chloro-4-(diethylamino-sulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	401	(CDCl ₃): 1.2 (t, 6H), 1.8 (s, 3H), 3.2 (q, 4H), 3.8 (s, 1H), 7.7 (dd, 1H), 7.9 (d, 1H), 8.6 (d, 1H), 9.3 (s, 1H).	E

31	R-N-[2-Chloro-4-(4- <i>t</i> -butyl-anilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide *	477	1.1 (s, 9H), 1.6 (s, 3H), 7.0 (d, 2H), 7.3 (d, 2H), 7.7 (d, 1H), 7.9 (s, 1H), 8.2 (d, 1H), 9.8 (s, 1H).	F
32	R-N-[2-Chloro-4-(anilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide *	421	1.6 (s, 3H), 7.0 (m, 2H), 7.2 (m, 3H), 7.7 (m, 1H), 7.9 (m, 1H), 8.2 (d, 1H), 9.8 (s, 1H).	F
33	R-N-[2-Chloro-4-(4-fluoro-anilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide *	439	1.6 (s, 3H), 7.1 (m, 4H), 7.7 (m, 1H), 7.9 (s, 1H), 8.2 (d, 1H), 9.8 (d, 1H).	F
34	R-N-[2-Chloro-4-(4-acetamid-anilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide *	478	1.6 (s, 3H), 2.0 (s, 3H), 7.0 (m, 2H), 7.4 (m, 2H), 7.7 (m, 1H), 7.9 (s, 1H), 8.2 (d, 1H), 9.8 (d, 1H).	F
35	R-N-[2-Chloro-4-(4-mesy lanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methyl propanamide *	499	1.6 (s, 3H), 3.1 (s, 3H), 7.3 (d, 2H), 7.8 (d, 2H), 7.85 (dd, 1H), 7.95 (d, 1H), 8.2 (d, 1H).	F
36	R-N-[2-Chloro-4-(4-sulphamoyl-anilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide *	500	1.6 (s, 3H), 7.2 (s, 2H), 7.3 (d, 2H), 7.7 (d, 2H), 7.8 (dd, 1H), 7.95 (d, 1H), 8.2 (d, 1H).	F
37	R-N-[2-Chloro-4-(3-iodoanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide *	547	1.6 (s, 3H), 7.0 (m, 2H), 7.4 (m, 2H), 7.7 (m, 1H), 7.9 (s, 1H), 8.2 (d, 1H), 9.8 (s, 1H).	F
38	R-N-[2-Chloro-4-(4-methylthio-anilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	467	1.6 (s, 3H), 2.4 (s, 3H), 7.0 (d, 2H), 7.2 (d, 2H), 7.7 (dd, 1H), 7.8 (d, 1H), 8.2 (d, 1H).	D

39	R-N-[2-Chloro-4-(4-benzyloxy-anilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methyl-propanamide	527	1.6 (s, 3H), 5.0 (s, 2H), 6.85 (d, 2H), 7.0 (d, 2H), 7.4 (m, 5H), 7.6 (dd, 1H), 7.8 (d, 1H), 8.5 (d, 1H).	D
40	R-N-(2-Chloro-4-sulphamoyl-phenyl)-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	345	1.6 (s, 3H), 7.4 (s, 2H), 7.8 (dd, 1H), 7.9 (d, 1H), 8.2 (d, 1H).	D
41	R-N-{2-Chloro-4-[N,N-bis-(2-hydroxyethyl)aminosulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methyl-propanamide	433	1.6 (s, 3H), 3.2 (t, 4H), 3.5 (m, 4H), 4.8 (t, 2H), 7.8 (dd, 1H), 7.9 (d, 1H), 8.25 (d, 1H).	D
42	R-N-(2-Chloro-4-{4-[N,N-bis-(2-hydroxyethyl)aminosulphonyl]anilinosulphonyl}phenyl)-3,3,3-trifluoro-2-hydroxy-2-methyl-propanamide	588	1.6 (s, 3H), 3.1 (t, 4H), 3.5 (m, 4H), 4.8 (t, 2H), 7.25 (d, 2H), 7.7 (d, 2H), 7.8 (dd, 1H), 7.9 (d, 1H), 8.2 (d, 1H).	D
43	R-N-{2-Chloro-4-[4-(morpholino-sulphonyl)anilinosulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	570	1.6 (s, 3H), 2.95 (m, 4H), 3.75 (m, 4H), 7.2 (d, 2H), 7.7 (d, 2H), 7.8 (dd, 1H), 7.9 (d, 1H), 8.6 (d, 1H).	D #
44	R-N-[2-Chloro-4-(R,S-2,3-dihydroxy-propylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	419	1.6 (s, 3H), 2.6 (m, 1H), 2.9 (m, 1H), 3.2 (m, 2H), 3.5 (m, 1H), 4.5 (t, 1H), 4.75 (d, 1H), 7.8 (dd, 1H), 7.9 (d, 1H), 8.2 (d, 1H).	D
45	R-N-[2-Fluoro-4-(2-fluoro-anilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methyl-propanamide [†]	423	CDCl ₃ : 1.75 (s, 3H), 6.75 (s, 1H), 6.9-7.0 (m, 1H), 7.1-7.2 (m, 2H), 7.5-7.6 (m, 3H), 8.5 (t, 1H), 8.9 (s, 1H).	G

46	R-N-[2-Fluoro-4-(2-methoxy-anilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methyl-propanamide [†]	435	CDCl ₃ : 1.75 (s, 3H), 3.7 (s, 3H), 6.8 (d, 1H), 6.9 (t, 1H), 7.0 (s, 1H), 7.05 (t, 1H), 7.5-7.6 (m, 3H), 8.5 (t, 1H), 8.8 (s, 1H).	G
47	R-N-[2-Fluoro-4-(2-methylthio-anilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methyl-propanamide [†]	451	CDCl ₃ : 1.75 (s, 3H), 2.25 (s, 3H), 3.4 (s, 1H), 7.1 (t, 1H), 7.25 (t, 1H), 7.35 (d, 1H), 7.5-7.6 (m, 3H), 8.5 (t, 1H), 8.8 (s, 1H).	G
48	R-N-[2-Fluoro-4-(3-fluoroanilino-sulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methyl-propanamide [†]	423	CDCl ₃ : 1.75 (s, 3H), 6.7 (s, 1H), 6.8-6.95 (m, 3H), 7.15-7.25 (m, 1H), 7.5-7.6 (m, 2H), 8.5 (t, 1H), 8.8 (br s, 1H).	G
49	R-N-[2-Fluoro-4-(3-methoxy-anilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methyl-propanamide [†]	435	CDCl ₃ : 1.75 (s, 3H), 3.75 (s, 3H), 6.5 (s, 1H), 6.6 (d, 1H), 6.65-6.7 (m, 2H), 7.15 (t, 1H), 7.5-7.65 (m, 2H), 8.5 (t, 1H), 8.8 (s, 1H).	G
50	R-N-[2-Fluoro-4-(3-methylthio-anilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methyl-propanamide [†]	451	CDCl ₃ : 1.75 (s, 3H), 2.45 (s, 3H), 6.5 (s, 1H), 6.8 (d, 1H), 6.9-7.05 (m, 2H), 7.15 (t, 1H), 7.5-7.6 (m, 2H), 8.5 (t, 1H), 8.8 (s, 1H).	G
51	R-N-[2-Fluoro-4-(4-fluoroanilino-sulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methyl-propanamide [†]	423	1.6 (s, 3H), 7.0-7.1 (m, 4H), 7.5-7.6 (m, 2H), 7.7 (s, 1H), 7.9 (t, 1H), 9.8 (s, 1H), 10.3 (br s, 1H).	G
52	R-N-[2-Fluoro-4-(4-methoxy-anilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methyl-propanamide [†]	435	CDCl ₃ : 1.75 (s, 3H), 3.8 (s, 3H), 6.4 (s, 1H), 6.6 (d, 2H), 7.0 (d, 2H), 7.4-7.6 (m, 2H), 8.5 (t, 1H), 8.9 (br s, 1H).	G

53	R-N-[2-Fluoro-4-(4-benzyloxy-anilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide [†]	511	CDCl ₃ : 1.75 (s, 3H), 5.0 (s, 2H), 6.25 (s, 1H), 6.8-6.9 (m, 2H), 6.9-7.0 (m, 2H), 7.3-7.5 (m, 7H), 8.5 (t, 1H), 8.85 (br s, 1H).	G
54	R-N-{2-Fluoro-4-[1-(<i>t</i> -butyloxy-carbonyl)piperazin-4-ylsulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide [†]	498	1.3 (s, 9H), 1.6 (s, 3H), 2.8-2.9 (m, 4H), 3.3-3.4 (m, 4H), 7.6 (d, 1H), 7.65 (d, 1H), 7.8 (s, 1H), 8.1 (t, 1H), 9.85 (s, 1H).	G
55	R-N-[2-Fluoro-4-(2-sulphamoyl-anilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide [†]	484	1.6 (s, 3H), 7.25 (t, 1H), 7.45-7.55 (m, 2H), 7.7-7.9 (m, 5H), 8.0 (t, 1H), 9.35 (br s, 1H), 9.8 (s, 1H).	G
56	R-N-[2-Fluoro-(4-sulphamoyl-anilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide [†]	484	1.6 (s, 3H), 7.15-7.25 (m, 5H), 7.6-7.8 (m, 5H), 7.9 (t, 1H), 9.8 (s, 1H).	G
57	R-N-[2-Fluoro-4-(3-benzyloxy-anilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide [†]	511	CDCl ₃ : 1.8 (s, 3H), 5.0 (s, 2H), 6.6 (d, 1H), 6.65 (s, 1H), 6.7-6.8 (m, 2H), 7.1 (t, 1H), 7.3-7.4 (m, 5H), 7.5-7.6 (m, 2H), 8.5 (t, 1H), 8.9 (s, 1H).	G
58	R-N-(2-Fluoro-4-(4-methylthio-anilinosulphonyl)phenyl)-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide [†]	451	1.6 (s, 3H), 2.4 (s, 3H), 7.05 (d, 2H), 7.1 (d, 2H), 7.5-7.6 (m, 2H), 7.7 (s, 1H), 7.9 (t, 1H), 9.8 (s, 1H), 10.25 (br s, 1H).	G
59	R-N-{2-Methyl-4-[1-(<i>t</i> -butyloxy-carbonyl)piperazin-4-ylsulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide [†]	494	CDCl ₃ : 1.4 (s, 9H), 1.7 (s, 3H), 2.35 (s, 3H), 2.9-3.0 (m, 4H), 3.4-3.6 (m, 4H), 4.4 (s, 1H), 7.5 (s, 1H), 7.6 (d, 1H), 8.3 (d, 1H), 8.7 (br s, 1H).	J

60	R-N-[2-Methyl-4-(4-methylthio-anilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methyl-propanamide [†]	447	CDCl ₃ : 1.7 (s, 3H), 2.25 (s, 3H), 2.4 (s, 3H), 6.6 (s, 1H), 7.0 (d, 2H), 7.15 (d, 2H), 7.55-7.65 (m, 2H), 8.2 (d, 1H), 8.5 (s, 1H).	J
----	--	-----	--	---

Ex	Compound	MS	Meth
61	R-N-[2-Chloro-4-(2-chloro-4-iodoanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methyl-propanamide *	581	F
62	R-N-[2-Chloro-4-(4-carbamoylanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methyl-propanamide *	464	F
63	R-N-[2-Chloro-4-(3,5-ditrifluoromethylanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide *	557	F
64	R-N-[2-Chloro-4-(2-fluoro-4-iodoanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methyl-propanamide *	565	F
65	R-N-[2-Chloro-4-(1,3-benzodioxol-5-ylmethylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide *	479	F
66	R-N-[2-Chloro-4-(pyrid-2-ylmethylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide *	436	F
67	R-N-[2-Chloro-4-(pyrid-3-ylmethylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide *	436	F
68	R-N-[2-Chloro-4-(pyrid-4-ylmethylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide *	436	F
69	R-N-[2-Chloro-4-(3-nitro-4-trifluoromethylanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide *	534	F
70	R-N-[2-Chloro-4-(2,4-dichloroanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methyl-propanamide *	489	F
71	R-N-[2-Chloro-4-(2-nitroanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide *	466	F

72	R-N-[2-Chloro-4-(3-bromoanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methyl-propanamide *	501	F
73	R-N-[2-Chloro-4-(3-chloroanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methyl-propanamide *	455	F
74	R-N-[2-Chloro-4-(3,4-dichloroanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methyl-propanamide *	489	F
75	R-N-[2-Chloro-4-(3,5-dichloroanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methyl-propanamide *	489	F
76	R-N-[2-Chloro-4-(3-trifluoromethylanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide *	489	F
77	R-N-[2-Chloro-4-(3,4-dimethylanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methyl-propanamide *	449	F
78	R-N-[2-Chloro-4-(4-cyanoanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methyl-propanamide *	446	F
79	R-N-[2-Chloro-4-(4-chloroanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methyl-propanamide *	455	F
80	R-N-[2-Chloro-4-(4-nitroanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methyl-propanamide *	466	F
81	R-N-[2-Chloro-4-(4-acetylanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methyl-propanamide *	463	F
82	R-N-[2-Chloro-4-(4-methylanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methyl-propanamide *	435	F
83	R-N-[2-Chloro-4-(benzylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide *	435	F
84	R-N-[2-Chloro-4-(3,4-dichlorobenzylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methyl-propanamide *	503	F
85	R-N-[2-Chloro-4-(4-trifluoromethylanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methyl-propanamide *	489	F

86	R-N-[2-Chloro-4-(3,4-dimethoxyanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide *	481	F
87	R-N-[2-Chloro-4-(2,4-difluorobenzylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methyl-propanamide *	471	F
88	R-N-[2-Chloro-4-(3-methylanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methyl-propanamide *	435	F
89	R-N-[2-Chloro-4-(4-fluorobenzylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methyl-propanamide *	453	F
90	R-N-[2-Chloro-4-(4-trifluoromethoxybenzylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide *	519	F
91	R-N-[2-Chloro-4-(4-chlorobenzylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide *	470	F
92	R-N-[2-Chloro-4-[4-(1,2,3-thiadiazol-4-yl)benzylaminosulphonyl]phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide *	519	F
93	R-N-[2-Chloro-4-(2-fluoro-4-trifluoromethylbenzylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide *	521	F
94	R-N-[2-Chloro-4-(4-chloro-3-trifluoromethylanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide *	523	F
95	R-N-[2-Chloro-4-(2-chloro-5-methylanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide [†]	469	D
96	R-N-[2-Chloro-4-(N-methylanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methyl-propanamide [†]	435	D
97	R-N-[2-Chloro-4-(N-methyl-4-methoxyanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methyl-propanamide [†]	465	D
98	R-N-[2-Chloro-4-(indolin-1-ylsulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide [†]	447	D
99	R-N-[2-Chloro-4-(N-ethylanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methyl-propanamide	449	D

100	R-N-[2-Fluoro-4-(2-chloro-6-methylanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methyl-propanamide [†]	453	G
101	R-N-[2-Fluoro-4-(indolen-1-ylsulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide [†]	431	G
102	R-N-[2-Fluoro-4-(N-methylanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methyl-propanamide [†]	419	G
103	R-N-[2-Fluoro-4-(N-ethylanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methyl-propanamide [†]	433	G
104	R-N-[2-Fluoro-4-(N-methyl-4-methoxyanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methyl-propanamide [†]	449	G
105	R-N-[2-Methyl-4-(indolin-1-ylsulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide [†]	427	J

* indicates that no 2,6-di-*t*-butylpyridine was used in the reaction.

† indicates that no aqueous work-up was used during the final step; instead the reaction mixture was concentrated, DCM (5 ml) was added and the solution loaded onto a Bond Elut column. This was then eluted with *iso*-hexane (50 ml) followed by 10% EtOAc/ DCM to yield the product.

indicates that this sulphonamide was obtained as a by-product from preparation of Example 42

Example 106

10 R-N-[2-Fluoro-4-(3-hydroxyanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide

A solution of R-N-[2-fluoro-4-(3-benzyloxyanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide (Example 57) (0.36g, 0.7 mmol) in ethanol (15 ml) was hydrogenated over 10% Pd/ C for 4h at ambient temperature. The catalyst was removed by filtration and the ethanol evaporated to yield the title compound as a solid (0.18g, 0.4 mmol).
 15 NMR: 1.6 (s, 3H), 6.4 (d, 1H), 6.5 (d, 1H), 6.6 (s, 1H), 7.0 (t, 1H), 7.6 (d, 2H), 7.7 (s, 1H), 7.95 (t, 1H), 9.45 (s, 1H), 9.8 (s, 1H); MS: 421.

Example 107**R-N-[2-Fluoro-4-(4-hydroxyanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide**

5 The procedure of Example 106 was repeated using R-N-[2-fluoro-4-(4-benzyloxyanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide (Example 53) (0.39g, 0.8 mmol) as the starting material to yield the title compound, as a solid (0.26g, 0.6 mmol). NMR: 1.6 (s, 3H), 6.6 (d, 2H); 6.85 (d, 2H), 7.05-7.15 (m, 1H), 7.4-7.5 (m, 2H), 7.7 (s, 1H), 7.9 (t, 1H), 9.2 (s, 1H), 9.8 (s, 1H); MS: 421.

10

Example 108**R-N-[2-Fluoro-4-(3-mesylanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide**

15 A solution of R-N-[2-fluoro-4-(3-methylthioanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide (Example 50) (360 mg, 0.8 mmol) and 70% 3-chloroperoxybenzoic acid (390 mg, 1.6 mmol) in DCM (20 ml) was stirred at ambient temperature overnight and then evaporated to dryness and the residue treated with aqueous sodium hydrogen carbonate solution (25 ml). The aqueous solution was extracted with EtOAc, the EtOAc extracts were washed with brine, dried and evaporated to yield the title compound as a solid (0.1g, 0.2 mmol).
20 NMR: 1.6 (s, 3H), 3.05 (s, 3H), 7.2-7.7 (m, 8H), 7.95 (t, 1H), 9.8 (s, 1H); MS: 483.

Example 109**R-N-[2-Fluoro-4-(2-mesylanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide**

25 The procedure of Example 108 was repeated using R-N-[2-fluoro-4-(2-methylthioanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide (Example 47) (290 mg, 0.6 mmol) as the starting material to yield the title compound as a solid (100 mg, 0.2 mmol). NMR: 1.6 (s, 3H), 3.25 (s, 3H), 7.25 (d, 1H), 7.4 (t, 1H), 7.5 (t, 1H), 7.6-7.8 (m, 3H), 7.85-7.95 (m, 2H), 8.05 (t, 1H), 9.85 (s, 1H); MS: 483.

Example 110**R-N-{2-Chloro-4-[4-(2-hydroxyethylthio)anilinosulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide**

5 A solution of S-3,3,3-trifluoro-2-hydroxy-2-methylpropanoyl chloride (Method P) (700 mg, 3.92 mmol) in DCM (25 ml) was added to a stirred mixture of 2-chloro-4-[4-(2-hydroxyethylthio)anilinosulphonyl]aniline (Method R) (630 mg, 1.76 mmol) and 2,6-di-*t*-butylpyridine (0.9 ml, 4.0 mmol) in DCM (50 ml). The resultant mixture was stirred at ambient temperature overnight and then washed with 1M aqueous hydrochloric acid, aqueous sodium hydrogen carbonate solution and brine then dried and evaporated to dryness. The residue was purified by column chromatography using 20% EtOAc in DCM to yield the title compound as a foam (330 mg, 0.66 mmol). NMR: 1.6 (s, 3H), 2.9 (t, 2H), 3.5 (m, 2H), 4.8 (t, 1H), 7.0 (d, 2H), 7.2 (d, 2H), 7.7 (dd, 1H), 7.8 (d, 1H), 8.2 (d, 1H); MS: 497.

15 Example 111**R-N-{2-Chloro-4-[4-(2-hydroxyethylsulphonyl)anilinosulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide**

A solution of R-N-{2-chloro-4-[4-(2-hydroxyethylthio)anilinosulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide (Example 110) (270 mg, 0.54 mmol) and 55% 3-chloroperoxybenzoic acid (340 mg, 1.08 mmol) in DCM (25 ml) was stirred at ambient temperature overnight and then evaporated to dryness and the residue treated with aqueous sodium hydrogen carbonate solution (25 ml). The aqueous solution was extracted with EtOAc, the EtOAc extracts were washed with brine, dried and evaporated. The residue was purified by column chromatography using 50% EtOAc in DCM to yield the title compound as a foam (144 mg, 0.27 mmol). NMR: 1.6 (s, 3H), 3.35 (t, 2H), 3.6 (m, 2H), 4.8 (t, 1H), 7.3 (d, 2H), 7.75 (d, 2H), 7.85 (dd, 1H), 7.95 (d, 1H), 8.25 (d, 1H); MS: 529.

Example 112**R-N-{2-Chloro-4-[4-(2-ethoxyethylthio)anilinosulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide**

A solution of S-3,3,3-trifluoro-2-hydroxy-2-methylpropanoyl chloride (Method P) (393 mg; 2.2 mmol) in DCM (25 ml) was added to a stirred mixture of 2-chloro-4-[4-(2-ethoxyethylthio)anilinosulphonyl]aniline (Method S) (770 mg, 2.0 mmol) and 2,6-di-*t*-butylpyridine (0.51 ml, 2.2 mmol) in DCM (25 ml). The resultant mixture was stirred at ambient temperature for 6h and then washed with 1M aqueous hydrochloric acid and brine, dried and evaporated to dryness. The residue was purified by column chromatography using 10% EtOAc in DCM to yield the title compound as a foam (520 mg, 0.99 mmol). NMR: 1.0 (t, 3H), 1.6 (s, 3H), 3.0 (t, 2H), 3.3-3.5 (m, 4H), 7.0 (d, 2H), 7.2 (d, 2H), 7.7 (dd, 1H), 7.8 (d, 1H), 8.2 (d, 1H); MS: 525.

Example 113**R-N-{2-Chloro-4-[4-(2-ethoxyethylsulphonyl)anilinosulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide**

The procedure of Example 111 was repeated using R-N-{2-chloro-4-[4-(2-ethoxyethylthio)anilinosulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide (Example 112) (460 mg, 0.87 mmol) as the starting material to obtain the title compound as a foam (260 mg, 0.46 mmol). NMR: 0.7 (t, 3H), 1.6 (s, 3H), 3.1 (t, 2H), 3.45 (t, 2H), 3.55 (t, 2H), 7.3 (d, 2H), 7.75 (d, 2H), 7.85 (dd, 1H), 8.0 (d, 1H), 8.3 (d, 1H); 557.

Example 114**R-N-[2-Chloro-4-(4-methoxycarbonyl)anilinosulphonyl]phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide**

A solution of S-3,3,3-trifluoro-2-hydroxy-2-methylpropanoyl chloride (Method P) (8.24 g, 46.2 mmol) in DCM (200 ml) was added to a stirred mixture of 2-chloro-4-(4-methoxycarbonyl)anilinosulphonyl)aniline (Method T) (13.6 g, 40.0 mmol) and 2,6-di-*t*-butylpyridine (10.3 ml, 46.2 mmol) in DCM (200 ml). The resultant mixture was stirred at ambient temperature overnight and then washed with 1M aqueous hydrochloric acid, aqueous sodium hydrogen carbonate solution and brine, dried and evaporated to dryness. The residue was purified by

column chromatography using 10% EtOAc in DCM to yield the title compound as a solid (16.6 g, 34.5 mmol). NMR: 1.6 (s, 3H), 3.8 (s, 3H), 7.2 (d, 2H), 7.8 (dd, 1H), 7.85 (d, 2H), 7.9 (d, 1H), 8.2 (d, 1H); MS: 479.

5 **Example 115**

R-N-[2-Chloro-4-(4-carboxyanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide

A solution of lithium hydroxide (428 mg, 11.6 mmol) in water (10 ml) was added to a solution of **R-N-[2-chloro-4-(4-methoxycarbonylanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide** (Example 114) (610 mg, 1.27 mmol) in methanol (20 ml). The resultant mixture was stirred at ambient temperature overnight and evaporated to dryness. The residue was dissolved in 1M aqueous hydrochloric acid (15 ml), the aqueous solution was extracted with EtOAc, the EtOAc extracts were washed with brine, dried and evaporated to yield the title compound as a foam (560 mg, 1.2 mmol); microanalysis found: C, 43.8; H, 3.0; N, 5.7%; $C_{17}H_{14}N_2F_3ClSO_6$ requires: C, 43.7; H, 3.0; N, 6.0%; NMR: 1.6 (s, 3H), 7.2 (d, 2H), 7.8 (d, 2H), 7.8 (dd, 1H), 7.95 (d, 1H), 8.2 (d, 1H); MS: 465.

Example 116

R-N-[2-Chloro-4-(piperazin-1-ylsulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide

A solution of **S-3,3,3-trifluoro-2-hydroxy-2-methylpropanoyl chloride** (Method P) (5.6 g, 31.2 mmol) in DCM (250 ml) was added to a stirred mixture of 2-chloro-4-[1-(*t*-butoxycarbonyl)piperazin-4-ylsulphonyl]aniline (Method U) (10.7 g, 28.4 mmol) and 2,6-di-*t*-butylpyridine (7.1 ml, 31.2 mmol) in EtOAc (750 ml). The resultant mixture was stirred at ambient temperature overnight, the DCM evaporated and the EtOAc layer washed with aqueous sodium hydrogen carbonate solution and brine, dried and evaporated to dryness. The residue was dissolved in 3M hydrogen chloride in EtOAc (70 ml), stirred at ambient temperature overnight and evaporated to dryness. The residue was treated with saturated sodium hydrogen carbonate solution, the aqueous solution was extracted with EtOAc, the EtOAc extracts were washed with brine, dried and evaporated. The residue was crystallised from EtOAc to yield the title compound

as a solid (7.5 g, 18.0 mmol). NMR: 1.6 (s, 3H), 2.7 (m, 4H), 2.8 (m, 4H), 7.7 (dd, 1H), 7.85 (d, 1H), 8.3 (d, 1H); MS: 414.

Example 117

5 R-N-[2-Chloro-4-(1-acetylpiperazin-4-ylsulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide

A solution of R-N-[2-chloro-4-(piperazin-1-ylsulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide (Example 116) (330 mg, 0.8 mmol) and triethylamine (0.33 ml) in DCM (20 ml) was added to a stirred solution of acetyl chloride (0.13 ml, 1.8 mmol) in DCM
 10 (15 ml). The resultant mixture was stirred at ambient temperature overnight and evaporated to dryness. The residue was dissolved in methanol (10 ml) and added to a stirred solution of lithium hydroxide (150 mg, 3.75 mmol) in water (5 ml). The resultant mixture was stirred at ambient temperature for 4h and then evaporated to dryness and the residue treated with 1M aqueous
 hydrochloric acid (15 ml). The aqueous solution was extracted with EtOAc, the EtOAc extracts
 15 were washed with brine, dried and evaporated to yield the title compound as a foam (81 mg, 0.18 mmol). NMR: 1.6 (s, 3H), 2.0 (s, 3H), 3.0 (m, 4H), 3.6 (m, 4H), 7.8 (dd, 1H), 7.9 (d, 1H), 8.4 (d, 1H); MS: 456.

Examples 118-125

20 The procedure of Example 117 was repeated using the appropriate acid chloride or sulphonyl chloride to replace the acetyl chloride to obtain the compounds described below.

Ex	Compound	MS	¹ H NMR
118	R-N-[2-Chloro-4-(1-mesylpiperazin-4-ylsulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	492	1.6 (s, 3H), 2.9 (s, 3H), 3.0 (m, 4H), 3.2 (m, 4H), 7.8 (dd, 1H), 7.9 (d, 1H), 8.4 (d, 1H).
119	R-N-{2-Chloro-4-[1-(4-mesylphenylsulphonyl)piperazin-4-ylsulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	632	1.6 (s, 3H), 3.0 (m, 8H), 3.3 (s, 3H), 7.7 (dd, 1H), 7.8 (d, 1H), 7.95 (d, 2H), 8.15 (d, 2H), 8.4 (d, 1H).

120	R-N-{2-Chloro-4-[1-(methoxyacetyl)-piperazin-4-ylsulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	486	1.6 (s, 3H), 2.95 (m, 4H), 3.2 (s, 3H), 3.5 (m, 4H), 4.0 (s, 2H), 7.8 (dd, 1H), 7.9 (d, 1H), 8.4 (d, 1H).
121	R-N-{2-Chloro-4-[1-(methoxypropionyl)-piperazin-4-ylsulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	500	1.6 (s, 3H), 2.5 (t, 2H), 2.95 (m, 4H), 3.2 (s, 3H), 3.5 (m, 4H), 3.6 (t, 2H), 7.8 (dd, 1H), 7.9 (d, 1H), 8.4 (d, 1H).
122	R-N-{2-Chloro-4-[1-(acetamidoacetyl)-piperazin-4-ylsulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	513	1.6 (s, 3H), 1.9 (s, 3H), 3.0 (m, 4H), 3.6 (m, 4H), 3.9 (d, 2H), 7.8 (dd, 1H), 7.9 (d, 1H), 8.4 (d, 1H).
123	R-N-{2-Chloro-4-[1-(hydroxyacetyl)-piperazin-4-ylsulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	472	1.6 (s, 3H), 2.8 (m, 2H), 3.0 (m, 4H), 3.45-3.6 (m, 4H), 7.8 (dd, 1H), 7.9 (d, 1H), 8.4 (d, 1H).
124	R-N-{2-Chloro-4-[1-(R,S-tetrahydrofuran-3-ylcarbonyl)piperazin-4-ylsulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	512	1.6 (s, 3H), 1.9 (m, 2H), 3.0 (m, 4H), 3.2 (m, 1H), 3.5-3.6 (m, 8H), 7.8 (dd, 1H), 7.9 (d, 1H), 8.4 (d, 1H).
125	R-N-{2-Chloro-4-[1-(morpholino-carbonyl)piperazin-4-ylsulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	527	1.6 (s, 3H), 2.9 (m, 4H), 3.1 (m, 4H), 3.2 (m, 4H), 3.5 (m, 4H), 7.8 (dd, 1H), 7.9 (d, 1H), 8.4 (d, 1H).

Example 126**R-N-{2-Chloro-4-[(1-aminoacetyl)piperazin-4-yl]sulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide**

- 5 A solution of R-N-[2-chloro-4-(piperazin-1-ylsulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide (Example 116) (330 mg, 0.8 mmol) in tetrahydrofuran (3 ml) was added to a stirred solution of (*t*-butoxycarbonyl)glycine (149 mg, 0.85 mmol) and carbonyldiimidazole (139 mg, 0.85 mmol) in tetrahydrofuran (8 ml). The resultant mixture was stirred at ambient temperature overnight and evaporated to dryness. The residue was dissolved in

0.7M aqueous citric acid (30 ml); the aqueous solution was extracted with EtOAc, the EtOAc extracts were washed with saturated aqueous sodium hydrogen carbonate solution and brine, dried and evaporated to dryness. The residue was dissolved in 3M hydrogen chloride in EtOAc (10 ml), stirred at ambient temperature for 3h and evaporated to dryness. The residue was treated with saturated sodium hydrogen carbonate solution; the aqueous solution was extracted with EtOAc, the EtOAc extracts were washed with brine, dried and evaporated to yield the title compound as a foam (230 mg, 0.49 mmol). NMR: 1.6 (s, 3H), 3.0 (m, 4H), 3.4 (m, 2H), 3.45-3.6 (m, 4H), 7.8 (dd, 1H), 7.9 (d, 1H), 8.4 (d, 1H); MS: 471.

10 Example 127

R-N-[2-Chloro-4-(N-acetylsulphamoyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide

A solution of R-N-(2-chloro-4-sulphamoylphenyl)-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide (Example 40) (150 mg, 0.43 mmol), acetic acid (62.4 mg, 1.04 mmol) and 4-dimethylaminopyridine (380 mg, 3.12 mmol) in DCM (25 ml) was stirred over 4A molecular sieve at ambient temperature for 3h. 1-(3-dimethylaminopropyl)-3-ethyl carbodiimide hydrochloride (300 mg, 1.57 mmol) was then added to this mixture. The resultant mixture was stirred at ambient temperature overnight and then filtered, washed with 1M aqueous hydrochloric acid, dried and evaporated to dryness. The residue was dissolved in methanol (10 ml) and added to a stirred solution of lithium hydroxide (92 mg, 2.3 mmol) in water (5 ml). The resultant mixture was stirred at ambient temperature for 4h and then evaporated to dryness and the residue treated with 1M aqueous hydrochloric acid (15 ml). The aqueous solution was extracted with EtOAc, the EtOAc extracts were washed with brine, dried and evaporated to yield the title compound as a foam (118 mg, 0.3 mmol); microanalysis found: C, 38.5; H, 3.6; N, 6.6%; $C_{12}H_{12}N_2F_3ClSO_5$, 0.5 EtOAc requires: C, 38.8; H, 3.7; N, 6.5%; NMR ($CDCl_3$): 1.8 (s, 3H), 2.1 (s, 3H), 7.9 (dd, 1H), 8.2 (d, 1H), 8.7 (s, 1H); MS: 387.

Examples 128-138

The procedure described in Example 127 was repeated using the appropriate acid to replace the acetic acid to obtain the compounds described below.

Ex	Example	MS
128	R-N-[2-Chloro-4-(<i>N</i> -benzoylsulphamoyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	449
129	R-N-{2-Chloro-4-[<i>N</i> -(2-chlorobenzoyl)sulphamoyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	483
130	R-N-{2-Chloro-4-[<i>N</i> -(3-iodobenzoyl)sulphamoyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	575
131	R-N-{2-Chloro-4-[<i>N</i> -(4-methylthiobenzoyl)sulphamoyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	495
132	R-N-{2-Chloro-4-[<i>N</i> -(4-cyanobenzoyl)sulphamoyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	474
133	R-N-{2-Chloro-4-[<i>N</i> -(4-benzyloxybenzoyl)sulphamoyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	555
134	R-N-{2-Chloro-4-[<i>N</i> -(4-methoxybenzoyl)sulphamoyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	479
135	R-N-{2-Chloro-4-[<i>N</i> -(4-trifluoromethylbenzoyl)sulphamoyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	517
136	R-N-{2-Chloro-4-[<i>N</i> -(2-chlorobenzylcarbonyl)sulphamoyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	497
137	R-N-{2-Chloro-4-[<i>N</i> -(4-dimethylaminosulphonylbenzoyl)sulphamoyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	556
138	R-N-{2-Chloro-4-[<i>N</i> -(4-mesylbenzoyl)sulphamoyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	527

Example 139

R-N-{2-Chloro-4-[*N*-(aminoacetyl)sulphamoyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide hydrochloride

5 A solution of R-N-(2-chloro-4-sulphamoylphenyl)-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide (Example 40) (345 mg, 1.0 mmol), (*tert*-butoxycarbonyl)glycine (420 mg,

2.4 mmol) and 4-dimethylaminopyridine (878 mg, 7.2 mmol) in DCM (35 ml) was stirred over 4A molecular sieve at ambient temperature for 3h. 1-(3-dimethylaminopropyl)-3-ethyl carbodiimide hydrochloride (688 mg, 3.6 mmol) was then added to this mixture. The resultant mixture was stirred at ambient temperature overnight and then filtered, washed with 0.7M aqueous citric acid and brine, dried and evaporated to dryness. The residue was dissolved in methanol (20 ml) and added to a stirred solution of lithium hydroxide (125 mg, 3.0 mmol) in water (10 ml). The resultant mixture was stirred at ambient temperature for 4h and then evaporated to dryness and the residue treated with 0.7M aqueous citric acid (15 ml). The aqueous solution was extracted with EtOAc, the EtOAc extracts were washed with brine, dried and evaporated. The residue was dissolved in 3M hydrogen chloride in EtOAc (5 ml), stirred at ambient temperature overnight and the solid filtered off to give the title compound (245 mg, 0.6 mmol); MS: 402.

Example 140

15 R-N-{2-Chloro-4-[N-(4-hydroxybenzoyl)sulphamoyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide

A solution of R-N-{2-chloro-4-[N-(4-benzyloxybenzoyl)sulphamoyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide (Example 133) (750 mg, 1.35 mmol) in EtOAc (40 ml) was hydrogenated over 10% Pd/ C for 4h at ambient temperature. The catalyst was removed by filtration and the EtOAc evaporated to yield the title compound as a foam (480 mg, 1.03 mmol). NMR: 1.6 (s, 3H), 6.7 (d, 2H), 7.7 (d, 2H), 7.8 (dd, 1H), 7.9 (d, 1H), 8.1 (d, 1H); MS: 465.

Example 141

25 R-N-[2-Chloro-4-(pyrrol-1-ylsulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide

A solution of R-N-(2-chloro-4-sulphamoylphenyl)-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide (Example 40) (345 mg, 1.0 mmol), 2,5-diethoxytetrahydrofuran (320 mg, 2.0 mmol) and 4-toluenesulphonic acid (190 mg, 1.0 mmol) in toluene (15 ml) was heated at 100°C overnight and evaporated to dryness. The residue was dissolved in EtOAc (50 ml) and washed with saturated sodium hydrogen carbonate solution and brine, dried and evaporated to dryness.

The residue was purified by column chromatography using 30% EtOAc in hexane to yield the title compound as a solid, (200 mg, 0.5 mmol). NMR: 1.6 (s, 3H), 6.35 (s, 2H), 7.35 (s, 2H), 7.9 (dd, 1H), 8.2 (d, 1H), 8.3 (d, 1H); MS: 395.

5 **Example 142**

R-N-[2-Chloro-4-[(2-morpholinoethylamino)sulphonyl]phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide

A solution of R-N-[2-chloro-4-(chlorosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide (Method Y) (366 mg, 1.0 mmol) in DCM (10 ml) was added to a stirred solution of 4-(2-aminoethyl)morpholine (286 mg, 2.2 mmol) in DCM (15 ml). The resultant mixture was stirred at ambient temperature overnight and then washed with aqueous sodium hydrogen carbonate solution and brine, dried and evaporated to dryness. The residue was purified by column chromatography using EtOAc to yield the title compound as a foam (384 mg, 0.84 mmol). NMR: 1.6 (s, 3H), 2.3 (m, 6H), 2.9 (m, 2H), 3.5 (m, 4H), 7.8 (dd, 1H), 7.9 (d, 1H), 8.2 (d, 1H); MS: 458.

Examples 143-208

The procedure described in Example 142 was repeated using the appropriate amine to replace the 4-(2-aminoethyl)morpholine using DCM or EtOAc as solvent to obtain the compounds described below.

Ex	Compound	MS	NMR
143	R-N-[2-Chloro-4-(4-hydroxypiperidino-sulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	429	1.4 (m, 2H), 1.6 (s, 3H), 1.75 (m, 2H), 2.8 (m, 2H), 3.2 (m, 2H), 3.5 (m, 1H), 4.6 (d, 1H), 7.8 (dd, 1H), 7.9 (d, 1H), 8.35 (d, 1H).
144	R-N-[2-Chloro-4-(2-hydroxyethylamino-sulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	389	1.6 (s, 3H), 2.8 (m, 2H), 3.4 (m, 2H), 4.7 (t, 1H), 7.7 (t, 1H), 7.8 (dd, 1H), 8.0 (d, 1H), 8.2 (d, 1H).

145	R-N-[2-Chloro-4-(2-R,S-hydroxypropyl-aminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide†1	403	1.0 (d, 3H), 1.6 (s, 3H), 2.65 (m, 2H), 3.6 (m, 1H), 4.65 (d, 1H), 7.7 (t, 1H), 7.8 (dd, 1H), 7.95 (d, 1H), 8.2 (d, 1H).
146	R-N-[2-Chloro-4-(2-R-hydroxypropylamino-sulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide†2	403	1.0 (d, 3H), 1.6 (s, 3H), 2.65 (m, 2H), 3.6 (m, 1H), 4.65 (d, 1H), 7.7 (t, 1H), 7.8 (dd, 1H), 7.95 (d, 1H), 8.2 (d, 1H).
147	R-N-[2-Chloro-4-(2-S-hydroxypropylamino-sulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide†3	403	1.0 (d, 3H), 1.6 (s, 3H), 2.65 (m, 2H), 3.6 (m, 1H), 4.65 (d, 1H), 7.7 (t, 1H), 7.8 (dd, 1H), 7.95 (d, 1H), 8.2 (d, 1H).
148	R-N-[2-Chloro-4-(2-piperidinoethyl-aminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	456	1.3 (m, 2H), 1.4 (m, 4H), 1.6 (s, 3H), 2.2 (m, 4H), 2.3 (t, 2H), 2.85 (t, 2H), 7.8 (dd, 1H), 7.9 (d, 1H), 8.3 (d, 1H).
149	R-N-[2-Chloro-4-(1-methylpiperazin-4-yl-sulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	428	1.6 (s, 3H), 2.15 (s, 3H), 2.45 (m, 4H), 2.95 (m, 4H), 7.75 (dd, 1H), 7.85 (d, 1H), 8.35 (d, 1H).
150	R-N-[2-Chloro-4-(trans/cis-4-hydroxy-cyclohexylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	443	1.1-1.5 (m, 8H), 1.6 (s, 3H), 3.0 (m, 1H), 4.4 (m, 1H), 7.75 (dd, 1H), 7.85 (d, 1H), 8.35 (d, 1H).

Ex	Compound	MS
151	R-N-[2-Chloro-4-(ethylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methyl-propanamide	373
152	R-N-[2-Chloro-4-(n-propylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	387

153	R-N-[2-Chloro-4-(<i>N</i> -methyl- <i>N</i> - <i>n</i> -propylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	401
154	R-N-[2-Chloro-4-(2-methoxyethylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	403
155	R-N-[2-Chloro-4-(3-methoxypropylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	417
156	R-N-[2-Chloro-4-(3-hydroxypropylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	403
157	R-N-{2-Chloro-4-[(ethoxycarbonylmethylaminocarbonylmethylamino)sulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methyl-propanamide	488
158	R-N-[2-Chloro-4-(methoxycarbonylethylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	431
159	R-N-[2-Chloro-4-(ethoxycarbonylpropylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	459
160	R-N-[2-Chloro-4-(2-acetamidoethylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	430
161	R-N-[2-Chloro-4-(3-morpholinopropylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	472
162	R-N-[2-Chloro-4-(<i>N</i> -methyl- <i>N</i> -allylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	399
163	R-N-[2-Chloro-4-(4-dimethylaminobutylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	444
164	R-N-[2-Chloro-4-(4-sulphamoylphenethylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	528
165	R-N-[2-Chloro-4-(2-methylthioethylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	419
166	R-N-[2-Chloro-4-(cyclopropylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	385

167	R-N-[2-Chloro-4-(cyclobutylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	399
168	R-N-{2-Chloro-4-[N,N-bis-(R,S-2-hydroxypropyl)aminosulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	461
169	R-N-{2-Chloro-4-[N-methyl-N-(2-hydroxyethyl)aminosulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	403
170	R-N-[2-Chloro-4-(thiazolidin-3-ylsulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	417
171	R-N-[2-Chloro-4-(pyrrolidin-1-ylsulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	399
172	R-N-{2-Chloro-4-[(R,S-1-hydroxymethyl-3-methylbutyl)aminosulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	445
173	R-N-[2-Chloro-4-(R,S-tetrahydrofuran-2-yl-methylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	429
174	R-N-[2-Chloro-4-(R,S-3-hydroxymethylpiperidinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	443
175	R-N-[2-Chloro-4-(thiomorpholinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	431
176	R-N-[2-Chloro-4-(R,S-2-hydroxymethylpiperidinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	443
177	R-N-{2-Chloro-4-[R,S-2-(2-hydroxyethyl)piperidinosulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	457
178	R-N-[2-Chloro-4-(pyrid-2-ylethylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	450
179	R-N-[2-Chloro-4-(R,S-1-phenyl-2-hydroxyethylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	465
180	R-N-[2-Chloro-4-(R,S-1-phenylethylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	449

181	R-N-[2-Chloro-4-(isopropylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	387
182	R-N-[2-Chloro-4-(R,S-1-methyl-2-hydroxyethylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	403
183	R-N-[2-Chloro-4-(R,S-1-ethyl-2-hydroxyethylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	417
184	R-N-[2-Chloro-4-(2-fluoroethylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	391
185	R-N-{2-Chloro-4-[2-(2-hydroxyethoxy)ethylaminosulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	433
186	R-N-[2-Chloro-4-(3-ethoxypropylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	431
187	R-N-[2-Chloro-4-(3-thiomethylpropylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	433
188	R-N-{2-Chloro-4-[(3-pyrrolin-1-yl)sulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	397
189	R-N-[2-Chloro-4-(R,S-1-hydroxymethyl-2-methylpropylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	431
190	R-N-[2-Chloro-4-(R,S-1-methyl-2-methoxyethylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	417
191	R-N-{2-Chloro-4-[N,N-bis-(2-methoxyethyl)aminosulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	461
192	R-N-[2-Chloro-4-(R,S-3-hydroxypyrrolidin-1-ylsulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	415
193	R-N-{2-Chloro-4-[(R,S-1-hydroxymethyl-3-methylthiopropyl)aminosulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	463
194	R-N-{2-Chloro-4-[4-(pyrrolidin-1-yl)piperidinosulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	482

195	R-N-[2-Chloro-4-(4-hydroxymethylpiperidinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	443
196	R-N-{2-Chloro-4-[N-methyl-N-(2-methoxyethyl)aminosulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	417
197	R-N-[2-Chloro-4-(1-hydroxyethylpiperazin-4-ylsulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	458
198	R-N-[2-Chloro-4-(3-imidazol-1-ylpropylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	453
199	R-N-{2-Chloro-4-[N-isopropyl-N-(2-hydroxyethyl)aminosulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	431
200	R-N-{2-Chloro-4-[N-(3-hydroxypropyl)-N-(2-hydroxyethyl)aminosulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	447
201	R-N-[2-Chloro-4-(N-methyl-N-R,S-tetrahydrofur-2-ylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	429
202	R-N-[2-Chloro-4-(R,S-1-ethylpiperidin-3-ylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	456
203	R-N-[2-Chloro-4-(3-dimethylaminopropylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	430
204	R-N-[2-Chloro-4-(5-hydroxypentylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	431
205	R-N-[2-Chloro-4-(4-hydroxybutylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	417
206	R-N-{2-Chloro-4-[N-methyl-N-(3-dimethylaminopropyl)aminosulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	444
207	R-N-{2-Chloro-4-[N-methyl-N-(2-dimethylaminoethyl)aminosulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	430
208	R-N-[2-Chloro-4-(1-methylhomopiperazin-4-ylsulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	442

†1The melting point for Example 145 was 117-118°C and the $[\alpha]_{\text{D}}^{20} +1.46^\circ$ (c, 10.2 in EtOH).

†2The melting point for Example 146 was 110-111°C and the $[\alpha]_{\text{D}}^{20} +0.85^\circ$ (c, 10.0 in EtOH).

†3The melting point for Example 147 was 134-135°C and the $[\alpha]_{\text{D}}^{20} +1.75^\circ$ (c, 10.7 in EtOH).

5 Example 209

10 R-N-[2-Chloro-4-(2-aminoethylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-
methylpropanamide hydrochloride.

A solution of R-N-[2-chloro-4-(chlorosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide (Method Y) (366 mg, 1.0 mmol) in DCM (15 ml) was added to a stirred solution of N-t-butoxycarbonyl ethylenediamine (350 mg, 2.2 mmol) in DCM (10 ml). The resultant mixture was stirred at ambient temperature for 4h, washed with 1M aqueous citric acid, saturated sodium hydrogen carbonate solution and brine, dried and evaporated to dryness. The residue was dissolved in 3M hydrogen chloride in EtOAc (5 ml) and stirred at ambient temperature overnight. The precipitated solid was filtered off, washed with EtOAc (5 ml) and dried to yield the title compound as a solid (362 mg, 0.63 mmol). NMR: 1.6 (s, 3H), 2.85 (m, 2H), 3.0 (m, 2H), 7.8 (dd, 1H), 8.0 (d, 1H), 8.3 (d, 1H); MS: 388.

Example 210

20 R-N-[2-Chloro-4-(2-dimethylaminoethylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-
methylpropanamide hydrochloride

A solution of R-N-[2-chloro-4-(chlorosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide (Method Y) (366 mg, 1.0 mmol) in DCM (10 ml) was added to a stirred solution of N,N-dimethylaminoethylamine (194 mg, 2.2 mmol) in DCM (15 ml). The resultant mixture was stirred at ambient temperature overnight and then washed with aqueous sodium hydrogen carbonate solution and brine, dried and evaporated to dryness. The residue was purified by conversion to the hydrochloride salt and recrystallization from EtOAc to yield the title compound (300mg, 0.72 mmol). NMR: 1.6 (s, 3H), 2.75 (s, 6H), 3.1 (s, 4H), 7.8 (dd, 1H), 8.0 (d, 1H), 8.3 (d, 1H); MS: 416.

Example 211**R-N-[2-Chloro-4-(4-aminoanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide**

A solution of R-N-[2-chloro-4-(chlorosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide (Method Y) (275 mg, 0.75 mmol) in DCM (15 ml) was added to a stirred solution of N-*t*-butoxycarbonyl-1,4-phenylenediamine (156 mg, 0.75 mmol) and pyridine (0.18 ml, 2.2 mmol) in DCM (10 ml). The resultant mixture was stirred at ambient temperature for 4h, washed with 1M aqueous citric acid, saturated sodium hydrogen carbonate solution and brine then dried and evaporated to dryness. The residue was dissolved in 3M hydrogen chloride in EtOAc (12 ml), stirred at ambient temperature overnight and evaporated to dryness. The residue was treated with saturated sodium hydrogen carbonate solution, the aqueous solution was extracted with EtOAc, the EtOAc extracts were washed with brine, dried and evaporated. The residue was purified by column chromatography using 25% EtOAc in DCM to yield the title compound as a foam (275 mg, 0.63 mmol). NMR: 1.6 (s, 3H), 5.0 (s, 2H), 6.4 (d, 2H), 6.7 (d, 2H), 7.6 (dd, 1H), 7.7 (d, 1H), 8.2 (d, 1H); MS: 436.

Example 212**R-N-[2-Chloro-4-(4-formylanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide**

A solution of R-N-[2-chloro-4-(chlorosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide (Method Y) (1.2 g, 3.28 mmol) in EtOAc (20 ml) was added to a stirred solution of 2-(4-aminophenyl)-1,3-dioxolane (Method V) (800 mg, 4.85 mmol) and pyridine (0.4 ml, 4.8 mmol) in EtOAc (80 ml). The resultant mixture was stirred at ambient temperature overnight, washed with 1M aqueous hydrochloric acid and brine, dried and evaporated to dryness. The residue was purified by column chromatography using 10% EtOAc in DCM to yield the title compound as a solid (1.1 g, 2.5 mmol). NMR: 1.6 (s, 3H), 7.3 (d, 2H), 7.8 (d, 2H), 7.85 (dd, 1H), 7.95 (d, 1H), 8.2 (d, 1H), 9.8 (s, 1H); MS: 449.

Example 213**R-N-[2-Chloro-4-(4-hydroxymethylanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide**

Sodium borohydride (38 mg, 1.0 mmol) was added to a solution of R-N-[2-chloro-4-(4-formylanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide (Example 212) (225 mg, 0.5 mmol) in ethanol (15 ml). The resultant mixture was stirred at ambient temperature overnight and evaporated to dryness. The residue was dissolved in 1M aqueous hydrochloric acid (15 ml), the aqueous solution was extracted with EtOAc, the EtOAc extracts were washed with brine, dried and evaporated to yield the title compound as a foam (220 mg, 0.5 mmol). NMR: 1.6 (s, 3H), 4.4 (s, 2H), 7.0 (d, 2H), 7.2 (d, 2H), 7.7 (dd, 1H), 7.85 (d, 1H) and 8.2 (d, 1H); MS: 451.

Example 214**R-N-{2-Chloro-4-[N-(2-hydroxyethyl)-N-(4-aminophenyl)aminosulphonyl]phenyl}3,3,3-trifluoro-2-hydroxy-2-methylpropanamide**

A solution of R-N-[2-chloro-4-(chlorosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide (Method Y) (580 mg, 1.58 mmol) in DCM (15 ml) was added to a stirred solution of 4-(2-hydroxyethylamino)aniline (Method W) (285 mg, 1.9 mmol) and pyridine (0.31 ml, 3.7 mmol) in DCM (15 ml). The resultant mixture was stirred at ambient temperature overnight, washed with 1M aqueous hydrochloric acid, dried and evaporated to dryness. The residue was dissolved in methanol (20 ml) and added to a stirred solution of lithium hydroxide (332 mg, 8.0 mmol) in water (10 ml). The resultant mixture was stirred at ambient temperature overnight and then evaporated to dryness and the residue treated with 1M aqueous hydrochloric acid to pH 7.0. The aqueous solution was extracted with EtOAc, the EtOAc extracts were washed with brine, dried and evaporated. The residue was purified by column chromatography using 50% EtOAc in DCM to yield the title compound as a foam (156 mg, 0.3 mmol). NMR: 1.6 (s, 3H), 3.35 (t, 2H), 3.5 (t, 2H), 4.7 (t, 1H), 5.2 (s, 2H), 6.4 (d, 2H), 6.7 (d, 2H), 7.6 (dd, 1H), 7.65 (d, 1H), 8.3 (d, 1H); MS: 480.

Example 215**R-N-{2-Chloro-4-[4-(2-hydroxyethylamino)anilinosulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide**

A solution of R-N-[2-chloro-4-(chlorosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide (Method Y) (770 mg, 2.1 mmol) in DCM (20 ml) was added to a stirred solution of 4-(N-2-tetrahydropyranyloxyethyl-t-butoxycarbonylamino)aniline (Method X) (620 mg, 1.85 mmol) and pyridine (0.34 ml, 4.08 mmol) in DCM (20 ml). The resultant mixture was stirred at ambient temperature overnight, washed with 1M aqueous citric acid, saturated sodium hydrogen carbonate solution and brine, dried and evaporated to dryness. The residue was dissolved in 3M hydrogen chloride in EtOAc (12 ml), stirred at ambient temperature for 4h and evaporated to dryness. The residue was treated with saturated sodium hydrogen carbonate solution, the aqueous solution was extracted with EtOAc, the EtOAc extracts were washed with brine, dried and evaporated. The residue was purified by column chromatography using 25% EtOAc in DCM to yield the title compound as a foam (260 mg, 0.45 mmol). NMR: 1.6 (s, 3H), 3.0 (m, 2H), 3.5 (m, 2H), 4.6 (t, 1H), 5.4 (t, 1H), 6.4 (d, 2H), 6.7 (d, 2H), 7.6 (dd, 1H), 7.7 (d, 1H), 8.2 (d, 1H); MS: 480.

Example 216**R-N-{2-Chloro-4-[4-(acetamididosulphonyl)anilinosulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide**

A solution of R-N-[2-chloro-4-(4-sulphamoylanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide (Example 36) (175 mg, 0.35 mmol), acetic acid (50 mg, 0.84 mmol) and 4-dimethylaminopyridine (306 mg, 2.51 mmol) in DCM (25 ml) was stirred over 4A molecular sieve at ambient temperature for 3h. To this mixture was added 1-(3-dimethylamino-propyl)-3-ethyl carbodiimide hydrochloride (242 mg, 1.26 mmol). The resultant mixture was stirred at ambient temperature overnight and then filtered, washed with 1M aqueous hydrochloric acid, dried and evaporated to dryness. The residue was dissolved in methanol (10 ml) and added to a stirred solution of lithium hydroxide (147 mg, 3.5 mmol) in water (5 ml). The resultant mixture was stirred at ambient temperature for 4h, evaporated to dryness and the residue treated with 1M aqueous hydrochloric acid (15 ml). The aqueous solution was extracted with EtOAc, the

EtOAc extracts were washed with brine, dried and evaporated to yield the title compound as a foam (140 mg; 0.26 mmol). NMR: 1.6 (s, 3H), 1.9 (s, 3H), 7.3 (d, 2H), 7.75 (d, 2H), 7.85 (dd, 1H), 7.95 (d, 1H), 8.25 (d, 1H); MS: 542.

15 **Example 217**

R-N-[2-Chloro-4-[4-(mesylaminocarbonyl)anilinosulphonyl]phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide

1-(3-Dimethylaminopropyl)-3-ethyl carbodiimide hydrochloride (195 mg, 1.02 mmol) was added to a solution of **R-N-[2-chloro-4-(4-carboxyanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide** (Example 115) (320 mg, 0.69 mmol), methane sulphonamide (79 mg, 0.82 mmol) and 4-dimethylaminopyridine (250 mg, 2.06 mmol) in DCM (25 ml). The resultant mixture was stirred at ambient temperature overnight, evaporated to dryness and the residue treated with 1M aqueous hydrochloric acid (15 ml). The aqueous solution was extracted with EtOAc, the EtOAc extracts were washed with brine, dried and evaporated to yield the title compound as a foam (130 mg, 0.24 mmol). NMR: 1.6 (s, 3H), 2.8 (s, 3H), 7.0 (d, 2H), 7.7 (d, 2H), 7.8 (dd, 1H), 7.9 (d, 1H), 8.2 (d, 1H); MS: 542.

Example 218

R-N-[2-Chloro-4-[4-(1-methylpiperazin-4-ylcarbonyl)anilinosulphonyl]phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide

Oxalyl chloride (0.07ml, 0.8 mmol) was added to a solution of **R-N-[2-chloro-4-(4-carboxyanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide** (Example 115) (314 mg, 0.67 mmol) in DCM (15 ml). The resultant mixture was stirred at ambient temperature overnight and evaporated to dryness. The residue was dissolved in EtOAc (5 ml) and added to a solution of 1-methylpiperazine (150 mg, 1.5 mmol) in EtOAc (5 ml). The resultant mixture was stirred at ambient temperature overnight, washed with 1M aqueous hydrochloric acid, saturated sodium hydrogen carbonate solution and brine, dried and evaporated to yield the title compound as a foam (355 mg, 0.65 mmol); MS: 547.

Examples 219-238:

The procedure of Example 218 was repeated using the appropriate amine to replace the 1-methylpiperazine to obtain the compounds described below.

Ex	Compound	MS
219	R-N-{2-Chloro-4-[4-(methylaminocarbonyl)anilinosulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	478
220	R-N-{2-Chloro-4-[4-(ethylaminocarbonyl)anilinosulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	492
221	R-N-{2-Chloro-4-[4-(n-propylaminocarbonyl)anilinosulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	506
222	R-N-{2-Chloro-4-[4-(isobutylaminocarbonyl)anilinosulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	520
223	R-N-{2-Chloro-4-[4-(allylaminocarbonyl)anilinosulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	504
224	R-N-{2-Chloro-4-[4-(dimethylaminocarbonyl)anilinosulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	492
225	R-N-{2-Chloro-4-[4-(2-methoxyethylaminocarbonyl)anilinosulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	522
226	R-N-{2-Chloro-4-[4-(2-dimethylaminoethylaminocarbonyl)anilinosulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	535
227	R-N-{2-Chloro-4-[4-(3-dimethylaminopropylaminocarbonyl)anilinosulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	549

Ex	Compound	MS	NMR
228	R-N-{2-Chloro-4-[4-(2-hydroxyethylaminocarbonyl)anilinosulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	508	1.6 (s, 3H), 3.4 (m, 2H), 3.5 (m, 2H), 7.2 (d, 2H), 7.8 (d, 2H), 7.85 (dd, 1H), 8.0 (d, 1H), 8.3 (d, 1H).

229	R-N-{2-Chloro-4-[4-(morpholinocarbonyl)anilinosulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	534	1.6 (s, 3H), 3.4 (m, 4H), 3.6 (m, 4H), 7.2 (d, 2H), 7.5 (d, 2H), 7.85 (dd, 1H), 7.95 (d, 1H), 8.3 (d, 1H).
230	R-N-{2-Chloro-4-[4-(R,S-2-hydroxy-propylaminocarbonyl)anilinosulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methyl-propanamide	522	1.1 (d, 3H), 1.6 (s, 3H), 3.2 (m, 2H), 3.8 (m, 1H), 7.2 (d, 2H), 7.8 (d, 2H), 7.85 (dd, 1H), 8.0 (d, 1H), 8.3 (d, 1H).
231	R-N-{2-Chloro-4-[4-(R,S-2-3-dihydroxy-propylaminocarbonyl)anilinosulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	538	1.6 (s, 3H), 3.4 (m, 4H), 3.6 (m, 1H), 7.2 (d, 2H), 7.8 (d, 2H), 7.85 (dd, 1H), 8.0 (d, 1H), 8.3 (d, 1H).
232	R-N-{2-Chloro-4-[4-(bis-2-hydroxyethyl-aminocarbonyl)anilinosulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methyl-propanamide	552	1.6 (s, 3H), 3.4 (m, 4H), 3.5 (m, 4H), 7.2 (d, 2H), 7.8 (d, 2H), 7.85 (dd, 1H), 8.0 (d, 1H), 8.3 (d, 1H).
233	R-N-{2-Chloro-4-[4-(trans/cis-4-hydroxycyclohexylaminocarbonyl)anilinosulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methyl-propanamide	562	1.6 (s, 3H), 1.3-1.9 (m, 8H), 3.8 (m, 2H), 7.2 (d, 2H), 7.8 (d, 2H), 7.85 (dd, 1H), 8.0 (d, 1H), 8.3 (d, 1H).
234	R-N-{2-Chloro-4-[4-(R,S-tetrahydrofur-2-ylmethylaminocarbonyl)anilinosulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	548	1.6 (s, 3H), 1.8 (m, 4H), 3.25 (t, 2H), 3.6 (m, 1H), 3.75 (m, 1H), 3.9 (m, 1H), 7.2 (d, 2H), 7.8 (d, 2H), 7.85 (dd, 1H), 8.0 (d, 1H), 8.3 (d, 1H).
235	R-N-{2-Chloro-4-[4-(R,S-1-methyl-2-hydroxyethylaminocarbonyl)anilinosulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	522	1.1 (d, 3H), 1.6 (s, 3H), 3.4 (m, 2H), 3.9 (m, 1H), 7.2 (d, 2H), 7.8 (d, 2H), 7.85 (dd, 1H), 8.0 (d, 1H), 8.3 (d, 1H).

236	<i>R-N</i> -{2-Chloro-4-[4-(<i>R,S</i> -1-methyl-2-methoxyethylaminocarbonyl)anilinosulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	536	1.1 (d, 3H), 1.6 (s, 3H), 3.2 (s, 3H), 3.4 (m, 2H), 4.1 (m, 1H), 7.2 (d, 2H), 7.8 (d, 2H), 7.85 (dd, 1H), 8.0 (d, 1H), 8.3 (d, 1H).
237	<i>R-N</i> -(2-Chloro-4-{4-[<i>N</i> -methyl- <i>N</i> -(2-hydroxyethyl)aminocarbonyl]anilinosulphonyl}phenyl)-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	522	1.6 (s, 3H), 2.9 (s, 3H), 3.4 (m, 2H), 3.5 (m, 2H), 7.2 (d, 2H), 7.8 (d, 2H), 7.85 (dd, 1H), 8.0 (d, 1H), 8.3 (d, 1H).
238	<i>R-N</i> -(2-Chloro-4-{4-[<i>N</i> -methyl- <i>N</i> -(2-methoxyethyl)aminocarbonyl]anilinosulphonyl}phenyl)-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	536	1.6 (s, 3H), 2.9 (s, 3H), 3.3 (s, 3H), 3.4 (m, 2H), 3.5 (m, 2H), 7.2 (d, 2H), 7.8 (d, 2H), 7.85 (dd, 1H), 8.0 (d, 1H), 8.3 (d, 1H).

Example 239

***R-N*-{2-Chloro-4-[4-(2-hydroxyethylaminosulphonyl)anilinosulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide**

A solution of *R-N*-[2-chloro-4-(4-fluorosulphonylanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide (Method O) (250 mg, 0.5 mmol), ethanolamine (92 mg, 1.5 mmol) and 4-dimethylaminopyridine (10 mg, 0.08 mmol) in ethanol (25 ml) was heated at 78°C overnight and evaporated to dryness. The residue was treated with 1M aqueous

hydrochloric acid (10 ml). The aqueous solution was extracted with EtOAc, the EtOAc extracts were washed with brine, dried and evaporated to yield the title compound as a foam (200 mg, 0.37 mmol). NMR: 1.6 (s, 3H), 2.8 (m, 2H), 3.3 (m, 2H), 4.6 (t, 1H), 7.3 (d, 2H), 7.7 (d, 2H), 7.8 (dd, 1H), 7.9 (d, 1H), 8.2 (d, 1H); MS: 544.

15 Examples 240-248

The procedure of Example 239 was repeated using the appropriate amine to replace the ethanolamine to obtain the compounds described below.

Ex	Compound	MS	NMR
240	<i>R-N</i> -{2-Chloro-4-[4-(<i>R,S</i> -2-3-dihydroxy-propylaminosulphonyl)anilinosulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	574	1.6 (s, 3H); 2.8 (m, 1H), 3.2 (m, 2H), 3.4 (m, 2H), 7.3 (d, 2H), 7.6 (d, 2H), 7.8 (dd, 1H), 7.9 (d, 1H), 8.2 (d, 1H).
241	<i>R-N</i> -{2-Chloro-4-[4-(<i>R,S</i> -2-hydroxypropylaminosulphonyl)anilinosulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	558	1.1 (d, 3H), 1.6 (s, 3H), 2.6 (m, 2H), 3.5 (m, 1H), 7.3 (d, 2H), 7.6 (d, 2H), 7.8 (dd, 1H), 7.9 (d, 1H), 8.2 (d, 1H).
242	<i>R-N</i> -{2-Chloro-4-[4-(2-dimethylaminoethylaminosulphonyl)anilinosulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide hydrochloride	571	1.6 (s, 3H), 2.7 (s, 6H), 3.1 (m, 4H), 7.3 (d, 2H), 7.6 (d, 2H), 7.8 (dd, 1H), 7.9 (d, 1H), 8.2 (d, 1H).
243	<i>R-N</i> -{2-Chloro-4-[4-(ethylaminosulphonyl)anilinosulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	528	0.9 (t, 3H), 1.6 (s, 3H), 2.7 (q, 2H), 7.3 (d, 2H), 7.6 (d, 2H), 7.8 (dd, 1H), 7.9 (d, 1H), 8.2 (d, 1H).
244	<i>R-N</i> -{2-Chloro-4-[4-(methylaminosulphonyl)anilinosulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	514	1.6 (s, 3H), 2.3 (d, 3H), 7.3 (d, 2H), 7.6 (d, 2H), 7.8 (dd, 1H), 7.9 (d, 1H), 8.2 (d, 1H).
245	<i>R-N</i> -{2-Chloro-4-[4-(allylaminosulphonyl)anilinosulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	540	1.6 (s, 3H), 3.4 (d, 2H), 5.0-5.2 (m, 2H), 5.65 (m, 1H), 7.3 (d, 2H), 7.6 (d, 2H), 7.8 (dd, 1H), 7.9 (d, 1H), 8.2 (d, 1H).
246	<i>R-N</i> -{2-Chloro-4-[4-(dimethylaminosulphonyl)anilinosulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	528	1.6 (s, 3H), 2.5 (s, 6H), 7.3 (d, 2H), 7.6 (d, 2H), 7.8 (dd, 1H), 7.9 (d, 1H), 8.2 (d, 1H).

247	R-N-{2-Chloro-4-[4-(R,S-3-hydroxy-pyrrolidin-1-ylsulphonyl)anilinosulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	570	1.6 (s, 3H), 1.65 (m, 2H), 2.9 (d, 1H), 3.2 (m, 3H), 4.1 (m, 1H), 7.3 (d, 2H), 7.6 (d, 2H), 7.8 (dd, 1H), 7.9 (d, 1H), 8.2 (d, 1H).
248	R-N-{2-Chloro-4-[4-(trans/cis-4-hydroxy-cyclohexylaminosulphonyl)anilinosulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	598	1.1-1.5 (m, 8H), 1.6 (s, 3H), 2.9 (d, 1H), 4.5 (m, 1H), 7.3 (d, 2H), 7.7 (d, 2H), 7.9 (dd, 1H), 8.1 (d, 1H), 8.3 (d, 1H).

Example 249

R-N-[2-Chloro-4-(5-trifluoromethylpyrid-2-ylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-

5 2-methylpropanamide

R-N-(2-chloro-4-chlorosulphonylphenyl)-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide (Method M) (293 mg, 0.80 mmol) and 2-amino-5-trifluoromethylpyridine (130 mg, 0.80 mmol) were dissolved in pyridine (2 ml). The resulting mixture was stirred at 85°C for 20h. The mixture was then cooled to room temperature and concentrated under vacuum to yield an oil, which was
 10 purified on a Bond Elut column to yield the title compound as a solid (285 mg, 0.58 mmol).
 NMR: 1.55 (s, 3H), 7.15 (d, 1H), 7.90 (m, 2H), 8.02 (m, 2H), 8.21 (d, 1H), 8.51 (s, 1H), 9.84 (s, 1H), 12.09 (br s, 1H); MS: 490.

Example 250-264

15 The procedure of Example 249 was repeated using the appropriate aminoheterocycle to replace the 2-amino-5-trifluoromethyl pyridine to obtain the compounds described below, in yields of 18-92%.

Ex	Example	MS	NMR
250	R-N-[2-Chloro-4-(pyrid-4-ylamino-sulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	422	1.59 (s, 3H), 6.91 (d, 2H), 7.76 (d, 1H), 7.84 (s, 1H), 7.90 (br s, 1H), 7.98 (d, 2H), 8.10 (d, 1H), 9.80 (s, 1H).

251	R-N-[2-Chloro-4-(pyrimidin-2-ylamino-sulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	423	1.53 (s, 3H), 7.00 (t, 1H), 7.89 (s, 1H), 7.93 (d, 1H), 8.00 (d, 1H), 8.18 (d, 1H), 8.47 (d, 2H), 9.80 (s, 1H), 12.0 (br s, 1H).
252	R-N-[2-Chloro-4-(5-methylpyrid-2-yl-aminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	436	1.47 (s, 3H), 2.02 (s, 3H), 7.02 (d, 1H), 7.51 (dd, 1H), 7.68-7.91 (m, 4H), 8.05 (d, 1H), 9.73 (br s, 1H).
253	R-N-[2-Chloro-4-(5-chloropyrid-2-yl-aminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	456	1.65 (s, 3H), 7.13 (d, 1H), 7.85 (d, 1H), 7.94 (d, 1H), 8.01 (s, 1H), 8.08 (s, 1H), 8.32 (m, 2H), 9.91 (s, 1H), 11.5 (br s, 1H).
254	R-N-[2-Chloro-4-(pyrid-2-ylamino-sulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	422	1.58 (s, 3H), 6.85 (t, 1H), 7.20 (d, 1H), 7.75 (t, 1H), 7.86 (d, 1H), 7.94 (m, 3H), 8.16 (d, 1H), 9.81 (s, 1H).
255	R-N-[2-Chloro-4-(2-carbamoylthien-2-yl-aminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide ¹¹	470	1.58 (s, 3H), 7.19 (d, 1H), 7.65-7.87 (m, 4H), 7.95 (s, 1H), 8.00 (s, 1H), 8.24 (d, 1H), 9.85 (s, 1H), 11.2 (br s, 1H).
256	R-N-[2-Chloro-4-(4,6-dimethylpyrid-2-yl-aminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methyl-propanamide	450	1.60 (s, 3H), 2.23 (s, 3H), 2.28 (s, 3H), 6.53 (s, 1H), 6.92 (br s, 1H) 7.75-7.96 (m, 3H), 8.13 (d, 1H), 9.82 (s, 1H).
257	R-N-[2-Chloro-4-(3,5-dichloropyrid-2-yl-aminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methyl-propanamide	490	
258	R-N-[2-Chloro-4-(pyrid-3-ylamino-sulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	422	1.58 (s, 3H), 7.30 (m, 1H), 7.50 (d, 1H), 7.73 (d, 1H), 7.88 (s, 1H), 7.95 (s, 1H), 8.27 (m, 3H), 9.83 (s, 1H).
259	R-N-[2-Chloro-4-(2-chloropyrid-3-yl-aminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	456	1.60 (s, 3H), 7.40 (m, 1H), 7.70 (d, 2H), 7.84 (s, 1H), 7.99 (s, 1H), 8.26 (m, 2H), 9.86 (s, 1H), 10.44 (s, 1H).

260	R-N-[2-Chloro-4-(4-dimethylamino-6-methylpyrimidin-2-ylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide ^{†2}	480	1.59 (s, 3H), 2.10 (s, 3H), 2.95 (s, 6H), 6.02 (s, 1H), 7.71 (d, 1H), 7.86 (s, 2H), 8.04 (d, 1H), 9.79 (s, 1H), 11.88 (br s, 1H).
261	R-N-[2-Chloro-4-(2,3-dimethylpyrazin-5-ylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide ^{†3}	451	1.58 (s, 3H), 2.32 (s, 6H), 7.94 (m, 2H), 8.05 (s, 1H), 8.10 (s, 1H), 8.22 (d, 1H), 9.88 (s, 1H), 11.31 (br s, 1H).
262	R-N-[2-Chloro-4-(3,5-dimethylisoxazol-4-ylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide ^{†4}	440	1.60 (s, 3H), 1.83 (s, 3H), 1.96 (s, 3H), 7.67 (dd, 1H), 7.78 (s, 1H), 8.23 (d, 1H).
263	R-N-[2-Chloro-4-(4,6-dimethylpyrimidin-2-ylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	451	1.59 (s, 3H), 2.25 (s, 6H), 6.72 (s, 1H), 7.91 (m, 2H), 8.12 (s, 1H), 8.17 (d, 1H), 9.83 (s, 1H).
264	R-N-[2-Chloro-4-(pyrazin-2-ylaminosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide	423	1.58 (s, 3H), 7.93 (m, 2H), 8.08 (s, 1H), 8.24 (m, 3H), 8.34 (s, 1H), 9.87 (s, 1H).

^{†1} The starting aminoheterocycle for Example 255, 2-amino-3-carbamoylthiophene, was prepared according to the procedure described in *J. Med. Chem.*, 1996, **39** (8), p1635.

^{†2} The starting aminoheterocycle for Example 260, 2-amino-4-dimethylamino-6-methylpyrimidine, was prepared according to the procedure described in US patent Application: US 77-5 769475 770217.

^{†3} The starting aminoheterocycle for Example 261, 2-amino-5,6-dimethylpyrazine, was prepared according to the procedure described in *Chem. Ber.*, 1967, **100** (2), p560-563.

^{†4} The starting aminoheterocycle for Example 262, 4-amino-3,5-dimethylisoxazole, was prepared according to the procedure described in *Helv. Chim. Acta*, 1991, **74** (3), p531.

Example 265

R-N-[2-Chloro-4-(N'-phenylureidosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide

A solution of R-N-(2-chloro-4-sulphamoylphenyl)-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide (Example 40) (345 mg, 1.0 mmol), phenyl isocyanate (122 mg, 1.02 mmol) and copper (I) chloride (5 mg, 0.052 mmol) in DMF (5 ml) was stirred at ambient temperature for 24h. The solution was poured onto water and this was acidified with 1M aqueous hydrochloric acid (1 ml). The precipitated solid was extracted with diethyl ether (25 ml) and the ether extract dried and evaporated. The residue was purified by column chromatography using 10% EtOAc in DCM to yield the title compound as a solid (70 mg, 0.15 mmol). NMR: 1.9 (s, 3H), 7.0 (t, 1H), 7.3 (t, 2H), 7.5 (m, 4H), 7.6 (d, 1H), 7.8 (dd, 1H), 7.9 (d, 1H); MS: 464.

Example 266

R-N-{2-Chloro-4-[N-methyl-N-(pyrazin-2-yl)aminosulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide

Potassium carbonate (65 mg, 0.47 mmol) and R-N-[2-chloro-4-(pyrazin-2-ylamino-sulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide (Example 264) (200 mg, 0.47 mmol) were added to acetone (10 ml). Methyl iodide (0.03 ml, 0.48 mmol) was added to the resulting suspension and the mixture was stirred at ambient temperature for 22h before being concentrated under vacuum. Water (15 ml) was added to the residue, which was acidified with 2M HCl and extracted with DCM (2x40 ml). The combined extracts were washed with brine (25 ml), dried and concentrated under vacuum to leave a foam, which was purified by column chromatography to yield the title compound as a foam (100 mg, 0.23 mmol). NMR: 1.60 (s, 3H), 3.25 (s, 3H), 7.66 (d, 1H), 7.82 (s, 1H), 8.27 (d, 1H), 8.42 (s, 1H), 8.52 (s, 1H), 8.80 (s, 1H); MS: 437.

Example 267**R-N-[2-Chloro-4-[N-methyl-N-(5-trifluoromethylpyrid-2-yl)aminosulphonyl]phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide**

Potassium carbonate (51 mg, 0.52 mmol) and R-N-[2-chloro-4-(5-trifluoromethylpyrid-2-yl)aminosulphonyl]phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide (Example 249) (180 mg; 0.37 mmol) were added to acetone (10 ml). Methyl iodide (0.023 ml, 0.37 mmol) was then added to the resulting suspension and the mixture was stirred at room temperature for 22h before being concentrated under vacuum. Water (2 ml) was added to the residue, which was acidified with 2M HCl and DCM (1 ml) was added. The mixture was absorbed onto a Bond Elut column and all organic components of the mixture were washed off with DCM. This resulting solution was concentrated under vacuum to yield a residue that was purified by column chromatography to yield the title compound as a foam (66 mg, 0.13 mmol). NMR: 1.60 (s, 3H), 3.38 (s, 3H), 7.74 (d, 1H), 7.78 (d, 1H), 7.93 (s, 1H), 8.05 (br s, 1H), 8.26 (d, 1H), 8.29 (d, 1H), 8.73 (s, 1H), 9.90 (br s, 1H); MS: 504.

Example 268**R-N-[2-Chloro-4-[N-ethyl-N-(pyrazin-2-yl)aminosulphonyl]phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide**

Potassium carbonate (64 mg, 0.65 mmol) and R-N-[2-chloro-4-(pyrazin-2-ylamino-sulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide (Example 264) (203 mg, 0.47 mmol) were added to acetone (10 ml). Ethyl iodide (0.078 ml, 0.98 mmol) was added to the resulting suspension and the mixture was stirred at room temperature for 96h before being concentrated under vacuum. 1M HCl (15 ml) was added to the residue which was then extracted with EtOAc (2x15 ml). The combined extracts were washed with brine (25 ml), dried and concentrated under vacuum to leave a foam which was purified by column chromatography to yield the title compound as a foam (78 mg, 0.17 mmol). NMR: 1.04 (t, 3H), 1.60 (s, 3H), 3.80 (q, 2H), 7.67 (d, 1H), 7.85 (s, 1H), 8.05 (br s, 1H), 8.28 (d, 1H), 8.48 (s, 1H), 8.57 (s, 1H), 8.78 (s, 1H), 9.89 (br s, 1H); MS: 451.

Example 269**R-N-{2-Chloro-4-[N-ethyl-N-(5-trifluoromethylpyrid-2-yl)aminosulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide**

Potassium carbonate (74 mg, 0.75 mmol) and R-N-[2-chloro-4-(5-trifluoromethylpyrid-2-yl)aminosulphonyl]phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide (Example 249) (246mg, 0.51 mmol) were added to acetone (10 ml). Ethyl iodide (0.045 ml, 0.055 mmol) was added to the resulting suspension and the mixture was stirred at room temperature for 22h. Ethyl iodide (0.045 ml, 0.055 mol) was added to the mixture which was then stirred at room temperature for a further 46h before being concentrated under vacuum. 1M HCl (5 ml) and EtOAc (1ml) were added to the residue, which was absorbed onto a Bond Elut column and all organic components of the mixture were washed off with EtOAc. This resulting solution was concentrated under vacuum to leave a residue that was purified by column chromatography to yield the title compound as a foam (81 mg, 0.16 mmol). NMR: 1.18 (m, 3H), 1.60 (s, 3H), 4.00 (m, 2H), 7.68 (d, 1H), 7.80 (d, 1H), 7.96 (s, 1H), 8.25 (d, 1H), 8.29 (d, 1H), 8.76 (s, 1H), 9.85 (br s, 1H); MS: 518.

Example 270**R-N-[2-Chloro-4-(4-hydroxyanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide**

A solution of R-N-[2-chloro-4-(4-benzyloxyanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide (Example 39) (1.85 g, 3.5 mmol) in EtOAc (200 ml) was hydrogenated over 10% Pd/ C for 4h at ambient temperature. The catalyst was removed by filtration and the EtOAc evaporated. The residue was purified by column chromatography using 10% EtOAc in DCM to yield the title compound as a foam (1.55 g, 3.5 mmol). NMR: 1.6 (s, 3H), 6.6 (d, 2H), 6.8 (d, 2H), 7.6 (dd, 1H), 7.75 (d, 1H), 8.2 (d, 1H); MS: 437.

Example 271**S-N-{2-chloro-4-[(4-sulphamoylanilino)sulphonyl]phenyl}-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide**

- A solution of R-3,3,3-trifluoro-2-hydroxy-2-methylpropanoylchloride (Method C1) (223 mg, 1.25mmol) in DCM (10 ml) was added to a stirred mixture of 2-chloro-4-[(4-sulphamoylanilino)sulphonyl]aniline (Method A1) (362 mg, 1.0mmol) and 2,6-di *tert* butylpyridine (0.28 ml, 1.25mmol) in EtOAc (25 ml). The resultant mixture was stirred at ambient temperature overnight, evaporated to dryness, the residue dissolved in EtOAc and washed with 1M aqueous hydrochloric acid, aqueous sodium hydrogen carbonate solution and brine, dried and evaporated to dryness.
- 10 The residue was purified by column chromatography using 25% EtOAc in hexane to give, as a foam, the title compound (270 mg, 0.54mmol). EA found: C, 39.4; H, 3.4; N, 7.4; S, 11.5%; C₁₆H₁₅N₃F₃ClS₂O₆ 0.5 EtOAc requires: C, 39.6; H, 3.5; N, 7.7; S, 11.7%; NMR: 1.6 (s, 3H), 7.2 (s, 2H), 7.25 (d, 2H), 7.65 (d, 2H), 7.8 (dd, 1H), 7.95 (d, 1H), 8.2 (d, 1H); MS: 500.

15 Example 272**R-N-[2-chloro-5-(2-chloroanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide**

- A solution of S-3,3,3-trifluoro-2-hydroxy-2-methylpropanoylchloride (Method P) (95 mg, 0.53mmol) in DCM (5 ml) was added to a stirred mixture of 2-chloro-5-(2-chloroanilinosulphonyl)aniline (Method B1) (140 mg, 0.44mmol) and 2,6-di-*t*-butylpyridine (0.12 ml, 0.53mmol) in DCM (10 ml). The resultant mixture was stirred at ambient temperature overnight, evaporated to dryness, the residue dissolved in EtOAc and washed with 1M aqueous hydrochloric acid and brine, dried and evaporated to dryness. The residue was purified by column chromatography using 5% EtOAc in DCM to give, as a foam, the title compound (178 mg,
- 20 0.39mmol). NMR: 1.6 (s, 3H), 7.2 (m, 3H), 7.4 (d, 1H), 7.45 (dd, 1H), 7.75 (d, 1H), 8.4 (d, 1H); MS: 455.
- 25

Preparation of Starting Materials

- The starting materials for the Examples above are either commercially available or are
- 30 readily prepared by standard methods from known materials. For example the following reactions

(Methods A-B1) are illustrations but not limitations of the preparation of some of the starting materials used in the above reactions.

Method A

5 2-Chloro-4-(morpholin Sulphonyl)anilino

A solution of *N*-acyl-2-chloro-4-chlorosulphonylaniline (536 mg, 2 mmol), morpholine (0.2 ml, 2.3 mmol), 4-dimethylaminopyridine (10 mg, 0.08 mmol) and pyridine (0.19 ml, 2.3 mmol) in DCM (25 ml) was stirred at ambient temperature for 4h. The mixture was washed with water and brine, dried and evaporated to dryness. The residue was dissolved in ethanol (25 ml) and 2M aqueous sodium hydroxide (5 ml) was added. The mixture was heated at reflux for 4h and then cooled to ambient temperature, evaporated to dryness and water (25 ml) was added to the residue. The aqueous solution was extracted with EtOAc; the organic phase was separated then washed with water, dried and evaporated to dryness to yield the title compound (500 mg, 1.8 mmol). MS: 275.

Method B

***N*-(2-Chloro-4-fluorosulphonylphenyl)-2-acetoxy-2-methylpropanamide**

A solution of 2-chloro-4-fluorosulphonylaniline (2.5 g, 12 mmol), 2-acetoxyisobutyryl chloride (2.35 g, 14 mmol) and pyridine (1.2 ml, 14.5 mmol), in DCM (50 ml) was stirred at ambient temperature for 18h. The mixture was washed with water, dried and evaporated to dryness. The residue was purified by column chromatography using 25% EtOAc/isohexane and the oily product, the title compound (2.0 g, 5.9 mmol), solidified on standing. MS: 336.

Method C

25 2-Chloro-4-(3-bromo-4-chloroanilinosulphonyl)aniline

A solution of *N*-acyl-2-chloro-4-chlorosulphonylaniline (370 mg, 1.38 mmol), 3-bromo-4-chloroaniline (250 mg, 1.2 mmol), 4-dimethylaminopyridine (10 mg) and pyridine (0.11 ml, 1.3 mmol) in DCM (10 ml) was stirred at ambient temperature for 4h. The solution was washed with water and brine, dried and evaporated to dryness. The residue was dissolved in ethanol (25 ml) and 2M aqueous sodium hydroxide (5 ml) was added. The mixture heated at reflux for 4h and

then cooled to ambient temperature, evaporated to dryness and water (25 ml) was added to the residue. The aqueous solution was extracted with EtOAc; the separated organic phase was washed with water, dried and evaporated to dryness to yield the title compound (500 mg, 1.1 mmol). MS: 435.

5

Method D

2-Chloro-4-(4-methoxyvanilinosulphonyl)aniline

A solution of *N*-acyl-2-chloro-4-chlorosulphonylaniline (536 mg, 2 mmol), 4-methoxyaniline (246 mg, 2 mmol), 4-dimethylaminopyridine (10 mg, 0.08 mmol) and pyridine
10 (0.19 ml, 2.3 mmol) in DCM (25 ml) was stirred at ambient temperature for 4h. The solution was washed with 1M aqueous hydrochloric acid and brine, dried and evaporated to dryness. The residue was dissolved in ethanol (25 ml) and 2M aqueous sodium hydroxide (5 ml) was added. The mixture heated at reflux for 4h and then cooled to ambient temperature and evaporated to dryness. Water (25 ml) was added to the residue and the solution was neutralised to pH 7.0 by
15 addition of 1M aqueous hydrochloric acid. The aqueous solution was extracted with EtOAc, the EtOAc extracts were washed with brine, dried and evaporated to dryness to yield the title compound (620 mg, 1.98 mmol). MS: 311.

Method E

2-Chloro-4-(allylaminosulphonyl)aniline

A solution of *N*-acyl-2-chloro-4-chlorosulphonylaniline (670 mg, 2.5 mmol) and allylamine (342 mg, 5.7 mmol), in DCM (25 ml) was stirred at ambient temperature for 4h. The mixture was washed with 1M aqueous hydrochloric acid and brine, dried and evaporated to dryness. The residue was dissolved in ethanol (25 ml) and 2M aqueous sodium hydroxide (5 ml)
25 was added. The mixture heated at reflux for 4h and then cooled to ambient temperature and evaporated to dryness. Water (25 ml) was added to the residue and the solution was neutralised to pH 7.0 by addition of 1M aqueous hydrochloric acid. The aqueous solution was extracted with EtOAc. The EtOAc extracts were washed with brine, dried and evaporated to dryness to yield the title compound (550 mg, 2.2 mmol). MS: 245.

Method F**2-Chloro-4-(thien-2-ylmethylaminosulphonyl)aniline**

A solution of *N*-acyl-2-chloro-4-chlorosulphonylaniline (320 mg, 1.2 mmol), 4-dimethylaminopyridine (10 mg, 0.08 mmol), pyridine (0.1 ml, 1.2 mmol) and 2-thiophenemethylamine (120 mg, 1.08 mmol) in DCM (8 ml) was stirred at ambient temperature for 18h. The mixture was washed with 1M aqueous hydrochloric acid and then the organic phase was concentrated. The residue was dissolved in ethanol (7 ml), 2M aqueous sodium hydroxide (2 ml) was added and the mixture was heated at 60°C for 6h. The solvent was evaporated and the residue dissolved in EtOAc, washed with water, and the solvent phase was concentrated to yield the title compound which was used, without further purification.

Method G**2-Fluoro-4-(2-fluoroanilinosulphonyl)aniline**

Method D was followed except that *N*-acyl-2-fluoro-4-chlorosulphonylaniline (Method H) was used in place of *N*-acyl-2-chloro-4-chlorosulphonylaniline, 2-fluoroaniline was used in place of 4-methoxyaniline and no 4-dimethylaminopyridine was used to yield the title compound.

Method H**20 *N*-Acyl-2-fluoro-4-chlorosulphonylaniline**

N-acyl-2-fluoro-4-sulphoaniline triethylamine salt (1:1) (Method I) (39 g, 0.12 mol) was added portion-wise, over 30mins, to POCl₃ (60 ml) at 0°C. The reaction mixture was stirred at room temperature for 15h and then poured slowly onto a stirred solution of ice-water. After stirring for 15mins the mixture was filtered to yield the title compound as a solid (26 g, 0.10 mol). NMR (CDCl₃): 2.25 (s, 3H), 7.55 (br s, 1H), 7.7 (dd, 1H), 7.75-7.80 (m, 1H), 8.65 (t, 1H); MS: 190.

Method I**N-Acyl-2-fluoro-4-sulphoaniline triethylamine salt (1:1)**

2-Fluoroaniline (40 g, 0.36 mol) was added dropwise to a stirred solution of concentrated sulphuric acid (60 ml) and the mixture heated to 190°C for 15h. The reaction mixture was then
5 cooled to room temperature and poured slowly onto a stirred solution of ice-water. After stirring for 15mins the mixture was filtered to yield a solid, 2-fluoro-4-sulphoaniline (41 g). NMR: 7.05 (t, 1H), 7.20-7.30 (m, 2H), 8.1 (br s, 3H); MS: 190 (M-H). The 2-fluoro-4-sulphoaniline (41 g) was dissolved in acetic anhydride (70 ml) and stirred in an ice bath. Triethylamine (22 g, 0.22 mol) was added very slowly with vigorous stirring (and concomittant release of heat). The
10 reaction mixture was left to stir for 14h, at which point a solid formed; this was filtered to yield the title compound (39 g, 0.12 mol). MS: 232.

Method J**1-(*t*-Butoxycarbonyl)-4-(4-amino-3-methylbenzene)sulphonylpiperazinide**

15 Method D was followed except that *N*-acyl-2-methyl-4-chlorosulphonylaniline (Method K) was used in place of *N*-acyl-2-chloro-4-chlorosulphonylaniline, 1-(*t*-butyloxycarbonyl) piperazine was used in place of 4-methoxyaniline and no 4-dimethylaminopyridine was used to yield the title compound.

20 Method K**N-Acyl-2-methyl-4-chlorosulphonylaniline**

N-Acyl-2-methyl-4-sulphoaniline triethylamine salt (1:1) (Method L) (35 g, 0.11 mol) was added portion-wise, over 30mins, to POCl₃ (50 ml) at 0°C. The reaction mixture was stirred at room temperature for 15h and then poured slowly onto a stirred solution of ice-water. After
25 stirring for 15mins the mixture was filtered to yield the title compound as a solid (25 g, 0.10 mol). MS: 246.

Method L**N-Acyl-2-methyl-4-sulphoaniline triethylamine salt (1:1)**

2-Methyl-4-sulphoaniline (30 g, 0.16 mol) was dissolved in acetic anhydride (50 ml) and stirred in an ice bath. Triethylamine (23 ml, 0.18 mol) was added very slowly with vigorous stirring (and concomittant release of heat). The reaction mixture was left to stir for 14h, at which point a solid formed; this was filtered to yield the title compound (35 g, 0.11 mol). NMR: 1.15 (t, 9H); 2.05 (s, 3H); 2.45-2.50 (m, 1H), 3.1 (q, 6H), 7.25-7.40 (m, 2H), 7.85 (t, 1H), 9.7 (br s, 1H);

MS: 228.

10 Method M**R-N-(2-Chloro-4-chlorosulphonylphenyl)-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide**

R-N-(2-Chlorophenyl)-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide (Method N) (13.8 g, 52 mmol) was added in portions to a cooled (0°C) solution of chlorosulphonic acid (25 ml) over 15mins and then the mixture was heated to 85°C. After 4.5h the reaction mixture was cooled in an ice bath and then poured very slowly onto a stirred ice-water mixture. After stirring for 15mins, the mixture was extracted with EtOAc (2x100 ml) and the combined organic layer washed with brine, dried and concentrated to yield a brown oil. This oil was purified by flash column chromatography using 10: 1, *iso*-hexane: EtOAc to yield the title compound as a pale yellow solid (11 g, 30 mmol). NMR: 1.6 (s, 3H), 7.55 (dd, 1H), 7.6 (d, 1H), 7.95 (d, 1H), 9.7 (br s, 1H); MS: 364.

Method N**R-N-(2-Chlorophenyl)-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide**

Acetyl chloride (11.7 ml, 164 mmol) was added dropwise to a stirred solution of the R-3,3,3-trifluoro-2-hydroxy-2-methylpropanoic acid (10 g, 63 mmol) in toluene (100 ml) cooled in an ice bath. The mixture was then heated to 80°C and the suspension dissolved to yield a clear solution. After 2h the reaction mixture was cooled and then concentrated to yield a slight brown oil. This oil was then redissolved in DCM (140 ml) and DMF (4 drops) was added followed by oxalyl chloride (6 ml, 69 mmol). The solution bubbled vigorously and the reaction mixture was left to stir. After 15h, this reaction mixture was added slowly to a stirred solution of 2-

chloroaniline (8.7 g, 68 mmol) and pyridine (5.5 ml, 68 mmol) in DCM (150 ml). After 15h stirring at room temperature, the resultant mixture was concentrated and the residue dissolved in methanol (500 ml). A solution of lithium hydroxide monohydrate (7.8 g, 0.19 mol) in water (120 ml) was then added and the mixture was stirred for 4h. The mixture was then concentrated and the residue acidified to pH 2 (by addition of concentrated hydrochloric acid). EtOAc (150 ml) was added and the mixture washed with water (2x100 ml) and brine, dried and evaporated to dryness. The residue was purified by flash column chromatography using 6: 1, *iso*-hexane: EtOAc to yield the title compound as a white solid (13.8 g, 52 mmol). NMR: 1.6 (s, 3H), 7.1-7.25 (m, 1H), 7.3-7.4 (m, 1H), 7.55 (dd, 1H), 7.8 (s, 1H), 8.0 (dd, 1H), 9.7 (br s, 1H); MS: 266.

10

Method O

R-N-[2-Chloro-4-(4-fluorosulphonylanilinosulphonyl)phenyl]-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide

A solution of R-N-(2-chloro-4-chlorosulphonylphenyl)-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide (Method M) (3.46 g, 9.45 mmol) in DCM (200 ml) was added to a stirred solution of 4-fluorosulphonylaniline (1.99 g, 11.3 mmol) and pyridine (1.52 ml, 18.2 mmol) in DCM (50 ml). The resultant mixture was stirred at ambient temperature overnight, evaporated to dryness and the residue treated with 1M aqueous hydrochloric acid (50 ml). The aqueous solution was extracted with EtOAc, the EtOAc extracts were washed with brine, dried and evaporated. The residue was purified by column chromatography using 30% EtOAc in DCM to yield the title compound as a foam (4.56 g, 9.05 mmol). NMR: 1.6 (s, 3H), 7.45 (d, 2H), 7.9 (dd, 1H), 8.0 (d, 1H), 8.05 (d, 2H), 8.3 (d, 1H); MS: 504.

Method P

S-3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylchloride

Oxalyl chloride (1.07 ml, 12 mmol) was added dropwise to a stirred suspension of (R)-(+)-2-hydroxy-2-methyl-3,3,3-trifluoropropanoic acid (Method Q) (1.95 g, 12 mmol) in DCM (42 ml) and DMF (0.8 ml). The mixture was stirred at ambient temperature for 2-15h to yield a solution of the title compound which was used in subsequent reactions without further purification.

Method Q**(R)-(+)-2-Hydroxy-2-methyl-3,3,3-trifluoropropanoic acid**

R/S-2-Hydroxy-2-methyl-3,3,3-trifluoropropanoic acid was resolved according to the
5 resolution method described in European Patent Application No. EP 524781 (described for the
preparation of the (S)-(-) acid) except that (1S, 2R)-norephedrine was used in place of (1R, 2S)-
norephedrine or (S)-(-)-1-phenylethylamine to yield the title compound, $[\alpha]_{\text{D}}^{20} +18.1^{\circ}$ (c, 8.8 in
MeOH); NMR analysis of the acid in the presence of (R)-(+)-1-phenylethylamine gave an
enantiomeric purity of >98%. NMR (CDCl_3): 1.27 (s, 3H) for the (R)-enantiomer, 1.21 (s, 3H)
10 for the (S)-enantiomer.

Method R**2-Chloro-4-[4-(2-hydroxyethylthio)anilinosulphonyl]aniline**

A solution of 4-acetamidothiophenol (8.35 g, 50.0 mmol), ethylene carbonate (5.5 g, 62.5
15 mmol) and sodium ethoxide (4.1 g, 60.0 mmol) in ethanol (250 ml) was heated at reflux
overnight and then cooled to ambient temperature and evaporated to dryness. The residue was
dissolved in diethyl ether (500 ml), washed with 1M aqueous hydrochloric acid and brine, dried
and evaporated to dryness. The residue was dissolved in ethanol (120 ml) and 2M aqueous
sodium hydroxide (60 ml) was added. The mixture was heated at reflux overnight and then
20 cooled to ambient temperature and evaporated to dryness. Water (25 ml) was added to the residue
and the solution was neutralised to pH 7 by the addition of 1M aqueous hydrochloric acid. The
aqueous solution was extracted with EtOAc, the EtOAc extracts were washed with brine, dried
and evaporated to dryness. The residue was dissolved in DCM (100 ml) and added to a solution
of 4-acetamido-3-chlorobenzenesulphonyl chloride (5.5 g, 20.5 mmol), 4-dimethylaminopyridine
25 (50 mg, 0.4 mmol) and pyridine (5.0 ml, 60 mmol) in DCM (100 ml). The mixture was stirred at
ambient temperature overnight, the solution was evaporated to dryness, the residue dissolved in
diethyl ether (450 ml), washed with 1M aqueous hydrochloric acid and brine, dried and
evaporated to dryness. The residue was dissolved in ethanol (100 ml) and 2M aqueous sodium
hydroxide (25 ml) was added. The mixture was heated at reflux overnight and then cooled to
30 ambient temperature and evaporated to dryness. Water (25 ml) was added to the residue and the

solution was neutralised to pH 7 by the addition of 1M aqueous hydrochloric acid. The aqueous solution was extracted with EtOAc, the EtOAc extracts were washed with brine, dried and evaporated to dryness. The residue was purified by column chromatography using 25% EtOAc in DCM to yield the title compound (630 mg, 1.8 mmol). NMR: 2.9 (t, 2H), 3.5 (m, 2H), 4.8 (t, 1H), 5.6.2 (s, 2H), 6.75 (d, 1H), 7.0 (d, 2H), 7.2 (d, 2H), 7.3 (dd, 1H), 7.45 (d, 1H); MS: 357.

Method S

2-Chloro-4-[4-(2-ethoxyethylthio)anilinosulphonyl]aniline

A solution of 4-acetamidothiophenol (8.35 g, 50.0 mmol), ethylene carbonate (5.5 g, 62.5 mmol) and sodium ethoxide (4.1 g, 60.0 mmol) in ethanol (250 ml) was heated at reflux overnight and then cooled to ambient temperature and evaporated to dryness. The residue was dissolved in diethyl ether (500 ml), washed with 1M aqueous hydrochloric acid and brine, dried and evaporated to dryness. The residue was dissolved in ethanol (120 ml) and 2M aqueous sodium hydroxide (60 ml) was added. The mixture was heated at reflux overnight and then cooled to ambient temperature and evaporated to dryness. Water (25 ml) was added to the residue and the solution was neutralised to pH 7 by the addition of 1M aqueous hydrochloric acid. The aqueous solution was extracted with EtOAc, the EtOAc extracts were washed with brine, dried and evaporated to dryness. The residue was dissolved in DCM (100 ml) and added to a solution of 4-acetamido-3-chlorobenzenesulphonyl chloride (5.5 g, 20.5 mmol), 4-dimethylaminopyridine (50 mg, 0.4 mmol) and pyridine (5.0 ml, 60 mmol) in DCM (100 ml). The mixture was stirred at ambient temperature overnight, the solution was evaporated to dryness, the residue dissolved in diethyl ether (450 ml), washed with 1M aqueous hydrochloric acid and brine, dried and evaporated to dryness. The residue was dissolved in ethanol (100 ml) and 2M aqueous sodium hydroxide (25 ml) was added. The mixture was heated at reflux overnight and then cooled to ambient temperature and evaporated to dryness. Water (25 ml) was added to the residue and the solution was neutralised to pH 7 by the addition of 1M aqueous hydrochloric acid. The aqueous solution was extracted with EtOAc, the EtOAc extracts were washed with brine, dried and evaporated to dryness. The residue was purified by column chromatography using 10% EtOAc in DCM to yield the title compound (1.53 g, 3.96 mmol). NMR: 1.0 (t, 3H), 3.0 (t, 2H), 3.3-3.5 (m, 4H), 6.2 (s, 2H), 6.75 (d, 1H), 7.0 (d, 2H), 7.2 (d, 2H), 7.3 (dd, 1H), 7.45 (d, 1H); MS: 385.

Method T**2-Chloro-4-[4-(methoxycarbonyl)anilinosulphonyl]aniline**

A solution of *N*-acyl-2-chloro-4-chlorosulphonylaniline (13.4 g, 50.0 mmol), methyl 4-aminobenzoate (7.55 g, 50.0 mmol), 4-dimethylaminopyridine (100 mg, 0.8 mmol) and pyridine (5.0 ml, 60 mmol) in DCM (350 ml) was stirred at ambient temperature for 4h. The solution was evaporated to dryness, the residue dissolved in EtOAc (1.0 L), washed with 1M aqueous hydrochloric acid and brine, dried and evaporated to dryness. The residue was dissolved in methanol (600 ml) and 2M aqueous sodium hydroxide (137 ml) was added. The mixture was heated at reflux for 4h and then cooled to ambient temperature and evaporated to dryness. Water (25 ml) was added to the residue and the solution was acidified to pH 1 by the addition of concentrated aqueous hydrochloric acid. The solution was evaporated to dryness, the residue dissolved in methanol (200 ml), the solution cooled to -35°C and treated with thionyl chloride (18.0 ml). The mixture was stirred at ambient temperature overnight and evaporated to dryness. The residue was dissolved in saturated sodium hydrogen carbonate solution, the aqueous solution was extracted with EtOAc, the EtOAc extracts were washed with brine, dried and evaporated to dryness to yield the title compound (13.6 g, 40.0 mmol); MS: 339.

Method U**2-Chloro-4-[1-(*t*-butoxycarbonyl)piperazin-4-ylsulphonyl]aniline**

A solution of *N*-acyl-2-chloro-4-chlorosulphonylaniline (10.0 g, 37.3 mmol), 1-(*t*-butoxycarbonyl)piperazine (7.0 g, 37.6 mmol) and triethylamine (7.77 ml, 56.4 mmol) in DCM (150 ml) was stirred at ambient temperature for 4h. The solution was evaporated to dryness, the residue dissolved in EtOAc (150 ml), washed with 0.7M aqueous citric acid, saturated aqueous sodium hydrogen carbonate solution and brine, dried and evaporated to dryness. The residue was dissolved in ethanol (100 ml) and 2M aqueous sodium hydroxide (87.5 ml) was added. The mixture was heated at reflux overnight and then cooled to ambient temperature and evaporated to dryness. Water (75 ml) was added to the residue and the mixture was filtered, the solid washed with water, dried and crystallised from EtOAc to yield the title compound as a solid (11.4 g, 30.3

mmol); M.pt. 243-244°C. NMR: 1.35 (s, 9H), 2.8 (m, 4H), 3.4 (m, 4H), 6.35 (s, 2H), 6.9 (d, 1H), 7.35 (dd, 1H), 7.5 (d, 1H); MS: 374.

Method V

5 2-(4-Aminophenyl)-1,3-dioxolane

To a mixture of 4-nitrobenzaldehyde (7.55 g, 50 mmol) and ethyleneglycol (4.65 g, 75.0 mmol) in toluene (125 ml) was added 4-toluenesulphonic acid (100 mg, 0.5 mmol) and the mixture heated under reflux using a Dean-Stark water separator for 2h. The solution was evaporated to dryness, the residue dissolved in EtOAc (125 ml), washed with saturated sodium hydrogen carbonate solution, dried and evaporated to dryness. The residue was dissolved in ethanol (200 ml) and hydrogenated over 10% Pd/ C for 4h at ambient temperature. The catalyst was removed by filtration, the ethanol evaporated to yield the title compound which was used without further purification.

15 Method W

4-(2-Hydroxyethylamino)aniline

A mixture of ethanolamine (2.29 g, 37.5 mmol) and 1-fluoro-4-nitrobenzene (3.53 g, 25.0 mmol) in dimethylsulphoxide (25 ml) was treated with potassium carbonate (6.9 g, 50 mmol) and heated at 90°C overnight. The mixture was cooled and poured onto water (250 ml), the precipitated solid filtered off, dried, dissolved in ethanol (50 ml) and hydrogenated over 10% Pd/ C for 4h at ambient temperature. The catalyst was removed by filtration, the ethanol evaporated and the residue purified by column chromatography using EtOAc to yield the title compound (740 mg, 5.0 mmol). NMR: 2.95 (t, 2H), 3.5 (t, 2H), 4.1 (s, 2H), 4.4 (s, 1H), 4.55 (t, 1H), 6.4 (m, 4H); MS: 153 (M+H)⁺.

25

Method X

4-(N-2-Tetrahydropyranyloxyethyl-*t*-butoxycarbonylamino)aniline

To a mixture of *N-t*-butoxycarbonyl ethanolamine (7.7 ml, 50 mmol) and 3,4-dihydropyran (5.7 ml, 62.5 mmol) in DCM (125 ml) was added 4-toluenesulphonic acid (100 mg, 0.5 mmol) and the mixture stirred at ambient temperature for 3h. The solution was washed with

saturated sodium hydrogen carbonate solution, dried and evaporated to dryness. The residue was filtered through a short pad of basic alumina with 10% EtOAc in hexane, the solvent evaporated to dryness, the residue dissolved in dimethylacetamide (5 ml) and added to a suspension of sodium hydride (400 mg of 50% dispersion in mineral oil, 10 mmol) in dimethylacetamide (5 ml). The mixture was heated at 50°C for 1h, cooled to ambient temperature and added to a solution of 1-fluoro-4-nitrobenzene (1.06 g, 7.5 mmol) in dimethylacetamide (5 ml) and heated at 50°C for 4h. The cooled mixture was poured onto water (150 ml), the aqueous solution was extracted with EtOAc, the EtOAc extracts were washed with brine, dried and evaporated. The residue was dissolved in ethanol (50 ml) and hydrogenated over 10% Pd/ C for 4h at ambient temperature. The catalyst was removed by filtration, the ethanol evaporated and the residue purified by column chromatography using 30% EtOAc in hexane to yield the title compound (640 mg, 1.9 mmol). NMR: 1.2-1.6 (m, 15H), 3.4-3.8 (m, 6H), 6.6 (d, 2H), 7.0 (d, 2H); MS: 337 (M+H)⁺.

15 Method Y

R-N-(2-chloro-4-chlorosulphonylphenyl)-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide

R-N-(2-chlorophenyl)-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide (Method Z) (4.71 g, 17.6 mmol) was added portion wise to a cooled (ice bath) and stirred solution of chlorosulphonic acid (9 ml). The resulting mixture was heated to 85°C for 270mins. The reaction mixture was then cooled to room temperature and poured onto an ice/ water slurry. The resulting aqueous mixture was extracted with EtOAc. The combined extracts were washed with brine, dried and were concentrated to yield an oil. Purification by column chromatography yielded the title compound as a solid (4.03 g, 11.0 mmol). NMR (CDCl₃): 1.80 (s, 3H), 3.46 (s, 1H), 8.00 (dd, 1H), 8.10 (d, 1H), 8.77 (d, 1H), 9.44 (s, 1H); MS: 364.

25

Method Z

R-N-(2-chlorophenyl)-3,3,3-trifluoro-2-hydroxy-2-methylpropanamide

2-Chloroaniline (3.2 ml) and 2,6-di-*t*-butylpyridine (7.4 ml) were added to a solution of S-3,3,3-trifluoro-2-hydroxy-2-methylpropanoyl chloride [produced *in situ* by reaction of R-3,3,3-trifluoro-2-hydroxy-2-methylpropanoic acid (4.74 g, 30 mmol) with 2M oxalyl chloride (16.5 ml,

30

33 mmol) in DCM (50 ml)] in DCM. The resulting solution was stirred for 20h at room temperature. The reaction mixture was then washed with water, brine, dried, evaporated and purified by column chromatography to yield the title compound as a solid (4.76 g, 17.8 mmol). NMR: 1.60 (s, 3H), 7.25 (t, 1H), 7.38 (t, 1H), 7.55 (d, 1H), 7.80 (s, 1H), 8.00 (d, 1H), 9.70 (s, 5 1H); MS: 266.

Method A1

2-chloro-4-[(4-sulphamoyl-anilino)sulphonyl]aniline

A solution of 4-acetamido-3-chlorobenzenesulphonyl chloride (2.68 g, 10.0mmol),
10 sulphanilamide (2.06 g, 12.0mmol), dimethylaminopyridine (1.22 g, 10.0mmol) and pyridine (2.40 ml, 10.0mmol) in EtOAc (150 ml) was stirred at ambient temperature for 3 days. The solution was washed with 1.0M aqueous hydrochloric acid and brine, dried and evaporated to dryness. The residue was dissolved in ethanol (100 ml) and 2M aqueous sodium hydroxide (22.1 ml) was added. The mixture was heated at reflux for 4 hours and then cooled to ambient
15 temperature and evaporated to dryness. Water (25 ml) was added to the residue and the solution was neutralised to pH 7.0 by the addition of 1M aqueous hydrochloric acid. The aqueous solution was extracted with EtOAc, the EtOAc extracts washed with brine, dried and evaporated to give, as a solid, the title compound (2.34 g, 6.5mmol). M.p. 197°-198°; NMR: 6.3 (s, 2H), 6.8 (d, 1H), 7.2 (s, 2H), 7.25 (d, 2H), 7.4 (dd, 2H), 7.55 (d, 1H), 7.65 (d, 2H); MS: 360.

20

Method B1

2-chloro-5-(2-chloroanilino-sulphonyl)aniline

A solution of 3-nitro-4-chlorobenzenesulphonyl chloride (768 mg, 3.0mmol), 2-chloroaniline (383 mg, 3.0mmol), 4-dimethylaminopyridine (10 mg, 0.08mmol) and pyridine
25 (0.30 ml, 3.6mmol) in DCM (35 ml) was stirred at ambient temperature overnight. The solution was evaporated to dryness, the residue dissolved in EtOAc and washed with 1.0M aqueous hydrochloric acid and brine, dried and evaporated to dryness. The residue was dissolved in EtOAc (30 ml) and hydrogenated over 10% Pd/C for 4h at ambient temperature. The catalyst was removed by filtration and the EtOAc evaporated to dryness. The residue was purified by column

chromatography on basic alumina using 5% methanol in EtOAc to give, as a solid, the title compound (140 mg, 0.44mmol). MS: 315.

Method C1

5 R-3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylchloride

Oxalyl chloride (1.07 ml, 12 mmol) was added dropwise to a stirred suspension of (S)-(-)-2-hydroxy-2-methyl-3,3,3-trifluoropropanoic acid (Method D1) (1.95 g, 12 mmol) in DCM (42 ml) and DMF (0.8 ml). The mixture was stirred at ambient temperature for 2-15h to yield a solution of the title compound which was used in subsequent reactions without further
10 purification.

Method D1

(S)-(-)-2-Hydroxy-2-methyl-3,3,3-trifluoropropanoic acid

R/S-2-Hydroxy-2-methyl-3,3,3-trifluoropropanoic acid was resolved according to the
15 resolution method described in European Patent Application No. EP 524781 to yield the title compound, $[\alpha]_D^{20} -18.6^\circ$ (c, 8.8 in MeOH).

Method E1

R/S-3,3,3-Trifluoro-2-hydroxy-2-methylpropanoylchloride

20 Oxalyl chloride (1.07 ml, 12 mmol) was added dropwise to a stirred suspension of R/S-2-hydroxy-2-methyl-3,3,3-trifluoropropanoic acid (1.95 g, 12 mmol) in DCM (42 ml) and DMF (0.8 ml). The mixture was stirred at ambient temperature for 2-15h to yield a solution of the title compound which was used in subsequent reactions without further purification.

25 Example 273

The following illustrate representative pharmaceutical dosage forms containing the compound of formula I, or a pharmaceutically acceptable salt or an *in vivo* hydrolysable ester thereof (hereafter compound X), for therapeutic or prophylactic use in humans:

(a) <u>Tablet I</u> <u>mg/tablet</u>		
	Compound X	100
	Lactose Ph.Eur.	182.75
5	Croscarmellose sodium	12.0
	Maize starch paste (5% w/v paste)	2.25
	Magnesium stearate	3.0
(b) <u>Tablet II</u> <u>mg/tablet</u>		
10	Compound X	50
	Lactose Ph.Eur.	223.75
	Croscarmellose sodium	6.0
	Maize starch	15.0
	Polyvinylpyrrolidone (5% w/v paste)	2.25
15	Magnesium stearate	3.0
(c) <u>Tablet III</u> <u>mg/tablet</u>		
	Compound X	1.0
	Lactose Ph.Eur.	93.25
20	Croscarmellose sodium	4.0
	Maize starch paste (5% w/v paste)	0.75
	Magnesium stearate	1.0
(d) <u>Capsule</u> <u>mg/capsule</u>		
25	Compound X	10
	Lactose Ph.Eur.	488.5
	Magnesium stearate	1.5

5	(e)	<u>Injection I</u>	<u>(50 mg/ml)</u>
		Compound X	5.0% w/v
		1M Sodium hydroxide solution	15.0% v/v
		0.1M Hydrochloric acid	(to adjust pH to 7.6)
		Polyethylene glycol 400	4.5% w/v
		Water for injection to 100%	
10	(f)	<u>Injection II</u>	<u>10 mg/ml)</u>
		Compound X	1.0% w/v
		Sodium phosphate BP	3.6% w/v
		0.1M Sodium hydroxide solution	15.0% v/v
		Water for injection to 100%	
15	(g)	<u>Injection III</u>	<u>(1mg/ml,buffered to pH6)</u>
		Compound X	0.1% w/v
		Sodium phosphate BP	2.26% w/v
		Citric acid	0.38% w/v
		Polyethylene glycol 400	3.5% w/v
20		Water for injection to 100%	


Note:

The above formulations may be obtained by conventional procedures well known in the pharmaceutical art. The tablets (a)-(c) may be enteric coated by conventional means, for example
25 to provide a coating of cellulose acetate phthalate.

CLAIMS:

What is claimed is:

1. A compound of the formula (I):



5

(I)

wherein:

Ring X is phenyl or a six membered heteroaryl ring containing one or two ring nitrogens where said nitrogens are optionally oxidised to form the N-oxide;

R¹ and **R²** are independently as defined in (a) or (b);

10 **R³** and **R⁴** are as defined in (c) or (d);

R⁵ is as defined in (e) or (f);

Y-Z is as defined in (g) or (h)

wherein:

(a) either **R¹** and **R²** are each selected independently from hydrogen, C₁₋₃alkyl, pyridyl and

15 phenyl which is unsubstituted or substituted by one or two substituents selected independently from C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkenyloxy, hydroxy, halo and cyano,

or **R¹** and **R²** together with the nitrogen atom to which they are attached form morpholino, thiomorpholino, piperidiny, pyrrolidinyl or imidazolyl;

(b) either **R¹** and **R²** are each selected independently from phenyl substituted by one or
20 more P (wherein P is as defined hereinbelow), phenyl substituted by one or more groups selected from C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkenyloxy, hydroxy, halo and cyano and additionally substituted by one or more groups selected from P, a heterocyclic group other than unsubstituted pyridyl which is optionally substituted on a ring carbon by one or more Q (wherein Q is as defined hereinbelow) and wherein if said heterocyclic group contains an -NH- moiety that nitrogen may

25 be optionally substituted by a group selected from D (wherein D is as defined hereinbelow), naphthyl optionally substituted by one or more Q, C₄₋₆alkyl, C₃₋₆cycloalkyl optionally substituted with one or more Q, C₂₋₆alkenyl, C₂₋₆alkynyl, C₁₋₆alkyl substituted by one or more V (wherein V is

as defined hereinbelow), R^6T - (wherein R^6 and T are as defined hereinbelow) and $R^7C_{1-6}alkylT$ - (where R^7 is as defined hereinbelow),

or R^1 and R^2 together with the nitrogen atom to which they are attached form a heterocyclic group other than unsubstituted morpholino, unsubstituted thiomorpholino,

5. unsubstituted piperidinyl, unsubstituted pyrrolidinyl or unsubstituted imidazolyl which is optionally substituted on a ring carbon by one or more Q (wherein Q is as defined hereinbelow) and wherein if said heterocyclic group contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from D (wherein D is as defined hereinbelow);

(c) either R^3 and R^4 are independently C_k alkyl optionally substituted by from 1 to $2k+1$ atoms selected from fluoro and chloro wherein k is 1-3, provided that R^3 and R^4 are not both methyl;

or R^3 and R^4 together with the carbon atom to which they are attached, form a C_m cycloalkyl ring optionally substituted by from 1 to $2m-2$ fluorine atoms wherein m is 3-5;

(d) R^3 and R^4 are both methyl;

(e) R^5 is hydrogen, C_{1-4} alkyl, halo C_{1-4} alkyl, C_{1-4} alkoxy, halo C_{1-4} alkoxy, cyano, nitro, C_{2-4} alkenyloxy or trifluoromethylthio;

(f) R^5 is halo, hydroxy, amino, C_{1-6} alkylamino, $(C_{1-6}alkyl)_2$ amino, C_{1-6} alkanoylamino, C_{1-6} alkanoyl(N- $C_{1-6}alkyl$)amino, $C_{1-6}alkylsulphonylamino$, $C_{1-6}alkylsulphonyl(N-C_{1-6}alkyl)amino$, thiol, $C_{1-6}alkylsulphanyl$, $C_{1-6}alkylsulphinyl$, $C_{1-6}alkylsulphonyl$, sulphamoyl,

N-($C_{1-6}alkyl$)aminosulphonyl, N-($C_{1-6}alkyl$) $_2$ aminosulphonyl, carboxy, carbamoyl, N-($C_{1-6}alkyl$)carbamoyl, N-($C_{1-6}alkyl$) $_2$ carbamoyl, C_{1-6} alkoxycarbonyl, formyl, C_{1-6} alkanoyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{5-6} alkyl, halo C_{5-6} alkyl, C_{5-6} alkoxy, halo C_{5-6} alkoxy or C_{5-6} alkenyloxy;

(g) Y-Z is -NHC(O)-, -OCH $_2$ -, -SCH $_2$ -, -NHCH $_2$ -, trans-vinylene, and ethynylene;

(h) Y-Z is -NHC(S)-;

R^6 is selected from C_{1-6} alkyl (optionally substituted with one or more R^8), C_{3-6} cycloalkyl optionally substituted with one or more R^8 , a heterocyclic group optionally substituted on a ring carbon by one or more R^8 and if said heterocyclic group contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from D (wherein D is as defined hereinbelow), phenyl optionally substituted by one or more R^8 , naphthyl optionally substituted by one or more R^8 ;

R^7 is a heterocyclic group optionally substituted on a ring carbon by one or more R^8 and if said heterocyclic group contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from D (wherein D is as defined hereinbelow), phenyl optionally substituted by one or more R^8 , naphthyl optionally substituted by one or more R^8 ;

5 R^8 is trifluoromethyl, C_{1-6} alkyl, halo, hydroxy, trifluoromethoxy, cyano, C_{1-6} alkoxy, formyl, C_{1-6} alkanoyl, C_{1-6} alkanoyloxy, amino, C_{1-6} alkylamino, $(C_{1-6}alkyl)_2$ amino, C_{1-6} alkanoylamino, C_{1-6} alkanoyl(N- $C_{1-6}alkyl$)amino, nitro, carboxy, carbamoyl, C_{1-6} alkoxycarbonyl, thiol, C_{1-6} alkylsulphanyl, C_{1-6} alkylsulphinyl, C_{1-6} alkylsulphonyl, sulphamoyl, N-($C_{1-6}alkyl$)aminosulphonyl, N-($C_{1-6}alkyl$)₂aminosulphonyl, carbamoyl $C_{1-6}alkyl$,
10 N-($C_{1-6}alkyl$)carbamoyl $C_{1-6}alkyl$, N-($C_{1-6}alkyl$)₂carbamoyl- $C_{1-6}alkyl$, hydroxy $C_{1-6}alkyl$, $C_{1-6}alkoxyC_{1-6}alkyl$, phenyl $C_{1-6}alkyl$ or phenyl $C_{1-6}alkoxy$;

P is selected from $C_{2-6}alkyl$ -M- substituted with one or more R^9 , $C_{2-6}alkenyl$ -M- optionally substituted with one or more R^9 , $C_{2-6}alkynyl$ -M- optionally substituted with one or more R^9 (with the proviso that in the three previous groups R^9 is not a substituent on the carbon atom attached to
15 M), R^{10} -CH₂-M-, R^{11} -M-, thiol, $C_{1-6}alkylsulphanyl$, $C_{1-6}alkylsulphinyl$, $C_{1-6}alkylsulphonyl$, sulphamoyl, nitro, carboxy, $C_{1-6}alkoxycarbonyl$, amino, $C_{1-6}alkylamino$, $(C_{1-6}alkyl)_2$ amino, carbamoyl, N-($C_{1-6}alkyl$)carbamoyl, N-($C_{1-6}alkyl$)₂carbamoyl, $C_{1-6}alkanoylamino$, $C_{1-6}alkanoyl$ (N- $C_{1-6}alkyl$)amino, trifluoromethyl, trifluoromethoxy, formyl, $C_{1-6}alkanoyl$, $C_{5-6}alkyl$, $C_{2-6}alkenyl$, $C_{2-6}alkynyl$, N-($C_{1-6}alkyl$)aminosulphonyl, hydroxymethyl, hydroxyacetyl,
20 N-($C_{1-6}alkyl$)₂aminosulphonyl, $C_{1-6}alkanoylaminosulphonyl$, $C_{1-6}alkanoyl$ (N- $C_{1-6}alkyl$)aminosulphonyl, $C_{1-6}alkylsulphonylaminocarbonyl$, $C_{1-6}alkylsulphonyl$ (N- $C_{1-6}alkyl$)aminocarbonyl, $C_{5-6}alkoxy$, $C_{5-6}alkenyloxy$, phenyl optionally substituted by one or more R^8 , naphthyl optionally substituted by one or more R^8 and a heterocyclic group optionally substituted on a ring carbon by one or more R^8 and if said heterocyclic group contains an -NH-
25 moiety that nitrogen may be optionally substituted by a group selected from D (wherein D is as defined hereinbelow);

Q is selected from any of the values defined for P, $C_{1-4}alkyl$, $C_{1-4}alkoxy$, $C_{2-4}alkenyloxy$, hydroxy, halo and cyano;

V is selected from any of the values defined for Q, phenyl optionally substituted by one or
30 more Q, naphthyl optionally substituted by one or more Q, a heterocyclic group optionally

substituted on a ring carbon by one or more Q and if said heterocyclic group contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from D (wherein D is as defined hereinbelow) or C₃₋₆cycloalkyl optionally substituted with one or more Q;

T is selected from -O-, -C(O)-, -NH-, -N(N-C₁₋₆alkyl)-, -C(O)NH-, -NHC(O)-, -C(O)N(N-C₁₋₆alkyl)-, -N(N-C₁₋₆alkyl)C(O)-, -SO₂-, -C(S)-, -C(S)NH-, -NHC(S)-, -C(S)N(N-C₁₋₆alkyl)- and -N(N-C₁₋₆alkyl)C(S)-;

M is selected from -O-, -N(R¹²)-, -C(O)-, -N(R¹²)C(O)-, -C(O)N(R¹²)-, -S(O)_n-, -OC(O)-, -C(O)O-, -N(R¹²)C(O)O-, -OC(O)N(R¹²)-, -C(S)N(R¹²)-, -N(R¹²)C(S)-, -SO₂N(R¹²)-, -N(R¹²)SO₂-, and -N(R¹²)C(O)N(R¹²)-, -N(R¹²)C(S)N(R¹²)-, -SO₂NHC(O)-, -SO₂N(R¹²)C(O)-, -C(O)NHSO₂-, -C(O)N(R¹²)SO₂- or M is a direct bond;

D is selected from C₁₋₆alkyl, C₁₋₆alkanoyl, C₁₋₆alkylsulphonyl, C₁₋₆alkoxycarbonyl, carbamoyl, N-(C₁₋₆alkyl)carbamoyl, N,N-(C₁₋₆alkyl)₂carbamoyl, benzoyl, (heterocyclic group)carbonyl, phenylsulphonyl, (heterocyclic group)sulphonyl, phenyl or a carbon linked heterocyclic group, and wherein any C₁₋₆alkyl group may be optionally substituted by one or more R⁹, and wherein any phenyl or heterocyclic group may be optionally substituted on a ring carbon by one or more groups selected from R⁸ and if a heterocyclic group contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from E;

E is selected from C₁₋₆alkyl, C₁₋₆alkanoyl, C₁₋₆alkylsulphonyl, C₁₋₆alkoxycarbonyl, carbamoyl, N-(C₁₋₆alkyl)carbamoyl, N,N-(C₁₋₆alkyl)₂carbamoyl, C₁₋₆alkoxyC₁₋₆alkanoyl, phenylC₁₋₆alkyl, benzoyl, phenylC₁₋₆alkanoyl, phenylC₁₋₆alkoxycarbonyl and phenylsulphonyl.

R⁹ is selected from hydroxy, amino, C₁₋₆alkylamino, (C₁₋₆alkyl)₂amino, carboxy, C₁₋₆alkoxy, carbamoyl, N-(C₁₋₆alkyl)carbamoyl, N-(C₁₋₆alkyl)₂carbamoyl, formyl, sulphamoyl, N-C₁₋₆alkylaminosulphonyl, N-(C₁₋₆alkyl)₂aminosulphonyl, C₁₋₆alkylsulphonylamino, C₁₋₆alkanoylamino, a heterocyclic group optionally substituted on a ring carbon by one or more R⁸ and if said heterocyclic group contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from D (wherein D is as defined hereinabove), phenyl optionally substituted by one or more R⁸, naphthyl optionally substituted by one or more R⁸, C₁₋₆alkylsulphanyl, C₁₋₆alkylsulphinyl and C₁₋₆alkylsulphonyl;

R¹⁰ is carboxy, carbamoyl, N-(C₁₋₆alkyl)carbamoyl, N-(C₁₋₆alkyl)₂carbamoyl, sulphamoyl, N-(C₁₋₆alkyl)aminosulphonyl, N-(C₁₋₆alkyl)₂aminosulphonyl, C₁₋₆alkylsulphanyl,

C₁₋₆alkylsulphinyl, C₁₋₆alkylsulphonyl, C₁₋₆alkoxycarbonyl, C₁₋₆alkanoylamino, a heterocyclic group optionally substituted on a ring carbon by one or more R⁸ and if said heterocyclic group contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from D (wherein D is as defined hereinabove), phenyl optionally substituted by one or more R⁸ or naphthyl optionally substituted by one or more R⁸;

R¹¹ is a heterocyclic group optionally substituted on a ring carbon by one or more R⁸ and if said heterocyclic group contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from D (wherein D is as defined hereinabove), phenyl optionally substituted by one or more R⁸, C₃₋₆cycloalkyl optionally substituted by one or more R⁸, or naphthyl optionally substituted by one or more R⁸;

R¹² is hydrogen or C₁₋₆alkyl optionally substituted with R¹³ with the proviso that R¹³ is not a substituent on the carbon attached to the nitrogen atom of M;


R¹³ is halo, hydroxy, amino, cyano, nitro, trifluoromethyl, trifluoromethoxy, C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, C₁₋₆alkylamino, (C₁₋₆alkyl)₂amino, C₁₋₆alkanoylamino, C₁₋₆alkanoyl(N-C₁₋₆alkyl)amino, C₁₋₆alkylsulphonylamino, C₁₋₆alkylsulphonyl(N-C₁₋₆alkyl)amino, thiol, C₁₋₆alkylsulphanyl, C₁₋₆alkylsulphinyl, C₁₋₆alkylsulphonyl, sulphamoyl, N-(C₁₋₆alkyl)aminosulphonyl, N-(C₁₋₆alkyl)₂aminosulphonyl, carboxy, carbamoyl, N-(C₁₋₆alkyl)carbamoyl, N-(C₁₋₆alkyl)₂carbamoyl, C₁₋₆alkoxycarbonyl, C₁₋₆alkanoyl or formyl;

n is 0-2;

with the proviso that where R¹ and R² are both as defined in (a), R³ and R⁴ are both as defined in (c), R⁵ is as defined in (e) and Ring X is phenyl, Y-Z must be -NHC(S)-; and pharmaceutically acceptable salts or *in vivo* hydrolysable esters thereof, and with the proviso that when R³ and R⁴ are both methyl, R⁵ is hydrogen, fluoro or chloro, Y-Z is ethynylene, X is phenyl and one of R¹ and R² is hydrogen and the other is pyrimidyl-NH-C(O)- or triazinyl-NH-C(O)- (wherein said triazine or pyrimidine is substituted by methyl, methoxy or dimethylamino) then the -SO₂NR¹R² moiety cannot be ortho to Y-Z; and provided said compound is not 4-(3-hydroxy-3-methyl-1-butyryl)-N-(3-methyl-2-pyridinyl)-benzenesulphonamide, N-{4-[N,N-bis-(*sec*-butyl)aminosulphonyl]phenyl}-2-hydroxy-2-methyl-3,3,3-trifluoropropanamide or N-{4-[N,N-bis-(*iso*-butyl)aminosulphonyl]phenyl}-2-hydroxy-2-methyl-3,3,3-trifluoropropanamide.

2. A compound of the formula (I) as claimed in claim 1 wherein R^3 and R^4 are independently C_k alkyl optionally substituted by from 1 to $2k+1$ atoms selected from fluoro and chloro wherein k is 1-3.
- 5 3. A compound of the formula (I) as claimed in claim 1 or 2 wherein one of R^3 and R^4 is methyl and the other is trifluoromethyl.
4. A compound of the formula (I) as claimed in claim 1, 2 or 3 wherein R^5 is selected from
10 halo, nitro, C_{1-4} alkyl, C_{1-4} alkoxy, C_{2-4} alkenyl, C_{2-4} alkynyl, hydroxy, hydrogen, amino, carboxy and sulphamoyl.
5. A compound of the formula (I) as claimed in claim 1, 2, 3 or 4 wherein R^5 is selected from fluoro and chloro.
- 15 6. A compound of the formula (I) as claimed in claim 1, 2, 3, 4 or 5 wherein X is phenyl.
7. A compound of the formula (I) as claimed in claim 1, 2, 3, 4, 5 or 6 wherein Y-Z is -NHC(O)-.
- 20 8. A compound of the formula (I) as claimed in claim 1, 2, 3, 4, 5, 6, or 7 wherein R^1 and R^2 are each independently selected from hydrogen, methyl, cyclopropyl, 4-hydroxycyclohexyl, 2-hydroxyethyl, 2-hydroxypropyl, 2,3-dihydroxypropyl, 4-(morpholinosulphonyl)phenyl, pyrid-3-yl, 2-carbamoylthien-3-yl, 2-chloropyrid-3-yl, 5-chloropyrid-2-yl, 5-methylpyrid-2-yl,
25 pyrimid-2-yl, 4,6-dimethylpyrimid-2-yl or 5,6-dimethylpyrazin-2-yl;
or R^1 and R^2 together with the nitrogen atom to which they are attached form 4-hydroxypiperidinyl or 1-(hydroxyacetyl)piperazin-4-yl.

9. A compound of formula (Ib):



(Ib)

wherein R^{1b} and R^{2b} are independently selected from:

- 5 i) hydrogen;
- ii) C₁₋₆alkyl or C₃₋₆cycloalkyl optionally substituted with one or more C₁₋₆alkyl, C₁₋₆alkoxy, C₂₋₆alkenyloxy, hydroxy, halo, cyano, C₁₋₆alkylsulphanyl, C₁₋₆alkylsulphinyl, C₁₋₆alkylsulphonyl, sulphamoyl, carboxy, C₁₋₆alkoxycarbonyl, amino, C₁₋₆alkylamino, (C₁₋₆alkyl)₂amino, carbamoyl, N-(C₁₋₆alkyl)carbamoyl, N-(C₁₋₆alkyl)₂carbamoyl, C₁₋₆alkanoylamino,
- 10 C₁₋₆alkanoyl(N-C₁₋₆alkyl)amino, C₁₋₆alkanoyl, C₂₋₆alkenyl, C₂₋₆alkynyl, N-(C₁₋₆alkyl)aminosulphonyl, hydroxymethyl, hydroxyacetyl or N-(C₁₋₆alkyl)₂aminosulphonyl;
- iii) a heterocyclic group selected from pyridyl, pyrimidyl, pyridazinyl or pyrazinyl, wherein said heterocyclic group is optionally substituted with one or more C₁₋₆alkyl, C₁₋₆alkoxy, C₂₋₆alkenyloxy, hydroxy, halo, cyano, C₁₋₆alkylsulphanyl, C₁₋₆alkylsulphinyl, C₁₋₆alkylsulphonyl,
- 15 sulphamoyl, carboxy, C₁₋₆alkoxycarbonyl, amino, C₁₋₆alkylamino, (C₁₋₆alkyl)₂amino, carbamoyl, N-(C₁₋₆alkyl)carbamoyl, N-(C₁₋₆alkyl)₂carbamoyl, C₁₋₆alkanoylamino, C₁₋₆alkanoyl(N-C₁₋₆alkyl)amino, trifluoromethyl, trifluoromethoxy, C₁₋₆alkanoyl, C₂₋₆alkenyl, C₂₋₆alkynyl, N-(C₁₋₆alkyl)aminosulphonyl, hydroxymethyl, hydroxyacetyl or N-(C₁₋₆alkyl)₂aminosulphonyl;
- 20 or R^{1b} and R^{2b} together with the nitrogen atom to which they are attached form piperidinyl or piperazinyl; wherein said piperidinyl and piperazinyl may be optionally substituted on a ring carbon by one or more groups selected from C₁₋₆alkyl, C₁₋₆alkoxy, C₂₋₆alkenyloxy, hydroxy, halo, cyano, C₁₋₆alkylsulphanyl, C₁₋₆alkylsulphinyl, C₁₋₆alkylsulphonyl, sulphamoyl, carboxy, C₁₋₆alkoxycarbonyl, amino, C₁₋₆alkylamino, (C₁₋₆alkyl)₂amino, carbamoyl,
- 25 N-(C₁₋₆alkyl)carbamoyl, N-(C₁₋₆alkyl)₂carbamoyl, C₁₋₆alkanoylamino, C₁₋₆alkanoyl(N-C₁₋₆alkyl)amino, C₁₋₆alkanoyl, C₂₋₆alkenyl, C₂₋₆alkynyl, N-(C₁₋₆alkyl)aminosulphonyl, hydroxymethyl, hydroxyacetyl or N-(C₁₋₆alkyl)₂aminosulphonyl;

and said piperazinyl may be optionally substituted on the ring nitrogen by a group selected from C₁₋₆alkanoyl, C₁₋₆alkylsulphonyl, C₁₋₆alkoxycarbonyl, carbamoyl, N-(C₁₋₆alkyl)carbamoyl and N,N-(C₁₋₆alkyl)₂carbamoyl; and wherein any C₁₋₆alkyl group may be optionally substituted by one or more groups selected from hydroxy, amino, C₁₋₆alkylamino, (C₁₋₆alkyl)₂amino, carboxy, C₁₋₆alkoxy, carbamoyl, N-(C₁₋₆alkyl)carbamoyl, N-(C₁₋₆alkyl)₂carbamoyl, sulphamoyl, N-C₁₋₆alkylaminosulphonyl, N-(C₁₋₆alkyl)₂aminosulphonyl, C₁₋₆alkylsulphonylamino, C₁₋₆alkanoylamino, C₁₋₆alkylsulphanyl, C₁₋₆alkylsulphinyl and C₁₋₆alkylsulphonyl. and pharmaceutically acceptable salts or *in vivo* hydrolysable esters thereof.


10. A pharmaceutical composition which comprises a compound of the formula (I) or (Ib) as claimed in claim 1 to 9 a pharmaceutically acceptable salt or an *in vivo* hydrolysable ester thereof, in association with a pharmaceutically acceptable excipient or carrier.

11. A compound of formula (I) or (Ib) as claimed in claim 1 to 9 and pharmaceutically acceptable salts or *in vivo* hydrolysable esters thereof for use as a medicament.

12. The use of a compound of the formula (I) or (Ib) as claimed in claim 1 to 9 or a pharmaceutically acceptable salt or an *in vivo* hydrolysable ester thereof in the manufacture of a medicament for use in the production of an elevation of PDH activity in a warm-blooded animal such as a human being.

13. A process for preparing a compound of formula (I) as claimed in claim 1 or a pharmaceutically acceptable salt or an *in vivo* hydrolysable ester thereof, which process comprises of:


(a) deprotecting a protected compound of formula (II)



(II)


where Pg is an alcohol protecting group;

(b) for a compound of formula (I) in which Y-Z is -NHC(O)-, by coupling an aniline of formula (III):



(III)

with an acid of formula (IV):




(IV)

wherein G is a hydroxyl group;

(c) by coupling an aniline of formula (III) with an activated acid derivative of formula (IV)


wherein G is a hydroxyl group optionally protected as an ester or ether;

(d) for a compound of formula (I) in which Y-Z is ethynylene, by reacting an alkyne of formula (V):



(V)

with a base, followed by treatment with a ketone of formula (VI):




(VI);

(e) for a compound of formula (I) in which Y-Z is *trans*-vinylene, by reducing a compound of


formula (I) in which Y-Z is ethynylene;

(f) for a compound of formula (I) in which Y-Z is *trans*-vinylene, by dehydration of a diol of formula (VII):



(VII);


(g) for a compound of formula (I) in which Y-Z is *trans*-vinylene, by base catalysed opening of an epoxide of formula (VIII):



(VIII);


(h) for a compound of formula (I) in which Y-Z is -NHCH₂-, by reducing a compound of formula (I) in which Y-Z is -NHC(O)-;

(i) for a compound of formula (I) in which Y-Z is -OCH₂-, -SCH₂- or -NHCH₂- by reacting an ethylene oxide of formula (IX):



(IX)


with a compound of formula (III) or a compound of formula (X):



15

(X) where J is -OH, -NH₂ or -SH;

(j) by reacting a compound of formula (XI):



(XI)

where K is a leaving atom or group, and in which Y-Z is OCH₂, SCH₂ or NHCH₂ or -NHC(O)-

with an amine of formula R^1R^2NH ;

(k) for a compound of formula (I) in which Y-Z is -NHC(S)-, by reacting a compound of formula (I) in which Y-Z is -NHC(O)- with a sulphur reagent;

and thereafter if necessary:

- 5 i) converting a compound of the formula (I) into another compound of the formula (I);
- ii) removing any protecting groups; or
- iii) forming a pharmaceutically acceptable salt or *in vivo* hydrolysable ester.